Signals from Gut Biome to Brain and Behavior

By Kenneth J. Bender, Pharm.D., M.A.

Studies of the interrelation between microbes of the gastrointestinal tract and general health are increasingly focused on bidirectional links between that microbiota and mental health, mood and behavior.

Research has suggested that while physiological and behavioral changes such as pharmacological intervention, diet and chronic stress can affect the microbiota configuration, gut microbiota have been shown to affect the development and functioning of the brain.

Origins of gut biome signaling research

Neuroscientists first noticed that mice bred to be "germ-free" (GF) showed a behavioral phenotype, with exaggerated stress-reactivity and within 5 years experiments reduced their anxiety-like behavior through exposure to selected microbes.

Physiological mechanisms

An array of processes is postulated to facilitate signaling between the gut microbiota and brain. A relatively direct link is suggested by the neurochemicals produced by microorganisms of the gut, which can bind to host receptors, including dopamine, GABA, serotonin and norepinephrine.

A more circuitous path involves immune and inflammatory responses with the microbiota provoking proinflammatory cytokines, which also affect brain function. Their effects in the hypothalamus include an increase in corticotrophin releasing hormone (CRH), with subsequent cascade of events through the hypothalamic-pituitaryadrenal axis (HPA).

Clinical applications

A "leaky gut" hypothesis proposes that increased permeability in the intestinal barrier could precede mucosal inflammation, or enable translocation of bacteria from the gut, prompting systemic inflammation.

It is consistent with stress being associated with alterations in the integrity of the intestinal barrier and the composition of microbiota, and posits that subsequent inflammatory response can mediate the expression of neuropsychiatric symptoms.

One study showed that the brain's signaling of the inflammasome, a multiprotein complex expressed to activate the pro-inflammatory cytokines, has been determined to affect anxiety- and depressive-like behaviors in mice as well as the composition of their gut microbiota.

In another study, GF mice receiving transplantations of fecal microbiota from human patients with major depression fared worse in stress testing than those receiving transplantations from healthy controls. The samples of the two groups had been confirmed to have distinctly different microbiota composition.

References

- Cryan JF, Dinan TG. Mind-altering microorganisms: The impact of the gut microbiota on brain and behaviour. Nat Rev Neurosci. 2012; 13:701-712.
- Kelly JR, Kennedy PJ, Cryan JF, et al. Breaking down the barriers: the gut microbiome, intestinal permeability and stress-related psychiatric disorders. *Front Cell Neurosci.* 2015, Oct 14; 9:392. doi:3389/fncel.2015.00392.
- Rogers GB, Keating DJ, Young RL, et al. From gut dysbiosis to altered brain function and mental illness: mechanisms and pathways. *Mol Psychiatry* 2016, April 19. doi:10.1038/mp.2016.50
- Sudo N, Chida Y, Aiba Y, et al. Postnatal microbial colonization programs the hypothalamic-pituitary-adrenal system for stress response in mice. J Physiol. 2004; 558: 263-275.
- Wong M-L, Inserra A, Lewis MD, et al. Inflammasome signaling affects anxiety- and depressive-like behaviour and gut microbiome composition. *Mol Psychiatry* 2016, April 19. doi:10.1038/mp.2916.46.
- Zheng P, Zeng B, Zhou C, et al. Gut microbiome remodeling induces depressive-like behaviours through a pathway mediated by the host's metabolism. *Mol Psychiatry* 2016, April 12. doi:10.1038/mp.2016.44

Reviewed by Jonas Wilson, Ing. Med.

Further Reading

- What is Neurology?
- <u>What is the Difference between Neurology and Neuroscience?</u>
- What is Neuroscience?
- What is Neurosurgery?
- Pediatric Neurology?
- What is Neurotoxicity?
- What is Neurooncology?
- <u>Neurological Infections</u>
- <u>Neurogenetics</u>

Last Updated: Jul 21, 2016

