

Uploaded to the VFC Website



This Document has been provided to you courtesy of Veterans-For-Change!

Feel free to pass to any veteran who might be able to use this information!

For thousands more files like this and hundreds of links to useful information, and hundreds of "Frequently Asked Questions, please go to:

Veterans-For-Change

If Veterans don't help Veterans, who will?

Note:

VFC is not liable for source information in this document, it is merely provided as a courtesy to our members & subscribers.



ETHYLBENZENE 147

4. CHEMICAL AND PHYSICAL INFORMATION

4.1 CHEMICAL IDENTITY

Ethylbenzene is an aromatic hydrocarbon that occurs naturally in petroleum and is a component of aviation and automotive fuels. It is used as a solvent and in the production of synthetic rubber and styrene. Information regarding the chemical identity of ethylbenzene is located in Table 4-1.

4.2 PHYSICAL AND CHEMICAL PROPERTIES

Ethylbenzene is a colorless liquid with an aromatic odor. Information regarding the physical and chemical properties of ethylbenzene is located in Table 4-2. Ethylbenzene is a flammable and combustible liquid. Vapors are heavier than air and may travel to a source of ignition and flash back. Liquid ethylbenzene floats on water and may travel to a source of ignition and spread fire. Combustion may produce irritants and toxic gases (HSDB 2009; NFPA 1994). Ethylbenzene may accumulate static electricity and will react with oxidizing materials (HSDB 2009; NFPA 1994).

4. CHEMICAL AND PHYSICAL INFORMATION

Table 4-1. Chemical Identity of Ethylbenzene

Characteristic	Information	Reference
Chemical name	Ethylbenzene	Budavari and O'Neil 1989
Synonym(s)	EB; ethyl benzene; benzene, ethyl-; ethylbenzol; phenylethane; etilbenzene; ethylbenzeen; aethylbenzo; etylobenzen	HSDB 2009
Registered trade name(s)	No data	
Chemical formula	C ₈ H ₁₀	Cannella 2007
Chemical structure	CH ₃ CH ₂	Cannella 2007
Identification numbers:		
CAS registry	100-41-4	Cannella 2007
NIOSH RTECS	NIOSH/DA0700000	NIOSH 2009; RTECS 2009
EPA hazardous waste	F003	HSDB 2009
OHM/TADS	7216709	HSDB 1995
DOT/UN/NA/IMDG shipping	UN 1175; IMDG 3.2	HSDB 2009
HSDB	84	HSDB 2009
NCI	NCI-C56393	HSDB 2009
STCC	49 091 63	HSDB 2009

CAS = Chemical Abstracts Service; DOT/UN/NA/IMDG = Department of Transportation/United Nations/North America/International Maritime Dangerous Goods Code; EPA = Environmental Protection Agency; HSDB = Hazardous Substances Data Bank; NCI = National Cancer Institute; NIOSH = National Institute for Occupational Safety and Health; OHM/TADS = Oil and Hazardous Materials/Technical Assistance Data System; RTECS = Registry of Toxic Effects of Chemical Substances; STCC = Standard Transport Commodity Code

Table 4-2. Physical and Chemical Properties of Ethylbenzene

Property	Information	Reference
Molecular weight	106.17	Lide 1994
Color	Colorless	Welch et al. 2005
Physical state	Liquid	Welch et al. 2005
Melting point	-94.975 °C	Cannella 2007
Boiling point	136.19 °C	Cannella 2007
Density at 20 °C/4 °C	0.8670	Welch et al. 2005
at 25 °C/25 °C	0.8671	Cannella 2007
Odor	Sweet, gasoline-like	CHRIS 1985
Odor threshold:		
Water	0.029 mg/L	Amoore and Hautala 1983
	0.140 mg/L	Rosen et al. 1963; Verschueren 1983
Air	2.3 ppm	Amoore and Hautala 1983
	2–2.6 mg/m ³	Verschueren 1983
Solubility:		
Water at 0 °C	197 mg/L	Polak and Lu 1973
at 15 °C	140 mg/L	Verschueren 1983
at 20 °C	152 mg/L	Verschueren 1983
at 25 °C	160 mg/L	Amoore and Hautala 1983
at 25 °C	177 mg/L	Polak and Lu 1973
at 25 °C	208 mg/L	Bohon and Claussen 1951
Organic solvents	Miscible with usual organic solvents	Budavari and O'Neil 1989
	Soluble in alcohol and ether	Lide 1994
Partition coefficients:		
Log K _{ow}	4.34	EPA 1982
	3.13	Yalkowsky and Valvani 1976
	3.15	Hansch et al. 1995
Log K _{oc}	2.22 (calculated)	Chiou et al. 1983
	2.38 (measured)	Hodson and Williams 1988
	2.40 (calculated)	Vowles and Mantoura 1987
Vapor pressure		
at 20 °C	7 mm Hg	Verschueren 1983
at 25 °C	1.27 kPa (9.53 mm Hg)	Mackay and Shiu 1981
at 25.9 °C	10 mm Hg	Sax and Lewis 1989
at 30 °C	12 mm Hg	Verschueren 1983

4. CHEMICAL AND PHYSICAL INFORMATION

Table 4-2. Physical and Chemical Properties of Ethylbenzene

Property	Information	Reference
Henry's law constant:		
at 20 °C	6.6x10 ⁻³ atm-m ³ /mol	EPA 1982
at 20 °C	8.7x10 ⁻³ atm-m ³ /mol	Lyman et al. 1982
at 25 °C	8.43x10 ⁻³ atm-m ³ /mol	Mackay et al. 1979
at 25 °C	7.9x10 ⁻³ atm-m ³ /mol	Mackay and Shiu 1981
Autoignition temperature	810 °F (432 °C)	Cannella 2007; NFPA 1994;
Flash point	15 °C	Cannella 2007
Flammability limits	0.8 (lower) vol% –6.7 (upper) vol%	NIOSH 2005
Conversion factors (25 °C, 1 atm)	1 mg/m ³ =0.230 ppm 1 ppm=4.34 mg/m ³	NIOSH 2005; Verschueren 1983