



Uploaded to VFC Website

~ November 2012 ~

This Document has been provided to you courtesy of Veterans-For-Change!

Feel free to pass to any veteran who might be able to use this information!

For thousands more files like this and hundreds of links to useful information, and hundreds of "Frequently Asked Questions, please go to:

[Veterans-For-Change](#)

*Veterans-For-Change is a 501(c)(3) Non-Profit Corporation
Tax ID #27-3820181*

If Veteran's don't help Veteran's, who will?

We appreciate all donations to continue to provide information and services to Veterans and their families.

https://www.paypal.com/cgi-bin/webscr?cmd=_s-xclick&hosted_button_id=WGT2M5UTB9A78

Note:

VFC is not liable for source information in this document, it is merely provided as a courtesy to our members.

Item ID Number 04245

Not Scanned

Author

Corporate Author

Report/Article Title Draft typescript; 227.71 Limiting Permissible Concentrations

Journal/Book Title

Year 0000

Month/Day

Color

Number of Images 2

Description Notes

227.71 Limiting Permissible Concentrations

A. 2,4-D and 2,4,5-T toxicity studies using freshwater and marine aquatic organisms were accomplished by the USAF Environmental Health Laboratory at Kelly AFB Texas (EHL/K). Also, sea water hydrolysis studies were performed. These studies together with an extensive literature review were used to characterize the expected effects and behavior of Orange herbicide components in aquatic systems. A thorough discussion of the findings ~~are~~ in Section F4, Part II of the Environmental Statement on the "Disposition of Orange Herbicide by Incineration".. In the discussion, the following generalizations were made:

(1) 2,4-D and 2,4,5-T are metabolized by fish.

(2) 2,4-D and 2,4,5-T do not undergo biomagnification through the food chain.

(3) Ester forms of 2,4-D and 2,4,5-T are usually more toxic to aquatic animals than are the parent acid forms so that hydrolysis is an important factor in reducing the toxicity of the esters.

(4) 2,4-D and 2,4,5-T and their N-butyl esters (NBE) rapidly disappear from sea water, via mechanisms of hydrolysis, metabolism and chemical interactions.

B. In acute toxicity studies by EHL(K), the 48 hr LC₅₀^{*} values in the shrimp (*Penaeus* sp.) were 5.6 ppm for 2,4-D NBE and 33 ppm for 2,4,5-T NBE. Oysters (*Crassostrea virginica*) were exposed to ~~apparent~~ concentrations of 2,4-D NBE ranging from 0.5 ppm to 85 ppm. The only acute effect observed was the death of one of the oysters (10%) in the highest concentration at 48 hours.

Using the above toxicity data ^{on shrimp (a sensitive species)} the limiting permissible concentrations are ~~at 0.01~~ ^{at 0.01}

^{of the LC₅₀} 0.056 ppm for 2,4-D NBE and 0.33 ppm for 2,4,5-T NBE. Because of the

highly toxic nature of 2,3,7,8 TCDD, limits are proposed to be 0.0001 ppm based on detection limits, rather than toxicity data.

*TL₅₀ and LC₅₀ (Tolerance Limit and Lethal Concentration) are concentration values statistically derived from the establishment of a dose-related response of experimental organisms to a toxicant. The LC is based on a measured response of death only. The TL is based on a count of unaffected organisms. The subscript number for both indicates the percent response expected for the calculated concentration. Therefore, in most cases, the TL₅₀ = LC₅₀ or the concentration in which 50% death is expected. Note that a more toxic chemical has a smaller LC₅₀.