



Uploaded to the VFC Website

▶▶ June 2014 ◀◀

This Document has been provided to you courtesy of Veterans-For-Change!

Feel free to pass to any veteran who might be able to use this information!

For thousands more files like this and hundreds of links to useful information, and hundreds of "Frequently Asked Questions, please go to:

[Veterans-For-Change](http://www.veteransforchange.org)

*Veterans-For-Change is a A 501(c)(3) Non-Profit Organization
Tax ID #27-3820181
CA Incorporation ID #3340400
CA Dept. of Charities ID #: CT-0190794*

If Veterans don't help Veterans, who will?

We appreciate all donations to continue to provide information and services to Veterans and their families.

https://www.paypal.com/cgi-bin/webscr?cmd=_s-xclick&hosted_button_id=WGT2M5UTB9A78

Note:

VFC is not liable for source information in this document, it is merely provided as a courtesy to our members & subscribers.



Item ID Number 05145



Not Scanned

Author

Corporate Author

Report/Article Title Handwritten Notes: Problems Encountered in
Determining Exposure to PCP

Journal/Book Title

Year 0000

Month/Day

Color



Number of Images 1

Description Notes

DMP - Barnes

~~Problems~~ Problems encountered in determining exposure to PCP.

Urine

Normality ~~is~~ must be established by region, sex, race, age, ^{weight} (urine volume, adipose level)

is further influenced

PCP - Geography - 80-85% positive detectable level > 5 ppb

Urine - 45
0.7-11 ppb mean 2 ppb

Serum

Humans are continuously exposed to

PCP in Adipose - $\bar{X} = 27$ ppb (17-52)

T.H. Shafiq, 1973

10(1): 57-63

Bulletin of Environmental Contamination & Toxicology

Florida

$\bar{X} = 140$ ppb

nd = 0.57 ppm

Ohe, T., 1979

22: 287-292

Bulletin

Bull. Environ. Contam. Toxicol.

Toxicol.

Lack Survey