



Uploaded to VFC Website

▶▶▶ July 2013 ◀◀◀

This Document has been provided to you courtesy of Veterans-For-Change!

Feel free to pass to any veteran who might be able to use this information!

For thousands more files like this and hundreds of links to useful information, and hundreds of "Frequently Asked Questions, please go to:

Veterans-For-Change

*Veterans-For-Change is a 501(c)(3) Non-Profit Corporation
Tax ID #27-3820181*

If Veteran's don't help Veteran's, who will?

We appreciate all donations to continue to provide information and services to Veterans and their families.

https://www.paypal.com/cgi-bin/webscr?cmd=_s-xclick&hosted_button_id=WGT2M5UTB9A78

Note:

VFC is not liable for source information in this document, it is merely provided as a courtesy to our members.



TOXICOLOGICAL PROFILE FOR COPPER

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Public Health Service
Agency for Toxic Substances and Disease Registry

September 2004

DISCLAIMER

The use of company or product name(s) is for identification only and does not imply endorsement by the Agency for Toxic Substances and Disease Registry.

UPDATE STATEMENT

A Toxicological Profile for Copper, Draft for Public Comment was released in September 2002. This edition supersedes any previously released draft or final profile.

Toxicological profiles are revised and republished as necessary. For information regarding the update status of previously released profiles, contact ATSDR at:

Agency for Toxic Substances and Disease Registry
Division of Toxicology/Toxicology Information Branch
1600 Clifton Road NE,
Mailstop F-32
Atlanta, Georgia 30333

FOREWORD

This toxicological profile is prepared in accordance with guidelines* developed by the Agency for Toxic Substances and Disease Registry (ATSDR) and the Environmental Protection Agency (EPA). The original guidelines were published in the *Federal Register* on April 17, 1987. Each profile will be revised and republished as necessary.

The ATSDR toxicological profile succinctly characterizes the toxicologic and adverse health effects information for the hazardous substance described therein. Each peer-reviewed profile identifies and reviews the key literature that describes a hazardous substance's toxicologic properties. Other pertinent literature is also presented, but is described in less detail than the key studies. The profile is not intended to be an exhaustive document; however, more comprehensive sources of specialty information are referenced.


The focus of the profiles is on health and toxicologic information; therefore, each toxicological profile begins with a public health statement that describes, in nontechnical language, a substance's relevant toxicological properties. Following the public health statement is information concerning levels of significant human exposure and, where known, significant health effects. The adequacy of information to determine a substance's health effects is described in a health effects summary. Data needs that are of significance to protection of public health are identified by ATSDR and EPA.

Each profile includes the following:

- (A) The examination, summary, and interpretation of available toxicologic information and epidemiologic evaluations on a hazardous substance to ascertain the levels of significant human exposure for the substance and the associated acute, subacute, and chronic health effects;
- (B) A determination of whether adequate information on the health effects of each substance is available or in the process of development to determine levels of exposure that present a significant risk to human health of acute, subacute, and chronic health effects; and
- (C) Where appropriate, identification of toxicologic testing needed to identify the types or levels of exposure that may present significant risk of adverse health effects in humans.

The principal audiences for the toxicological profiles are health professionals at the Federal, State, and local levels; interested private sector organizations and groups; and members of the public.

This profile reflects ATSDR's assessment of all relevant toxicologic testing and information that has been peer-reviewed. Staff of the Centers for Disease Control and Prevention and other Federal scientists have also reviewed the profile. In addition, this profile has been peer-reviewed by a nongovernmental panel and was made available for public review. Final responsibility for the contents and views expressed in this toxicological profile resides with ATSDR.


Julie Louise Gerberding, M.D., M.P.H.
Administrator
Agency for Toxic Substances and
Disease Registry

*Legislative Background

The toxicological profiles are developed in response to the Superfund Amendments and Reauthorization Act (SARA) of 1986 (Public Law 99-499) which amended the Comprehensive Environmental Response, Compensation, and Liability Act of 1980 (CERCLA or Superfund). This public law directed ATSDR to prepare toxicological profiles for hazardous substances most commonly found at facilities on the CERCLA National Priorities List and that pose the most significant potential threat to human health, as determined by ATSDR and the EPA. The availability of the revised priority list of 275 hazardous substances was announced in the *Federal Register* on November 7, 2003 (68 FR 63098). For prior versions of the list of substances, see *Federal Register* notices dated April 17, 1987 (52 FR 12866); October 20, 1988 (53 FR 41280); October 26, 1989 (54 FR 43619); October 17, 1990 (55 FR 42067); October 17, 1991 (56 FR 52166); October 28, 1992 (57 FR 48801); February 28, 1994 (59 FR 9486); April 29, 1996 (61 FR 18744); November 17, 1997 (62 FR 61332); October 21, 1999 (64 FR 56792) and October 25, 2001 (66 FR 54014). Section 104(i)(3) of CERCLA, as amended, directs the Administrator of ATSDR to prepare a toxicological profile for each substance on the list.

QUICK REFERENCE FOR HEALTH CARE PROVIDERS

Toxicological Profiles are a unique compilation of toxicological information on a given hazardous substance. Each profile reflects a comprehensive and extensive evaluation, summary, and interpretation of available toxicologic and epidemiologic information on a substance. Health care providers treating patients potentially exposed to hazardous substances will find the following information helpful for fast answers to often-asked questions.

Primary Chapters/Sections of Interest

Chapter 1: Public Health Statement: The Public Health Statement can be a useful tool for educating patients about possible exposure to a hazardous substance. It explains a substance's relevant toxicologic properties in a nontechnical, question-and-answer format, and it includes a review of the general health effects observed following exposure.

Chapter 2: Relevance to Public Health: The Relevance to Public Health Section evaluates, interprets, and assesses the significance of toxicity data to human health.

Chapter 3: Health Effects: Specific health effects of a given hazardous compound are reported by type of health effect (death, systemic, immunologic, reproductive), by route of exposure, and by length of exposure (acute, intermediate, and chronic). In addition, both human and animal studies are reported in this section.

NOTE: Not all health effects reported in this section are necessarily observed in the clinical setting. Please refer to the Public Health Statement to identify general health effects observed following exposure.

Pediatrics: Four new sections have been added to each Toxicological Profile to address child health issues:

Section 1.6	How Can (Chemical X) Affect Children?
Section 1.7	How Can Families Reduce the Risk of Exposure to (Chemical X)?
Section 3.7	Children's Susceptibility
Section 6.6	Exposures of Children

Other Sections of Interest:

Section 3.8	Biomarkers of Exposure and Effect
Section 3.11	Methods for Reducing Toxic Effects

ATSDR Information Center

Phone: 1-888-42-ATSDR or (404) 498-0110 **Fax:** (770) 488-4178

E-mail: atsdric@cdc.gov **Internet:** <http://www.atsdr.cdc.gov>

The following additional material can be ordered through the ATSDR Information Center:

Case Studies in Environmental Medicine: Taking an Exposure History—The importance of taking an exposure history and how to conduct one are described, and an example of a thorough exposure history is provided. Other case studies of interest include *Reproductive and Developmental Hazards; Skin Lesions and Environmental Exposures; Cholinesterase-Inhibiting Pesticide Toxicity*; and numerous chemical-specific case studies.

Managing Hazardous Materials Incidents is a three-volume set of recommendations for on-scene (prehospital) and hospital medical management of patients exposed during a hazardous materials incident. Volumes I and II are planning guides to assist first responders and hospital emergency department personnel in planning for incidents that involve hazardous materials. Volume III—*Medical Management Guidelines for Acute Chemical Exposures*—is a guide for health care professionals treating patients exposed to hazardous materials.

Fact Sheets (ToxFAQs) provide answers to frequently asked questions about toxic substances.

Other Agencies and Organizations

The National Center for Environmental Health (NCEH) focuses on preventing or controlling disease, injury, and disability related to the interactions between people and their environment outside the workplace. Contact: NCEH, Mailstop F-29, 4770 Buford Highway, NE, Atlanta, GA 30341-3724 • Phone: 770-488-7000 • FAX: 770-488-7015.

The National Institute for Occupational Safety and Health (NIOSH) conducts research on occupational diseases and injuries, responds to requests for assistance by investigating problems of health and safety in the workplace, recommends standards to the Occupational Safety and Health Administration (OSHA) and the Mine Safety and Health Administration (MSHA), and trains professionals in occupational safety and health. Contact: NIOSH, 200 Independence Avenue, SW, Washington, DC 20201 • Phone: 800-356-4674 or NIOSH Technical Information Branch, Robert A. Taft Laboratory, Mailstop C-19, 4676 Columbia Parkway, Cincinnati, OH 45226-1998 • Phone: 800-35-NIOSH.

The National Institute of Environmental Health Sciences (NIEHS) is the principal federal agency for biomedical research on the effects of chemical, physical, and biologic environmental agents on human health and well-being. Contact: NIEHS, PO Box 12233, 104 T.W. Alexander Drive, Research Triangle Park, NC 27709 • Phone: 919-541-3212.

Referrals

The Association of Occupational and Environmental Clinics (AOEC) has developed a network of clinics in the United States to provide expertise in occupational and environmental issues. Contact: AOEC, 1010 Vermont Avenue, NW, #513, Washington, DC 20005 • Phone: 202-347-4976 • FAX: 202-347-4950 • e-mail: AOEC@AOEC.ORG • Web Page: <http://www.aoec.org/>.

The American College of Occupational and Environmental Medicine (ACOEM) is an association of physicians and other health care providers specializing in the field of occupational and environmental medicine. Contact: ACOEM, 55 West Seegers Road, Arlington Heights, IL 60005 • Phone: 847-818-1800 • FAX: 847-818-9266.

CONTRIBUTORS

CHEMICAL MANAGER(S)/AUTHOR(S):

Alfred Dorsey, DVM
ATSDR, Division of Toxicology, Atlanta, GA

Lisa Ingerman, Ph.D.
Steven Swarts, Ph.D.
Syracuse Research Corporation, Syracuse, NY

THE PROFILE HAS UNDERGONE THE FOLLOWING ATSDR INTERNAL REVIEWS:

1. Health Effects Review. The Health Effects Review Committee examines the health effects chapter of each profile for consistency and accuracy in interpreting health effects and classifying end points.
2. Minimal Risk Level Review. The Minimal Risk Level Workgroup considers issues relevant to substance-specific Minimal Risk Levels (MRLs), reviews the health effects database of each profile, and makes recommendations for derivation of MRLs.
3. Data Needs Review. The Research Implementation Branch reviews data needs sections to assure consistency across profiles and adherence to instructions in the Guidance.

PEER REVIEW

A peer review panel was assembled for copper (September 2002 profile). The panel consisted of the following members:

1. Dr. Jonathan H. Freedman, Center for Environmental Genomes, Duke University, Durham, North Carolina;
2. Dr. Paul Mushak, PB Associates, Durham, North Carolina; and
3. Dr. Robert B. Ruckner, School of Medicine, Department of Nutrition, University of California at Davis, Davis, California.
4. Dr. Edward Massaro, U.S. Environmental Protection Agency, Reproductive Toxicology Facility, Durham, North Carolina

These experts collectively have knowledge of copper's physical and chemical properties, toxicokinetics, key health end points, mechanisms of action, human and animal exposure, and quantification of risk to humans. All reviewers were selected in conformity with the conditions for peer review specified in Section 104(I)(13) of the Comprehensive Environmental Response, Compensation, and Liability Act, as amended.

Scientists from the Agency for Toxic Substances and Disease Registry (ATSDR) have reviewed the peer reviewers' comments and determined which comments will be included in the profile. A listing of the peer reviewers' comments not incorporated in the profile, with a brief explanation of the rationale for their exclusion, exists as part of the administrative record for this compound. A list of databases reviewed and a list of unpublished documents cited are also included in the administrative record.

The citation of the peer review panel should not be understood to imply its approval of the profile's final content. The responsibility for the content of this profile lies with the ATSDR.

CONTENTS

DISCLAIMER	ii
UPDATE STATEMENT	iii
FOREWORD	v
QUICK REFERENCE FOR HEALTH CARE PROVIDERS.....	vii
CONTRIBUTORS	ix
PEER REVIEW	xi
CONTENTS.....	xiii
LIST OF FIGURES	xvii
LIST OF TABLES.....	xix
1. PUBLIC HEALTH STATEMENT.....	1
1.1 WHAT IS COPPER?.....	1
1.2 WHAT HAPPENS TO COPPER WHEN IT ENTERS THE ENVIRONMENT?.....	2
1.3 HOW MIGHT I BE EXPOSED TO COPPER?	3
1.4 HOW CAN COPPER ENTER AND LEAVE MY BODY?	6
1.5 HOW CAN COPPER AFFECT MY HEALTH?	6
1.6 HOW CAN COPPER AFFECT CHILDREN?	7
1.7 HOW CAN FAMILIES REDUCE THE RISK OF EXPOSURE TO COPPER?	7
1.8 IS THERE A MEDICAL TEST TO DETERMINE WHETHER I HAVE BEEN EXPOSED TO COPPER?	8
1.9 WHAT RECOMMENDATIONS HAS THE FEDERAL GOVERNMENT MADE TO PROTECT HUMAN HEALTH?.....	8
1.10 WHERE CAN I GET MORE INFORMATION?.....	9
2. RELEVANCE TO PUBLIC HEALTH	11
2.1 BACKGROUND AND ENVIRONMENTAL EXPOSURES TO COPPER IN THE UNITED STATES.....	11
2.2 SUMMARY OF HEALTH EFFECTS	12
2.3 MINIMAL RISK LEVELS (MRLs)	15
3. HEALTH EFFECTS	21
3.1 INTRODUCTION.....	21
3.2 DISCUSSION OF HEALTH EFFECTS BY ROUTE OF EXPOSURE	21
3.2.1 Inhalation Exposure	22
3.2.1.1 Death	22
3.2.1.2 Systemic Effects	22
3.2.1.3 Immunological and Lymphoreticular Effects	25
3.2.1.4 Neurological Effects	25
3.2.1.5 Reproductive Effects	25
3.2.1.6 Developmental Effects	31
3.2.1.7 Cancer.....	31
3.2.2 Oral Exposure.....	32
3.2.2.1 Death	32
3.2.2.2 Systemic Effects	32
3.2.2.3 Immunological and Lymphoreticular Effects	62
3.2.2.4 Neurological Effects	63
3.2.2.5 Reproductive Effects	63

3.2.2.6	Developmental Effects	63
3.2.2.7	Cancer	64
3.2.3	Dermal Exposure	64
3.2.3.1	Death	64
3.2.3.2	Systemic Effects	65
3.2.3.3	Immunological and Lymphoreticular Effects	65
3.2.3.4	Neurological Effects	65
3.2.3.5	Reproductive Effects	65
3.2.3.6	Developmental Effects	65
3.2.3.7	Cancer	65
3.2.4	Other Routes of Exposure	66
3.3	GENOTOXICITY	66
3.4	TOXICOKINETICS	68
3.4.1	Absorption	68
3.4.1.1	Inhalation Exposure	68
3.4.1.2	Oral Exposure	70
3.4.1.3	Dermal Exposure	72
3.4.2	Distribution	72
3.4.2.1	Inhalation Exposure	72
3.4.2.2	Oral Exposure	72
3.4.2.3	Dermal Exposure	73
3.4.3	Metabolism	73
3.4.4	Elimination and Excretion	74
3.4.4.1	Inhalation Exposure	74
3.4.4.2	Oral Exposure	74
3.4.4.3	Dermal Exposure	74
3.4.4.4	Other Routes of Exposure	74
3.4.5	Physiologically Based Pharmacokinetic (PBPK)/Pharmacodynamic (PD) Models	74
3.5	MECHANISMS OF ACTION	76
3.5.1	Pharmacokinetic Mechanisms	76
3.5.2	Mechanisms of Toxicity	78
3.5.3	Animal-to-Human Extrapolations	79
3.6	TOXICITIES MEDIATED THROUGH THE NEUROENDOCRINE AXIS	80
3.7	CHILDREN'S SUSCEPTIBILITY	81
3.8	BIOMARKERS OF EXPOSURE AND EFFECT	85
3.8.1	Biomarkers Used to Identify or Quantify Exposure to Copper	86
3.8.2	Biomarkers Used to Characterize Effects Caused by Copper	87
3.9	INTERACTIONS WITH OTHER CHEMICALS	88
3.10	POPULATIONS THAT ARE UNUSUALLY SUSCEPTIBLE	89
3.11	METHODS FOR REDUCING TOXIC EFFECTS	91
3.11.1	Reducing Peak Absorption Following Exposure	91
3.11.2	Reducing Body Burden	92
3.11.3	Interfering with the Mechanism of Action for Toxic Effects	93
3.12	ADEQUACY OF THE DATABASE	93
3.12.1	Existing Information on Health Effects of Copper	93
3.12.2	Identification of Data Needs	95
3.12.3	Ongoing Studies	102
4.	CHEMICAL AND PHYSICAL INFORMATION	105
4.1	CHEMICAL IDENTITY	105
4.2	PHYSICAL AND CHEMICAL PROPERTIES	105

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL	111
5.1 PRODUCTION	111
5.2 IMPORT/EXPORT	117
5.3 USE.....	117
5.4 DISPOSAL	118
6. POTENTIAL FOR HUMAN EXPOSURE	121
6.1 OVERVIEW.....	121
6.2 RELEASES TO THE ENVIRONMENT	124
6.2.1 Air	129
6.2.2 Water.....	135
6.2.3 Soil	141
6.3 ENVIRONMENTAL FATE.....	142
6.3.1 Transport and Partitioning.....	143
6.3.1.1 Ambient Air.....	143
6.3.1.2 Ambient Waters.....	144
6.3.1.3 Ambient Soils	146
6.3.1.4 Bioconcentration and Biomagnification	149
6.3.2 Transformation and Degradation	151
6.3.2.1 Air.....	151
6.3.2.2 Water	151
6.3.2.3 Sediment and Soil.....	155
6.3.2.4 Other Media.....	158
6.4 LEVELS MONITORED OR ESTIMATED IN THE ENVIRONMENT	158
6.4.1 Air	158
6.4.2 Water.....	163
6.4.3 Sediment and Soil	167
6.4.4 Other Environmental Media	171
6.5 GENERAL POPULATION AND OCCUPATIONAL EXPOSURE	182
6.6 EXPOSURES OF CHILDREN	183
6.7 POPULATIONS WITH POTENTIALLY HIGH EXPOSURES	184
6.8 ADEQUACY OF THE DATABASE.....	185
6.8.1 Identification of Data Needs	185
6.8.2 Ongoing Studies	189
7. ANALYTICAL METHODS	193
7.1 BIOLOGICAL MATERIALS.....	193
7.2 ENVIRONMENTAL SAMPLES.....	193
7.3 ADEQUACY OF THE DATABASE.....	197
7.3.1 Identification of Data Needs	198
7.3.2 Ongoing Studies	198
8. REGULATIONS AND ADVISORIES	201
9. REFERENCES	207
10. GLOSSARY	267

APPENDICES

A. ATSDR MINIMAL RISK LEVELS AND WORKSHEETSA-1

B. USER'S GUIDE B-1

C. ACRONYMS, ABBREVIATIONS, AND SYMBOLS..... C-1

D. INDEXD-1

LIST OF FIGURES

3-1. Levels of Significant Exposure to Copper – Inhalation.....	28
3-2. Levels of Significant Exposure to Copper – Oral.....	50
3-3. Conceptual Representation of a Physiologically Based Pharmacokinetic (PBPK) Model for a Hypothetical Chemical Substance	77
3-4. Existing Information on Health Effects of Copper.....	94
6-1. Frequency of NPL Sites with Copper Contamination	122

LIST OF TABLES

3-1. Levels of Significant Exposure to Copper – Inhalation.....	26
3-2. Levels of Significant Exposure to Copper – Oral.....	28
3-3. Genotoxicity of Copper <i>In Vivo</i>	67
3-4. Genotoxicity of Copper <i>In Vitro</i>	69
3-5. Ongoing Studies on Copper.....	103
4-1. Chemical Identity of Copper	106
4-2. Chemical Identity of Copper Sulfate	107
4-3. Physical and Chemical Properties of Copper and Copper Sulfate.....	109
5-1. Facilities that Produce, Process, or Use Copper	112
5-2. Facilities that Produce, Process, or Use Copper Compounds.....	114
6-1. Releases to the Environment from Facilities that Produce, Process, or Use Copper.....	125
6-2. Releases to the Environment from Facilities that Produce, Process, or Use Copper Compounds ...	127
6-3. Summary of Copper Concentrations in Environmental Media	130
6-4. Global Emissions of Copper from Natural Sources ($\times 10^6$ kg/year).....	131
6-5. Global Emissions of Copper from Anthropogenic Sources ($\times 10^6$ kg/year).....	132
6-6. Concentrations of Copper in Particulate Matter (<10 μm) Generated from Various Sources.....	134
6-7. Concentrations of Copper in Water	139
6-8. Concentrations of Copper in Air.....	160
6-9. Particle Size Distributions and Total Copper Concentrations in Dust Collected at Four Mine Waste Pump Sites in Butte, Montana	162
6-10. Concentration of Copper in Surface Soils of the United States (in ppm-Dry Weight [dw], Equivalent to mg/kg-dw).....	168
6-11. Geometric Means of Selected Soil Elements and Associated Soil Parameters in U.S. Surface Soils by Land Resource Regions.....	169
6-12. Copper Content of Selected Foods (mg/kg)	173
6-13. Copper Content of Human Tissues and Body Fluids	178

6-14. Agricultural Sources of Copper Contamination in Soils	181
6-15. Ongoing Studies on Environmental Fate and the Potential for Human Exposure to Copper	190
7-1. Analytical Methods for Determining Copper in Biological Materials	194
7-2. Analytical Methods for Determining Copper in Environmental Samples	195
8-1. Regulations and Guidelines Applicable to Copper	202

1. PUBLIC HEALTH STATEMENT

This public health statement tells you about copper and the effects of exposure to it.

The Environmental Protection Agency (EPA) identifies the most serious hazardous waste sites in the nation. These sites are then placed on the National Priorities List (NPL) and are targeted for long-term federal clean-up activities. Copper has been found in at least 906 of the 1,647 current or former NPL sites. Although the total number of NPL sites evaluated for this substance is not known, the possibility exists that the number of sites at which copper is found may increase in the future as more sites are evaluated. This information is important because these sites may be sources of exposure and exposure to this substance may harm you.

When a substance is released either from a large area, such as an industrial plant, or from a container, such as a drum or bottle, it enters the environment. Such a release does not always lead to exposure. You can be exposed to a substance only when you come in contact with it and your body is able to absorb it. You may be exposed by breathing, eating, or drinking the substance, or by skin contact.

If you are exposed to copper, many factors will determine whether you will be harmed. These factors include the dose (how much), the duration (how long), and how you come in contact with it. You must also consider any other chemicals you are exposed to and your age, sex and other genetic traits, diet, family traits, lifestyle, and state of health, including pregnancy and developmental stage of embryo/fetus.

1.1 WHAT IS COPPER?

Copper is a reddish metal that occurs naturally in rock, soil, water, sediment, and, at low levels, air. Its average concentration in the earth's crust is about 50 parts copper per million parts soil (ppm) or, stated another way, 50 grams of copper per 1,000,000 grams of soil (1.8 ounces or 0.11 pounds of copper per 2,200 pounds of soil). Copper also occurs naturally in all plants and animals. It is an essential element for all known living organisms including humans and other

1. PUBLIC HEALTH STATEMENT

animals at low levels of intake. At much higher levels, toxic effects can occur. The term copper in this profile not only refers to copper metal, but also to compounds of copper that may be in the environment.

Metallic copper can be easily molded or shaped. The reddish color of this element is most commonly seen in the U.S. penny, electrical wiring, and some water pipes. It is also found in many mixtures of metals, called alloys, such as brass and bronze. Many compounds (substances formed by joining two or more chemicals) of copper exist. These include naturally occurring minerals as well as manufactured chemicals. The most commonly used compound of copper is copper sulfate. Many copper compounds can be recognized by their blue-green color.

Copper is extensively mined and processed in the United States and is primarily used as the metal or alloy in the manufacture of wire, sheet metal, pipe, and other metal products. Copper compounds are most commonly used in agriculture to treat plant diseases, like mildew, or for water treatment and as preservatives for wood, leather, and fabrics. For more information on the properties and uses of copper, please see Chapters 4 and 5.

1.2 WHAT HAPPENS TO COPPER WHEN IT ENTERS THE ENVIRONMENT?

Copper can enter the environment through releases from the mining of copper and other metals, and from factories that make or use copper metal or copper compounds. Copper can also enter the environment through waste dumps, domestic waste water, combustion of fossil fuels and wastes, wood production, phosphate fertilizer production, and natural sources (for example, windblown dust, from native soils, volcanoes, decaying vegetation, forest fires, and sea spray). Therefore, copper is widespread in the environment. About 1,400,000,000 pounds (640,000,000,000 grams) of copper were released into the environment by industries in 2000. Copper is often found near mines, smelters, industrial settings, landfills, and waste disposal sites.

When copper is released into soil, it can become strongly attached to the organic material and other components (e.g., clay, sand, etc.) in the top layers of soil and may not move very far when it is released. When copper and copper compounds are released into water, the copper that

1. PUBLIC HEALTH STATEMENT

dissolves can be carried in surface waters either in the form of copper compounds or as free copper or, more likely, copper bound to particles suspended in the water. Even though copper binds strongly to suspended particles and sediments, there is evidence to suggest that some water-soluble copper compounds do enter groundwater. Copper that enters water eventually collects in the sediments of rivers, lakes, and estuaries. Copper is carried on particles emitted from smelters and ore processing plants, and is then carried back to earth through gravity or in rain or snow. Copper is also carried into the air on windblown metallurgical dust. Indoor release of copper comes mainly from combustion processes (for example, kerosene heaters).

Elemental copper does not break down in the environment. Copper can be found in plants and animals, and at high concentrations in filter feeders such as mussels and oysters. Copper is also found in a range of concentrations in many foods and beverages that we eat and drink, including drinking water. You will find additional information on the fate of copper in the environment in Chapters 5 and 6.

1.3 HOW MIGHT I BE EXPOSED TO COPPER?

Copper is common in the environment. You may be exposed to copper by breathing air, drinking water, eating food, and by skin contact with soil, water and other copper-containing substances. Most copper compounds found in air, water, sediment, soil and rock are strongly attached to dust and dirt or imbedded in minerals. You can take copper into your body upon ingestion of water or soil that contains copper or by inhalation of copper-containing dust. Some copper in the environment is less tightly bound to soil or particles in water and may be soluble enough in water to be taken up by plants and animals. In the general population, soluble copper compounds (those that dissolve in water), which are most commonly used in agriculture, are more likely to threaten your health. When soluble copper compounds are released into lakes and rivers, they generally become attached to particles in the water within approximately 1 day. This could lessen your exposure to copper in water, depending on how strongly the copper is bound to the particles and how much of the particles settle into lake and river sediments. However, fine particles have an enormous surface area and can remain suspended for prolonged periods of

1. PUBLIC HEALTH STATEMENT

time. Therefore, at high fine particle concentrations, both exposure and uptake can be considerable even under conditions of tight copper binding to the suspended particulates.

The concentration of copper in air ranges from a few nanograms (1 nanogram equals 1/1,000,000,000 of a gram or 4/100,000,000,000 of an ounce) in a cubic meter of air (ng/m^3) to about 200 ng/m^3 . A cubic meter (m^3) is approximately 25% larger than a cubic yard. Near smelters, which process copper ore into metal, concentrations may reach 5,000 ng/m^3 . You may breathe high levels of copper-containing dust if you live or work near copper mines or processing facilities.

You may be exposed to levels of soluble copper in your drinking water that are above the acceptable drinking water standard of 1,300 parts copper per billion parts of water (ppb), especially if your water is corrosive and you have copper plumbing and brass water fixtures. The average concentration of copper in tap water ranges from 20 to 75 ppb. However, many households have copper concentrations of over 1,000 ppb. That is more than 1 milligram per liter of water. This is because copper is dissolved from copper pipes and brass faucets when the water sits in the pipes overnight. After the water is allowed to run for 15–30 seconds, the concentration of copper in the water decreases below the acceptable drinking water standard.

The concentration of copper in lakes and rivers ranges from 0.5 to 1,000 ppb with an average concentration of 10 ppb. The average copper concentration in groundwater (5 ppb) is similar to that in lakes and rivers; however, monitoring data indicate that some groundwater contains levels of copper (up to 2,783 ppb) that are well above the standard of 1,300 ppb for drinking water. This copper is generally bound to particles in the water. Lakes and reservoirs recently treated with copper compounds to control algae or receive cooling water from a power plant can have high concentrations of dissolved copper. Once in natural water, much of this copper soon attaches to particles or convert to other forms that can settle into sediments. This can limit exposure to copper unless the sediments are stirred; for example, by the resuspension and swallowing of sediments by swimmers in recreational waters.

1. PUBLIC HEALTH STATEMENT

Garden products containing copper that are used to control certain plant diseases are also a potential source of exposure through contact with skin or if they are accidentally swallowed. For example, you can find copper compounds in some fungicides.

Soil generally contains between 2 and 250 ppm copper, although concentrations close to 17,000 ppm have been found near copper and brass production facilities. High concentrations of copper may be found in soil because dust from these industries settles out of the air, or wastes from mining and other copper industries are disposed of on the soil. Another common source of copper in soil results from spreading sludge from sewage treatment plants. This copper generally stays strongly attached to the surface layer of soil. You may be exposed to this copper by skin contact. Children may also be exposed to this copper by hand to mouth contact and eating the contaminated dirt and dust.

Food naturally contains copper. You eat and drink about 1 milligram (1/1,000 of a gram or 4/100,000 ounces) of copper every day.

While some hazardous waste sites on the NPL contain high levels of copper, we do not always know how high it is above natural levels. We also do not know what form it is in at most of these sites. However, evidence suggests that most copper at these sites is strongly attached to soil.

You may be exposed to copper in the workplace. If you work in the industry of mining copper or processing the ore, you are exposed to copper by breathing copper-containing dust or by skin contact. If you grind or weld copper metal, you may breathe high levels of copper dust and fumes. Occupational exposure to forms of copper that are soluble or not strongly attached to dust or dirt would most commonly occur in agriculture, water treatment, and industries such as electroplating, where soluble copper compounds are used. Exposure to copper in air in the workplace is regulated and is set to be below concentrations that can be harmful to you.

For more information on the potential for exposure to copper, please refer to Chapter 6.

1. PUBLIC HEALTH STATEMENT

1.4 HOW CAN COPPER ENTER AND LEAVE MY BODY?

Copper can enter your body when you drink water or eat food, soil, or other substances that contain copper. Copper can also enter your body if you breathe air or dust containing copper. Copper may enter the lungs of workers exposed to copper dust or fumes.

Copper rapidly enters the bloodstream and is distributed throughout the body after you eat or drink it. Certain substances in foods eaten with copper can affect the amount of copper that enters the bloodstream from the gastrointestinal tract. Your body is very good at blocking high levels of copper from entering the bloodstream. We do not know how much copper enters the body through the lungs or skin. Copper then leaves your body in feces and urine, mostly in feces. It takes several days for copper to leave your body. Generally, the amount of copper in your body remains constant (the amount that enters your body equals the amount that leaves). More information on how copper enters and leaves the body is presented in Chapter 3.

1.5 HOW CAN COPPER AFFECT MY HEALTH?

Scientists use many tests to protect the public from harmful effects of toxic chemicals and to find ways for treating persons who have been harmed.

One way to learn whether a chemical will harm people is to determine how the body absorbs, uses, and releases the chemical. For some chemicals, animal testing may be necessary. Animal testing may also help identify health effects such as cancer or birth defects. Without laboratory animals, scientists would lose a basic method for getting information needed to make wise decisions that protect public health. Scientists have the responsibility to treat research animals with care and compassion. Scientists must comply with strict animal care guidelines because laws today protect the welfare of research animals.

Copper is essential for good health. However, exposure to higher doses can be harmful. Long-term exposure to copper dust can irritate your nose, mouth, and eyes, and cause headaches, dizziness, nausea, and diarrhea. If you drink water that contains higher than normal levels of

1. PUBLIC HEALTH STATEMENT

copper, you may experience nausea, vomiting, stomach cramps, or diarrhea. Intentionally high intakes of copper can cause liver and kidney damage and even death. We do not know if copper can cause cancer in humans. EPA does not classify copper as a human carcinogen because there are no adequate human or animal cancer studies.

More detailed information on the health effects of copper in animals and humans can be found in Chapter 3.

1.6 HOW CAN COPPER AFFECT CHILDREN?

This section discusses potential health effects in humans from exposures during the period from conception to maturity at 18 years of age.

Exposure to high levels of copper will result in the same types of effects in children and adults. We do not know if these effects would occur at the same dose level in children and adults. Studies in animals suggest that children may have more severe effects than adults; we do not know if this would also be true in humans. There is a very small percentage of infants and children who are unusually sensitive to copper. We do not know if copper can cause birth defects or other developmental effects in humans. Studies in animals suggest that ingestion of high levels of copper may cause a decrease in fetal growth.

1.7 HOW CAN FAMILIES REDUCE THE RISK OF EXPOSURE TO COPPER?

The greatest potential source of copper exposure is through drinking water, especially in water that is first drawn in the morning after sitting in copper piping and brass faucets overnight. To reduce copper in drinking water, run the water for at least 15–30 seconds before using it. Additionally, if there is concern about the concentration of copper in drinking water exceeding the minimum value of 1,300 ppb, families should have their water tested.

1. PUBLIC HEALTH STATEMENT

If your doctor finds that you have been exposed to substantial amounts of copper, ask whether your children might also have been exposed. Your doctor might need to ask your state health department to investigate.

1.8 IS THERE A MEDICAL TEST TO DETERMINE WHETHER I HAVE BEEN EXPOSED TO COPPER?

Copper is normally found in all tissues of the body, blood, urine, feces, hair, and nails. High levels of copper in the blood, urine, hair, and nails can show that you have been exposed to higher than normal levels of copper. Tests to measure copper levels in the body are not usually available at a doctor's office because they require special equipment, but the doctor can send samples to a specialty laboratory. Although these tests can show that you have been exposed to higher than normal copper levels, they can not be used to predict the extent of exposure or potential health effects. More detailed information on the measurement of copper is provided in Chapters 3 and 7.

1.9 WHAT RECOMMENDATIONS HAS THE FEDERAL GOVERNMENT MADE TO PROTECT HUMAN HEALTH?

The federal government develops regulations and recommendations to protect public health. Regulations *can* be enforced by law. The EPA, the Occupational Safety and Health Administration (OSHA), and the Food and Drug Administration (FDA) are some federal agencies that develop regulations for toxic substances. Recommendations provide valuable guidelines to protect public health, but *cannot* be enforced by law. The Agency for Toxic Substances and Disease Registry (ATSDR) and the National Institute for Occupational Safety and Health (NIOSH) are two federal organizations that develop recommendations for toxic substances.

Regulations and recommendations can be expressed as “not-to-exceed” levels, that is, levels of a toxic substance in air, water, soil, or food that do not exceed a critical value that is usually based on levels that affect animals; they are then adjusted to levels that will help protect humans.

1. PUBLIC HEALTH STATEMENT

Sometimes these not-to-exceed levels differ among federal organizations because they used different exposure times (an 8-hour workday or a 24-hour day), different animal studies, or other factors.

Recommendations and regulations are also updated periodically as more information becomes available. For the most current information, check with the federal agency or organization that provides it. Some regulations and recommendations for copper include the following:

The EPA has determined that drinking water should not contain more than 1.3 mg copper per liter of water (1.3 mg/L). The EPA has also developed regulations on the amount of copper that industry is allowed to release.

The OSHA has set a limit of 0.1 milligrams/cubic meter (mg/m³) for copper fumes (vapor generated from heating copper) and 1.0 mg/m³ for copper dusts (fine metallic copper particles) and mists (aerosols of soluble copper) in workroom air to protect workers during an 8-hour work shift (40-hour workweek).

The Food and Nutrition Board of the Institute of Medicine has developed recommended dietary allowances (RDAs) of 340 micrograms (µg) of copper per day for children aged 1–3 years, 440 µg/day for children aged 4–8 years, 700 µg/day for children aged 9–13 years, 890 µg/day for children aged 14–18 years, and 900 µg/day for adults. This provides enough copper to maintain health. Further information on regulations and guidelines pertaining to copper is provided in Chapter 8.

1.10 WHERE CAN I GET MORE INFORMATION?

If you have any more questions or concerns, please contact your community or state health or environmental quality department, or contact ATSDR at the address and phone number below.

1. PUBLIC HEALTH STATEMENT

ATSDR can also tell you the location of occupational and environmental health clinics. These clinics specialize in recognizing, evaluating, and treating illnesses that result from exposure to hazardous substances.

Toxicological profiles are also available on-line at www.atsdr.cdc.gov and on CD-ROM. You may request a copy of the ATSDR ToxProfiles™ CD-ROM by calling the toll-free information and technical assistance number at 1-888-42ATSDR (1-888-422-8737), by e-mail at atsdric@cdc.gov, or by writing to:

Agency for Toxic Substances and Disease Registry
Division of Toxicology
1600 Clifton Road NE
Mailstop F-32
Atlanta, GA 30333
Fax: 1-770-488-4178

Organizations for-profit may request copies of final Toxicological Profiles from the following:

National Technical Information Service (NTIS)
5285 Port Royal Road
Springfield, VA 22161
Phone: 1-800-553-6847 or 1-703-605-6000
Web site: <http://www.ntis.gov/>

2. RELEVANCE TO PUBLIC HEALTH

2.1 BACKGROUND AND ENVIRONMENTAL EXPOSURES TO COPPER IN THE UNITED STATES

Copper is a metallic element that occurs naturally as the free metal, or associated with other elements in compounds that comprise various minerals. Most copper compounds occur in +1 Cu(I) and +2 Cu(II) valence states. Copper is primarily used as a metal or an alloy (e.g., brass, bronze, gun metal). Copper sulfate is used as a fungicide, algicide, and nutritional supplement. Copper particulates are released into the atmosphere by windblown dust; volcanic eruptions; and anthropogenic sources, primarily copper smelters and ore processing facilities. Copper particles in the atmosphere will settle out or be removed by precipitation, but can be resuspended into the atmosphere in the form of dust. The mean concentration of copper in ambient air in the United States ranges from 5 to 200 ng/m³. Copper is released into waterways by natural weathering of soil and rocks, disturbances of soil, or anthropogenic sources (e.g., effluent from sewage treatment plants). Copper concentrations in drinking water vary widely as a result of variations in pH and hardness of the water supply; the levels range from a few ppbs to 10 ppm. The mean concentration of copper in soil in the United States ranges from 5 to 70 mg/kg. The estimated daily intake of copper from food is 1.0–1.3 mg/day for adults (0.014–0.019 mg/kg/day).

The general population is exposed to copper through inhalation, consumption of food and water, and dermal contact with air, water, and soil that contains copper. The primary source of copper intake is the diet; however, the amount of copper in the diet usually does not exceed the average dietary requirements (RDAs) for copper. Drinking water is the primary source of excess copper. Populations living near sources of copper emissions, such as copper smelters and refineries and workers in these and other industries may also be exposed to high levels of copper in dust by inhalation. Copper concentrations in soils near copper emission sources could be sufficiently high to result in significantly high intakes of copper in young children who ingest soil. For example, copper concentrations of 2,480–6,912 ppm have been measured near copper smelters. These levels of copper in soils would result in the intake of 0.74–2.1 mg copper per day in a child ingesting 300 mg of soil. Copper has been identified in at least 906 of the 1,647 hazardous waste sites that have been proposed for inclusion on the EPA NPL.

2. RELEVANCE TO PUBLIC HEALTH

2.2 SUMMARY OF HEALTH EFFECTS

Copper is an essential nutrient that is incorporated into a number of metalloenzymes involved in hemoglobin formation, drug/xenobiotic metabolism, carbohydrate metabolism, catecholamine biosynthesis, the cross-linking of collagen, elastin, and hair keratin, and the antioxidant defense mechanism. Copper-dependent enzymes, such as cytochrome c oxidase, superoxide dismutase, ferroxidases, monoamine oxidase, and dopamine β -monooxygenase, function to reduce activated oxygen species or molecular oxygen. Symptoms associated with copper deficiency in humans include normocytic, hypochromic anemia, leukopenia, and osteoporosis; copper deficiency is rarely observed in the U.S. general population. In the United States, the median intake of copper from food is 0.93–1.3 mg/day for adults (0.013–0.019 mg Cu/kg body weight/day using a 70-kg reference body weight). A recommended dietary allowance (RDA) of 0.9 mg/day (0.013 mg/kg/day) has recently been established.

Copper is readily absorbed from the stomach and small intestine. After nutritional requirements are met, there are several mechanisms that prevent copper overload. Excess copper absorbed into gastrointestinal mucosal cells induces the synthesis of and binds to the metal binding protein metallothionein. This bound copper is excreted when the cell is sloughed off. Copper that eludes binding to intestinal metallothionein is transported to the liver. It is stored in the liver bound to liver metallothionein, from which it is ultimately released into bile and excreted in the feces. Although copper homeostasis plays an important role in the prevention of copper toxicity, exposure to excessive levels of copper can result in a number of adverse health effects including liver and kidney damage, anemia, immunotoxicity, and developmental toxicity. Many of these effects are consistent with oxidative damage to membranes or macromolecules. Copper can bind to the sulfhydryl groups of several enzymes, such as glucose-6-phosphatase and glutathione reductase, thus interfering with their protection of cells from free radical damage.

One of the most commonly reported adverse health effect of copper is gastrointestinal distress. Nausea, vomiting, and/or abdominal pain have been reported, usually occurring shortly after drinking a copper sulfate solution, beverages that were stored in a copper or untinned brass container, or first draw water (water that sat in the pipe overnight). The observed effects are not usually persistent and gastrointestinal effects have not been linked with other health effects. Animal studies have also reported gastrointestinal effects (hyperplasia of forestomach mucosa) following ingestion of copper sulfate in the diet. Copper is also irritating to the respiratory tract. Coughing, sneezing, runny nose, pulmonary fibrosis, and increased vascularity of the nasal mucosa have been reported in workers exposed to copper dust.

2. RELEVANCE TO PUBLIC HEALTH

The liver is also a sensitive target of toxicity. Liver damage (necrosis, fibrosis, abnormal biomarkers of liver damage) have been reported in individuals ingesting lethal doses of copper sulfate. Liver effects have also been observed in individuals diagnosed with Wilson's disease, Indian childhood cirrhosis, or idiopathic copper toxicosis (which includes Tyrollean infantile cirrhosis). These syndromes are genetic disorders that result in an accumulation of copper in the liver; the latter two syndromes are associated with excessive copper exposure. Inflammation, necrosis, and altered serum markers of liver damage have been observed in rats fed diets with copper sulfate levels that are at least 100 times higher than the nutritional requirement. Damage to the proximal convoluted tubules of the kidney has also been observed in rats. The liver and kidney effects usually occur at similar dose levels; however, the latency period for the kidney effects is longer than for the liver effects.

There is some evidence from animal studies to suggest that exposure to airborne copper or high levels of copper in drinking water can damage the immune system. Impaired cell-mediated and humoral-mediated immune function have been observed in mice. Studies in rats, mice, and mink suggest that exposure to high levels of copper in the diet can result in decreased embryo and fetal growth.

The carcinogenicity of copper has not been adequately studied. An increase in cancer risk has been found among copper smelters; however, the increased risk has been attributed to concomitant exposure to arsenic. Increased lung and stomach cancer risks have also been found in copper miners. However, a high occurrence of smoking and exposure to radioactivity, silica, iron, and arsenic obscure the association of copper exposure with carcinogenesis. Animal studies have not found increased cancer risks in orally exposed rats or mice. The IARC has classified the pesticide, copper 8-hydroxyquinoline, in Group 3, unclassifiable as to carcinogenicity in humans and EPA has classified copper in Group D, not classifiable as to human carcinogenicity

A more detailed discussion of the critical targets of copper toxicity, the gastrointestinal tract and the liver, follows.

Gastrointestinal Effects. The available human and animal data suggest that the gastrointestinal tract is a sensitive target of toxicity. There are numerous reports of nausea, vomiting, and/or abdominal pain in humans ingesting beverages contaminated with copper or water containing copper sulfate. These symptoms typically occur shortly after ingestion and are not persistent. The results of three single exposure studies suggest that the threshold for gastrointestinal symptoms is between 4 and 6 ppm, which is equivalent to doses of 0.11 mg/kg and 0.017–0.018 mg Cu/kg. Nausea, vomiting, and/or abdominal

2. RELEVANCE TO PUBLIC HEALTH

pain also appear to be the most sensitive end point following repeated exposure to copper in drinking water. These symptoms were reported by adults drinking water containing ≥ 3 ppm copper as copper sulfate (0.0731 mg Cu/kg/day) for 1–2 weeks or 4 ppm copper as copper sulfate (0.091 mg Cu/kg/day) for 2 months. Similar gastrointestinal effects were observed in adults ingesting copper oxide in drinking water. Although gastrointestinal irritation may play a role in the observed gastrointestinal effects, data from ferrets and monkeys suggest that vagal afferent fibers and 5-HT₃ and 5-HT₄ receptors are involved in copper-induced emesis.

Hepatic Effects. In humans, copper-induced hepatic damage is dependent on several factors including genetics, age, and copper intake. Liver damage is rarely reported in adults; the few reported cases of liver damage (centrilobular necrosis, jaundice, and increased aspartate aminotransferase activity) have been associated with intentional ingestion of a lethal dose of copper sulfate. In infants and children, reported liver effects are usually manifested in one of three syndromes: Wilson's disease, Indian childhood cirrhosis, and idiopathic copper toxicosis. Wilson's disease is an autosomal recessive genetic disorder associated with impaired copper metabolism. Dietary exposure to higher than normal levels of copper does not appear to be necessary for the manifestation of liver damage. Some heterozygous carriers of Wilson's disease also have elevated hepatic levels of copper and increased urinary excretion, although adverse health effects have not been reported in these individuals. There is evidence that Indian childhood cirrhosis and idiopathic copper toxicosis are also caused by a genetic defect that is transmitted in an autosomal recessive mode. However, unlike Wilson's disease, manifestation of the disease is associated with exposure to unusually high levels of dietary copper from milk stored in copper or brass containers or from drinking water. The clinical age of onset is usually between 6 months and 5 years, and the observed liver effects include pericellular fibrosis, abnormal biochemical markers of liver damage (e.g., increased serum aspartate aminotransferase and alkaline phosphatase activities and serum bilirubin levels), and very high levels of copper in the liver. In general, the potential hepatotoxicity of copper has not been extensively investigated in healthy humans. No effect levels of 0.14–0.17 and 0.315 mg Cu/kg/day for liver effects in adults and infants (3–12 months of age), respectively, had been reported in intermediate-duration studies (2–9 months); these studies used serum chemistry biomarkers (e.g., alanine aminotransferase, aspartate aminotransferase) to assess liver damage. Two community survey studies also found no evidence of liver damage in infants living in households with 0.8 ppm copper in drinking water. The results of the three studies involving infants should be interpreted cautiously due to the high drop out rate, small number of subjects examined for possible liver damage, and the dismissal of anomalous findings as secondary to infection rather than possibly indicative of copper toxicity.

2. RELEVANCE TO PUBLIC HEALTH

Adverse liver effects have been observed in rats exposed to dietary copper levels that were more than 100 times higher than the nutritional requirement. The liver effects included inflammation, necrosis, and abnormal serum chemistry markers of liver damage. Rats appear to develop a tolerance to copper doses of 180–<550 mg Cu/kg/day. Tolerance is defined as “the ability to endure the continued or increasing administration of a toxicant and the capacity to exhibit less response to a test dose than previous.” As the levels of hepatic copper increase, so does the severity of the damage until peak copper levels are reached. After about 3–5 weeks of exposure, the copper levels begin to decline and are maintained at a steady level for the remainder of the exposure period. When the hepatic levels decline, regeneration of hepatic tissue is observed, and continued exposure or exposure to higher doses does not result in more tissue damage. The decline in hepatic copper levels and regeneration of damaged tissue occurs early at higher doses. At doses >550 mg Cu/kg/day, the liver becomes permanently overloaded and chronic hepatitis develops.

2.3 MINIMAL RISK LEVELS (MRLs)

Estimates of exposure levels posing minimal risk to humans (Minimal Risk Levels or MRLs) have been made for copper. An MRL is defined as an estimate of daily human exposure to a substance that is likely to be without an appreciable risk of adverse effects (noncarcinogenic) over a specified duration of exposure. MRLs are derived when reliable and sufficient data exist to identify the target organ(s) of effect or the most sensitive health effect(s) for a specific duration within a given route of exposure. MRLs are based on noncancerous health effects only and do not consider carcinogenic effects. MRLs can be derived for acute, intermediate, and chronic duration exposures for inhalation and oral routes. Appropriate methodology does not exist to develop MRLs for dermal exposure.

Although methods have been established to derive these levels (Barnes and Dourson 1988; EPA 1990), uncertainties are associated with these techniques. Furthermore, ATSDR acknowledges additional uncertainties inherent in the application of the procedures to derive less than lifetime MRLs. As an example, acute inhalation MRLs may not be protective for health effects that are delayed in development or are acquired following repeated acute insults, such as hypersensitivity reactions, asthma, or chronic bronchitis. As these kinds of health effects data become available and methods to assess levels of significant human exposure improve, these MRLs will be revised.

A User's Guide has been provided at the end of this profile (see Appendix B). This guide should aid in the interpretation of the tables and figures for Levels of Significant Exposure and the MRLs.

2. RELEVANCE TO PUBLIC HEALTH

Inhalation MRLs

The available data on the toxicity of inhaled copper were considered inadequate for derivation of acute-, intermediate-, or chronic-duration inhalation MRLs. Data on the inhaled toxicity of copper in humans following acute-duration exposure are limited to a report of workers developing metal fume fever while cutting brass pipe with an electric cutting tool in a poorly ventilated area (Armstrong et al. 1983); exposure levels were not reported. Respiratory effects and impaired immune function have been observed in mice following a single 3-hour exposure to 3.3 mg Cu/m³ as copper sulfate or repeated exposure (3 hours/day, 5 days/week for 1–2 weeks) to 0.12–0.13 mg Cu/m³ as copper sulfate (Drummond et al. 1986). The Drummond et al. (1986) study was not selected as the basis of an acute-duration inhalation MRL because a small number of animals was tested (four per group) and a limited number of end points (respiratory tract and immune function) were examined. Intermediate-duration data are limited to studies by Johansson et al. (1983, 1984), which did not find any histological alterations in the lungs or functional or morphological alterations in alveolar macrophages of rabbits exposed to copper chloride. As with the acute-duration data, the limited number of end points examined precludes deriving an intermediate-duration inhalation MRL. The chronic-duration database for copper consists of two occupational exposure studies reporting respiratory (Askergren and Mellgren 1975; Suciú et al. 1981) and gastrointestinal (Suciú et al. 1981) irritation, hepatic effects (Suciú et al. 1981), and possible neurological and reproductive effects (Suciú et al. 1981). Chronic-duration inhalation MRLs cannot be derived from these studies due to poor exposure characterization and/or lack of controls.

Oral MRLs

- An MRL of 0.01 mg/kg/day has been derived for acute-duration oral exposure (1–14 days) to copper.

The available human and animal acute-duration studies strongly suggest that the gastrointestinal tract is the most sensitive target of copper toxicity. Numerous studies and case reports have reported nausea, vomiting, and/or abdominal pain in humans immediately following ingestion of copper-contaminated water or other beverages (Araya et al. 2001, 2003a, 2003b, 2003c; Chuttani et al. 1965; Gotteland et al. 2001; Knobloch et al. 1994; Nicholas and Brist 1968; Olivares et al. 2001; Pizarro et al. 1999, 2001; Spitalny et al. 1984). In human studies involving a single exposure to copper following an overnight fast, adverse gastrointestinal effects (nausea, vomiting, abdominal pain, and/or diarrhea) have been observed at doses of 0.011–0.03 mg Cu/kg (Araya et al. 2001, 2003a, 2003c; Gotteland et al. 2001; Olivares et al. 2001). Under these experimental conditions, the apparent threshold appears to fall between 0.011 and

2. RELEVANCE TO PUBLIC HEALTH

0.017 mg Cu/kg (Araya et al. 2001, 2003a; Olivares et al. 2001). Slightly higher thresholds for gastrointestinal symptoms were observed in two acute-duration repeated exposure studies in which subjects used a copper-containing water as their primary source of drinking water for 1 or 2 weeks (Pizarro et al. 1999, 2001). In the 2-week study, 60 women were given copper sulfate containing water to be used for drinking and cooking purposes. No significant alterations in serum biomarkers of liver damage (alanine aminotransferase, aspartate aminotransferase, γ -glutamyl transferase) were observed in the subjects at the end of the study. An increased occurrence of nausea, vomiting, and/or abdominal pain was observed when the women were exposed to 3 ppm copper as copper sulfate (0.0731 mg Cu/kg/day) (Pizarro et al. 1999); no significant increases in the incidence of gastrointestinal symptoms were noted at 1 ppm (0.0272 mg Cu/kg/day). Nausea, vomiting, and/or abdominal pain were also reported by women ingesting water containing 5 ppm (0.096 mg Cu/kg/day) as copper sulfate or copper oxide for 1 week (Pizarro et al. 2001). Animal studies support the identification of the gastrointestinal tract as a sensitive target of toxicity. Hyperplasia of the forestomach mucosa was observed in rats exposed to 44 mg Cu/kg/day as copper sulfate in the diet (NTP 1993) and in mice exposed to 197 mg Cu/kg/day as copper sulfate in the diet (NTP 1993). At higher doses, liver and kidney damage have been observed (Haywood 1980; Haywood and Comerford 1980; Haywood et al. 1985b; NTP 1993).

The Pizarro et al. (1999) 2-week study was selected as the basis of the acute-duration oral MRL for copper. This study identified no-observed-adverse-effect level (NOAEL) and lowest-observed-adverse-effect level (LOAEL) values of 0.0272 and 0.0731 mg Cu/kg/day for increases in the incidence of nausea, vomiting, and/or abdominal pain. Although the LOAEL values identified in the single exposure studies (Araya et al. 2001, 2003; Olivares et al. 2001) are slightly lower than the NOAEL identified in the Pizarro et al. (1999) study, the Pizarro et al. (1999) study was selected as the critical study because it is a longer-duration study and it more closely mimics an exposure scenario of a population drinking copper-contaminated drinking water. The NOAEL was divided by an uncertainty factor of 3 (to account for human variability) to yield an acute-duration oral MRL of 0.01 mg Cu/kg/day. The observed gastrointestinal effects were probably due to direct contact; thus, only a partial uncertainty factor of 3 was used to account for human variability because toxicokinetic differences among individuals should not affect sensitivity. The acute-duration MRL is intended to protect against the health effects associated with exposure to copper-contaminated drinking water; it assumes that the affected population will have a normal intake of copper from the diet.

2. RELEVANCE TO PUBLIC HEALTH

- An MRL of 0.01 mg/kg/day has been derived for intermediate-duration oral exposure (15–365 days) to copper.

There are limited data on the intermediate-duration toxicity of copper in humans. Araya et al. (2003b) exposed groups of 327–355 adults to <0.01 (control group), 2, 4, or 6 ppm copper sulfate in water for 2 months. The subjects prepared the copper sulfate solution to be used at home by mixing a stock copper sulfate solution with tap water; this solution was used for drinking water and preparing beverages and soups. Exposure to copper sulfate resulted in increases in the occurrence of gastrointestinal symptoms; the incidence was significantly higher than controls at 6 ppm when the data were analyzed using the chi-square test with Bonferroni correction and at 4 ppm when the Bonferroni correction was not used. Only one test was used to assess whether exposure to copper results in adverse gastrointestinal effects (reported symptoms); thus, the Bonferroni correction is not needed for this end point. Therefore, the 4 ppm concentration is identified as the LOAEL and the 2 ppm concentration as the NOAEL. The study authors reported copper intakes for 48–49 subjects per group who provided blood samples; no information on selection criteria were provided. The copper intakes were 0, 0.042, 0.091, and 0.17 mg Cu/kg/day for the control, 2, 4, and 6 ppm groups, respectively. The dietary intake of copper was not measured in this study; however, Araya et al. (2003b) noted that copper intake found in a survey of other community residents was 0.9 mg Cu/day. No significant alterations in copper status or liver function (as assessed by serum alanine aminotransferase, aspartate aminotransferase, and γ -glutamyl transferase activities) were observed in a subset of subjects from each group. In a study by Pratt et al. (1985), a group of seven adults were administered 10 mg Cu/day (0.14 mg Cu/kg/day) as copper gluconate in a capsule for 12 weeks. No significant alterations in serum markers of liver damage (cholesterol and triglyceride levels and serum aspartate aminotransferase, alkaline phosphatase, γ -glutamyl transferase, and lactate dehydrogenase activities) were found. Similarly, no alterations in total bilirubin or serum alanine aminotransferase, aspartate aminotransferase, or γ -glutamyl transferase activities were observed in infants exposed to 0.315 mg Cu/kg/day for 9 months (Olivares et al. 1998). Zietz et al. (2003a, 2003b) also did not find evidence of liver damage in infants living in households with water concentrations of 0.8 ppm and higher. The Pratt et al. (1985), Olivares et al. (1998), and Zietz et al. (2003a, 2003b) studies did not report significant alterations in the occurrence of gastrointestinal disturbances and the study design did not include symptoms questionnaires, although the high dropout rate observed in the Olivares et al. (1998) study may have been related to gastrointestinal effects. Severe liver damage (pericellular fibrosis and increased serum aminotransferase and alkaline phosphatase activities) has been observed in children with a genetic susceptibility to high levels of copper in the liver. The liver was a critical target of toxicity in rats exposed to very high levels of copper in diet (greater than 100 times the nutritional requirement). Inflammation, necrosis, and increased alanine and aspartate aminotransferases activities have been

2. RELEVANCE TO PUBLIC HEALTH

reported in rats at exposure levels of 16 mg Cu/kg/day as copper sulfate in the diet (Haywood 1980, 1985; Haywood and Comerford 1980; Haywood and Loughran 1985; Haywood et al. 1985a; NTP 1993). No liver effects were observed at 8 mg Cu/kg/day (NTP 1993). Histological alterations in stomach, indicative of irritation (hyperplasia of the squamous mucosa on the limiting ridge separating the forestomach from the glandular stomach), have also been observed in rats and mice exposed to 33 or 267 mg Cu/kg/day, respectively, as copper sulfate in the diet for 13 weeks (NTP 1993).

An intermediate-duration oral MRL of 0.01 mg Cu/kg/day was derived for copper based on gastrointestinal effects using the data from the Araya et al. (2003b) study. This study identified NOAEL and LOAEL values of 0.042 and 0.091 mg Cu/kg/day, respectively; these copper doses were in excess of normal dietary intake. The NOAEL was divided by an uncertainty factor of 3 (to account for human variability) to yield an intermediate-duration oral MRL of 0.01 mg Cu/kg/day. As with the acute-duration MRL, the intermediate-duration MRL is intended to protect against exposure to excess copper in drinking water and assumes a normal copper dietary intake.

The database on the chronic oral toxicity of copper is inadequate for derivation of a MRL. Massie and Aiello (1984) reported a 15% decrease in the lifespan in mice exposed to 4.2 mg Cu/kg/day as copper gluconate in drinking water.

3. HEALTH EFFECTS

3.1 INTRODUCTION

The primary purpose of this chapter is to provide public health officials, physicians, toxicologists, and other interested individuals and groups with an overall perspective on the toxicology of copper. It contains descriptions and evaluations of toxicological studies and epidemiological investigations and provides conclusions, where possible, on the relevance of toxicity and toxicokinetic data to public health.

A glossary and list of acronyms, abbreviations, and symbols can be found at the end of this profile.

3.2 DISCUSSION OF HEALTH EFFECTS BY ROUTE OF EXPOSURE

To help public health professionals and others address the needs of persons living or working near hazardous waste sites, the information in this section is organized first by route of exposure (inhalation, oral, and dermal) and then by health effect (death, systemic, immunological, neurological, reproductive, developmental, genotoxic, and carcinogenic effects). These data are discussed in terms of three exposure periods: acute (14 days or less), intermediate (15–364 days), and chronic (365 days or more).

Levels of significant exposure for each route and duration are presented in tables and illustrated in figures. The points in the figures showing no-observed-adverse-effect levels (NOAELs) or lowest-observed-adverse-effect levels (LOAELs) reflect the actual doses (levels of exposure) used in the studies. LOAELs have been classified into "less serious" or "serious" effects. "Serious" effects are those that evoke failure in a biological system and can lead to morbidity or mortality (e.g., acute respiratory distress or death). "Less serious" effects are those that are not expected to cause significant dysfunction or death, or those whose significance to the organism is not entirely clear. ATSDR acknowledges that a considerable amount of judgment may be required in establishing whether an end point should be classified as a NOAEL, "less serious" LOAEL, or "serious" LOAEL, and that in some cases, there will be insufficient data to decide whether the effect is indicative of significant dysfunction. However, the Agency has established guidelines and policies that are used to classify these end points. ATSDR believes that there is sufficient merit in this approach to warrant an attempt at distinguishing between "less serious" and "serious" effects. The distinction between "less serious" effects and "serious" effects is considered to be important because it helps the users of the profiles to identify levels of exposure at which

3. HEALTH EFFECTS

major health effects start to appear. LOAELs or NOAELs should also help in determining whether or not the effects vary with dose and/or duration, and place into perspective the possible significance of these effects to human health.

The significance of the exposure levels shown in the Levels of Significant Exposure (LSE) tables and figures may differ depending on the user's perspective. Public health officials and others concerned with appropriate actions to take at hazardous waste sites may want information on levels of exposure associated with more subtle effects in humans or animals (LOAELs) or exposure levels below which no adverse effects (NOAELs) have been observed. Estimates of levels posing minimal risk to humans (Minimal Risk Levels or MRLs) may be of interest to health professionals and citizens alike.

A User's Guide has been provided at the end of this profile (see Appendix B). This guide should aid in the interpretation of the tables and figures for Levels of Significant Exposure and the MRLs.

3.2.1 Inhalation Exposure

3.2.1.1 Death

No studies were located regarding death of humans or animals following inhalation exposure to copper.

3.2.1.2 Systemic Effects

No studies were located regarding cardiovascular, musculoskeletal, renal, dermal, or body weight effects in humans or animals following inhalation exposure to copper.

Respiratory, gastrointestinal, hematological, hepatic, endocrine, and ocular effects were observed in humans. Respiratory effects have also been observed in animals exposed to copper sulfate aerosols.

Respiratory Effects. In humans, copper is a respiratory irritant. Workers exposed to copper dust report a number of symptoms that are suggestive of respiratory irritation, including coughing, sneezing, thoracic pain, and runny nose (Askergren and Mellgren 1975; Suciú et al. 1981). In the Suciú et al. (1981) study of 75–100 workers involved in sieving copper, lung radiographs revealed linear pulmonary fibrosis, and in some cases, nodulation. During the first year of operation, the workers were exposed to 434 mg Cu/m³; the exposure levels declined each year, and by year 3, the levels were 111 mg Cu/m³. In

3. HEALTH EFFECTS

sheet metal workers exposed to patina dust (copper-hydroxide-nitrate, copper-hydroxide-sulfate, copper silicate, copper oxide), 6 of the 11 examined workers had increased vascularity and superficial epistatic vessels in the nasal mucosa (Askergren and Mellgren 1975); no exposure levels were reported.

Copper is considered the etiologic agent in the occupational disease referred to as “vineyard sprayer’s lung”. This disease, which is observed in vineyard workers spraying an antimildew agent containing 1–2.5% copper sulfate neutralized with hydrated lime, was first described in humans by Cortez Pimentel and Marques (1969). In most cases, published information on this disease comes from case reports (Cortez Pimentel and Marques 1969; Cortez Pimentel and Menezes 1975; Stark 1981; Villar 1974; Villar and Nogueira 1980) with no concentration-response information. Common findings (obtained by alveolar lavage and biopsy) include intraalveolar desquamation of macrophages, formation of histiocytic and noncaseating granulomas containing inclusions of copper, and healing of lesions in the form of fibrohyaline nodules, very similar to those found in silicosis (Cortez Pimentel and Marques 1969; Plamenac et al. 1985). Higher incidences of abnormal columnar cells, squamous metaplasia without atypia, copper containing macrophages, eosinophilia, and respiratory spirals were found in the sputa of smoking and nonsmoking vineyard sprayers, as compared to rural workers from the same geographic region who did not work in the vineyards (Plamenac et al. 1985).

The potential of copper to induce respiratory effects has been tested in mice, hamsters, and rabbits. Decreased cilia beating was observed in Syrian-Golden hamsters exposed to 3.3 mg Cu/m³ as copper sulfate for 3 hours (Drummond et al. 1986); this effect was not observed in similarly exposed CD-1 mice. Repeated exposure resulted in alveolar thickening in CD-1 mice exposed to 0.12 mg Cu/m³ as copper sulfate for 3 hours/day, 5 days/week for 1–2 weeks (Drummond et al. 1986); the severity of the effect increased with the duration of exposure. In rabbits (strain not reported) exposed to 0.6 mg Cu/m³ as copper chloride for 6 hours/day, 5 days/week for 4–6 weeks, the only histological alteration in the lungs was a slight increase in alveolar type II cell volume density (Johansson et al. 1984); this effect was not considered adverse. No functional or morphological alterations were observed in the alveolar macrophages of similarly exposed rabbits (Johansson et al. 1983).

Gastrointestinal Effects. In workers involved in grinding and sieving copper dust, anorexia, nausea, and occasional diarrhea were reported (Suciu et al. 1981); exposure levels ranged from 111 to 434 mg Cu/m³ over a 3-year period. It is likely that the observed gastrointestinal effects were due to oral exposure to copper. Ingestion probably resulted from mucocilliary clearance of copper particles deposited in the nasopharyngeal and tracheobronchial regions of the respiratory tract.

3. HEALTH EFFECTS

No studies were located regarding gastrointestinal effects in animals following inhalation exposure to copper.

Hematological Effects. Decreased hemoglobin and erythrocyte levels have been observed in workers exposed to airborne copper levels of 0.64–1.05 mg/m³. Results of hair analysis reveal that the workers were also exposed to iron, lead, and cadmium (Finelli et al. 1981).

No studies were located regarding hematological effects in animals following inhalation exposure to copper.

Hepatic Effects. Hepatomegaly was observed in workers involved in grinding and sieving copper dust (Suciu et al. 1981); the exposure levels ranged from 111 to 434 mg Cu/m³.

No studies were located regarding hepatic effects in animals following inhalation exposure to copper.

Endocrine Effects. Seven cases of enlargement of the sella turcica, nonsecretive hypophyseal adenoma, accompanied by obesity, arterial hypertension, and "red facies" were observed in a group of 100 workers exposed to 111–434 mg Cu/m³ as copper dust (Suciu et al. 1981). The study authors noted that there was a possibility that the clinical manifestations of hypophyseal adenoma or of Cushing's syndrome may have been the result of a disturbance of copper metabolism. The significance of this effect and its relationship to copper exposure cannot be determined.

Ocular Effects. Eye irritation has been reported by workers exposed to copper dust (Askergren and Mellgren 1975). The irritation is likely due to direct contact with the copper rather than a systemic effect resulting from inhalation exposure.

Other Systemic Effects. A few studies have reported metal fume fever, a 24–48-hour illness characterized by chills, fever, aching muscles, dryness in the mouth and throat, and headache, in workers exposed to copper dust or fumes (Armstrong et al. 1983; Gleason 1968). Gleason (1968) reported airborne copper dust concentrations of 0.075–0.12 mg/m³. It has been suggested that other metals present in the workplace may have been the causative agent for the metal fume fever, rather than copper. This is supported by the small number of reports of metal fume fever despite the extensive use of copper in many industries (Borak et al. 2000).

3. HEALTH EFFECTS

3.2.1.3 Immunological and Lymphoreticular Effects

No studies were located regarding immunological effects in humans following inhalation exposure to copper.

An acute exposure study in mice reported an impaired immune response following exposure to copper sulfate and a bacterial challenge (Drummond et al. 1986). Increased mortality and decreased survival time were observed in CD-1 mice challenged by an aerosol of *Streptococcus zooepidemicus* following 0.56 mg Cu/m³ for 3 hours or 0.13 mg Cu/m³ for 3 hours/day, 5 days/week for 2 weeks. Decreased bactericidal activity of alveolar macrophages was also observed in mice exposed to 3.3 mg Cu/m³ for 3 hours or 0.12 mg Cu/m³ for 3 hours/day, 5 days/week for 2 weeks following exposure to an aerosol of *Klebsiella pneumonia*.

These LOAEL values for immunological effects are recorded in Table 3-1 and plotted in Figure 3-1.

3.2.1.4 Neurological Effects

Only one study examining neurological effects was located. Headache, vertigo, and drowsiness were reported in factory workers exposed to 111–434 mg/m³ copper dust (Suciu et al. 1981).

3.2.1.5 Reproductive Effects

Sexual impotence was reported in 16% of workers (75–100 workers examined) exposed to 111–434 mg/m³ copper dust during grinding and sieving operations (Suciu et al. 1981). The significance of this finding is difficult to assess because a control group was not used.

No studies were located regarding reproductive effects in animals following inhalation exposure to copper.

Table 3-1 Levels of Significant Exposure to Copper - Inhalation

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	NOAEL (mg/m ³)	LOAEL		Reference Chemical Form
					Less Serious (mg/m ³)	Serious (mg/m ³)	
ACUTE EXPOSURE							
Systemic							
1	Mouse	3 hr	Resp	3.3			Drummond et al. 1986
2	Mouse	1-2 wk 5d/wk 3hr/d	Resp		0.12 (alveoli thickening)		Drummond et al. 1986
3	Hamster	3 hr	Resp	1.21	3.3 (decr cilia beating frequency)		Drummond et al. 1986
4	Hamster	1-2 wk 5d/wk 3hr/d	Resp	0.13			Drummond et al. 1986
Immuno/ Lymphoret							
5	Mouse	1-2 wk 5d/wk 3hr/d			0.12 (decr bactericidal activity)	0.13 (decr mean survival time)	Drummond et al. 1986
6	Mouse	3 hr			3.3 (decr bactericidal activity)	0.56 (decr mean survival time)	Drummond et al. 1986

Table 3-1 Levels of Significant Exposure to Copper - Inhalation

(continued)

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL		Reference Chemical Form
				NOAEL (mg/m ³)	Less Serious (mg/m ³)	
INTERMEDIATE EXPOSURE						
Systemic						
7	Rabbit (NS)	1 mo 5d/wk 6hr/d	Resp	0.6 M		Johansson et al. 1983 copper chloride
8	Rabbit (NS)	4-6 wk 5d/wk 6hr/d	Resp	0.6 M		Johansson et al. 1984 copper chloride
CHRONIC EXPOSURE						
Systemic						
9	Human	8 hr/d, 5 d/wk	Hemato		0.64 (decr hemoglobin and erythrocyte levels)	Finelli et al. 1981 NS

^aThe number corresponds to entries in Figure 3-1.

d = day(s); decr = decreased; hemato = hematological; hr = hour(s); LOAEL = lowest-observed-adverse-effect level; NOAEL = no-observed-adverse-effect level; (NS) = not specified; Resp = respiratory; wk = week(s); yr = year(s)

Figure 3-1. Levels of Significant Exposure to Copper- Inhalation
Acute (≤14 days)




Figure 3-1. Levels of Significant Exposure to Copper- Inhalation (*Continued*)

Intermediate (15-364 days)





Figure 3-1. Levels of Significant Exposure to Copper- Inhalation (*Continued*)
 Chronic (≥ 365 days)



3. HEALTH EFFECTS

3.2.1.6 Developmental Effects

No studies were located regarding developmental effects in humans and animals following inhalation exposure to copper.

3.2.1.7 Cancer

There are limited data for humans and no data for animals on the carcinogenicity of inhaled copper. Although a number of studies have examined cancer risk among copper smelters, these papers are not discussed because the cancer risk has been attributed to exposure to arsenic rather than to copper. In a study of over 6,700 male workers at a Chinese copper mine, significant increases in the risk of cancer (all sites combined) (standardized mortality ratio [SMR] =123; 95% confidence interval [CI] =109–139), stomach cancer (SMR=131; 95% CI=105–161), and lung cancer (SMR=147; 95% CI=112–189) were observed (Chen et al. 1993). The cancer risk increased with increasing duration of employment and time since first exposure and was also higher in workers employed in the 1950s when there was a dramatic increase in production, dry drilling methods were used, and there was poor underground ventilation. Radon and radon daughters were detected in the underground mines; between 1960 and 1990, radioactivity levels of 1.29×10^{-11} Ci/L were measured. To assess the relative contribution of radon and radon daughters to the lung cancer risk, the workers were divided into two groups: underground miners and workers involved in drilling (presumably above ground). Increases in lung cancer risk were observed in both groups, thus suggesting that exposure to radioactivity was not the primary source of increased cancer risk. The copper ore also contained silica, iron, manganese, arsenic, titanium, and sulfur. The study authors noted that the arsenic level in the copper was relatively low (0.061%) and did not likely contribute to the lung cancer risk; however, the lung cancer risk from exposure to silica and iron could not be ruled out. A significant increase in the risk of silicosis was observed in the miners. In a 7-year follow-up of this cohort (Chen et al. 1995), the risk of all sites of cancer (SMR=129; 95% CI=117–142), stomach cancer (SMR=141; 95% CI=116–169), and lung cancer (SMR=152; 95% CI=123–187) were still significantly elevated. This study also conducted a smoking survey and found that a higher percentage of the miners were smokers (71.7%) than in the control population of local residents (64.3%); this increased smoking rate, along with exposure to radioactivity, silica, iron, and arsenic may have contributed to the increased cancer risk.

3. HEALTH EFFECTS

3.2.2 Oral Exposure**3.2.2.1 Death**

A number of deaths have been reported in individuals intentionally ingesting large doses of copper sulfate (Chuttani et al. 1965). Thirteen of 53 individuals died after ingesting 6–637 mg/kg copper; because the amount of copper sulfate was self-reported, the estimated doses may be inaccurate. The deaths were attributed to shock and hepatic and/or renal complications. Deaths, probably due to central nervous system depression and hepatic and renal failure, have also been reported in individuals ingesting “spiritual green water”, which contains ≥ 100 mg copper sulfate/L (Akintonwa et al. 1989).

Increased mortality was observed in rats fed a diet containing 4,000 ppm of copper (133 mg Cu/kg/day) for 1 week. Anorexia, possibly the result of taste aversion, contributed to the deaths (Boyden et al. 1938). Weanling rats exposed to 300 mg Cu/kg/day as Cu(II) in the diet (6,000 ppm) died after 2 weeks (Haywood 1985). The deaths were attributed to extensive centrilobular necrosis.

Lifetime exposure to 42.5 mg Cu/kg/day as copper gluconate in drinking water resulted in a 12.8% reduction of the maximal lifespan (from 986 to 874 days) in mice (Massie and Aiello 1984).

The doses associated with deaths in the Haywood (1985) and Massie and Aiello (1984) studies are recorded in Table 3-2 and plotted in Figure 3-2.

3.2.2.2 Systemic Effects

No studies were located regarding endocrine, dermal, ocular, or metabolic effects in humans or animals following oral exposure to copper.

The highest NOAEL values and all reliable LOAEL values from each reliable study for systemic effects in each species and duration category are recorded in Table 3-2 and plotted in Figure 3-2.

Respiratory Effects. Data on the potential of copper to induce respiratory effects are limited to the NTP (1993) study that found no histological alterations in the lungs of rats exposed to 285 or 134 mg Cu/kg/day as copper sulfate in the diet for 14 or 90 days, respectively, or in mice exposed to 717 or 814 mg Cu/kg/day for 14 or 90 days.

Table 3-2 Levels of Significant Exposure to Copper - Oral

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL			Reference Chemical Form
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)	Serious (mg/kg/day)	
ACUTE EXPOSURE							
Death							
1	Rat (Wistar)	2-15 wk (F)				550 M (increased mortality)	Haywood 1985 NS
2	Rat (Fischer- 344)	14 d (W)				31 F (100% mortality)	NTP 1993 copper sulfate
3	Mouse (B6C3F1)	14 d (W)				62 M (increased mortality)	NTP 1993 copper sulfate
Systemic							
4	Human	once (W)	Gastro	0.011	0.017 (nausea, vomiting, diarrhea, or abdominal pain)		Araya et al. 2001 copper sulfate
5	Human	once (W)	Gastro	0.012 F	0.018 F (nausea)		Araya et al. 2003a copper sulfate
6	Human	once (W)	Gastro		0.046 (nausea, delayed gastric emptying)		Araya et al. 2003c copper sulfate
7	Human	once (W)	Gastro		0.03 (nausea and vomiting)		Gotteland et al. 2001 copper sulfate

Table 3-2 Levels of Significant Exposure to Copper - Oral

(continued)

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL		Reference Chemical Form
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)	
8	Human	once (W)	Gastro		6 (vomiting)	Karlsson and Noren 1965 copper sulfate
9	Human	once (W)	Gastro		0.08 M (vomiting, diarrhea)	Nicholas and Brist 1968 NS
10	Human	once (W)	Gastro	0.0057	0.011 (nausea)	Olivares et al. 2001 copper sulfate
11	Human	2 wks (W)	Gastro	0.0272 ^b F	0.0731 F (abdominal pain, nausea, and/or vomiting)	Pizarro et al. 1999 copper sulfate
12	Human	1 wk (W)	Gastro		0.096 F (nausea, vomiting, and/or abdominal pain)	Pizarro et al. 2001 copper sulfate and copper oxide
13	Rat (NS)	1-2 wk (F)	Hepatic		300 M (parenchymal cell hypertrophy)	Haywood 1980 copper sulfate
			Renal	300 M		

Table 3-2 Levels of Significant Exposure to Copper - Oral

(continued)

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL		Reference Chemical Form
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)	
14	Rat (NS)	1-2 wk (F)	Hepatic		300 M (increased alanine aminotransferase activity)	Haywood and Comerford 1980 copper sulfate
15	Rat (Wistar)	1-2 wk (F)	Hepatic		450 M (hepatocellular necrosis)	Haywood et al. 1985a NS
			Renal		450 M (copper-containing droplets and granules in proximal tubule cells)	
16	Rat (Wistar)	2 wk (F)	Renal		200 M (droplets in proximal tubule lumen)	Haywood et al. 1985b NS

Table 3-2 Levels of Significant Exposure to Copper - Oral

(continued)

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL		Reference Chemical Form	
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)		Serious (mg/kg/day)
17	Rat (Fischer- 344) (W)	14 d	Resp	29 M			NTP 1993 copper sulfate
			Cardio	29 M			
			Gastro	29 M			
			Hepatic	29 M			
			Renal		10 M (protein droplets in epithelial cells of proximal tubule)		
Bd Wt	26 F						

Table 3-2 Levels of Significant Exposure to Copper - Oral

(continued)

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL		Reference Chemical Form	
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)		Serious (mg/kg/day)
18	Rat (Fischer- 344) (F)	14 d	Resp	285 F			NTP 1993 copper sulfate
			Cardio	285 F			
			Gastro	23 F	44 F (hyperplasia of forestomach mucosa)		
			Hemato	93 F	196 F (depletion of hematopoietic cells in bone marrow)		
			Hepatic	92 M	198 M (inflammation)		
			Renal	46 M	92 M (increased protein droplets in cortical tubules)		
			Bd Wt	93 F	196 F (18% decrease in body weight gain)		

Table 3-2 Levels of Significant Exposure to Copper - Oral

(continued)

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL		Reference Chemical Form	
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)		Serious (mg/kg/day)
19	Mouse (B6C3F1)	14 d (W)	Resp	24 M			NTP 1993 copper sulfate
			Cardio	24 M			
			Gastro	24 M			
			Hepatic	24 M			
			Renal	24 M			
			Bd Wt	24 M			

Table 3-2 Levels of Significant Exposure to Copper - Oral

(continued)

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL		Reference Chemical Form	
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)		Serious (mg/kg/day)
20	Mouse (B6C3F1)	14 d (F)	Resp	717 M			NTP 1993 copper sulfate
			Cardio	717 M			
			Gastro	92 M	197 M (hyperplasia of forestomach mucosa)		
			Hepatic	717 M			
			Renal	717 M			
			Bd Wt	717 M			
INTERMEDIATE EXPOSURE							
Systemic							
21	Human	daily 2 months (W)	Gastro	0.042 ^C	0.091 (gastrointestinal symptoms)		Araya et al. 2003b copper sulfate
			Hepatic	0.17			

Table 3-2 Levels of Significant Exposure to Copper - Oral

(continued)

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL			Reference Chemical Form
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)	Serious (mg/kg/day)	
22	Human	9 months (W)	Gastro	0.319			Olivares et al. 1998 copper sulfate
			Hepatic	0.319			
			Bd Wt	0.319			
23	Human	12 wks (C)	Gastro	0.14			Pratt et al. 1985 copper gluconate
			Hemato	0.14			
			Hepatic	0.14			
24	Rat (Fischer- 344) (F)	3 mo	Hepatic	66 M	89 M (increased number of necroinflammatory foci in the liver)		Aburto et al. 2001b copper sulfate
			Bd Wt	114 M	140 M (15% decreased in terminal body weight)		

Table 3-2 Levels of Significant Exposure to Copper - Oral

(continued)

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL		Reference Chemical Form
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)	
25	Rat (Sprague- Dawley)	30-58 d (F)	Hepatic	20 F		Cristofori et al. 1992 NS
			Renal	20 F		
26	Rat (Sprague- Dawley)	90 d (W)	Hepatic		8 M (increased aspartate aminotransferase activity)	Epstein et al. 1982 copper acetate
			Bd Wt	8 M		
27	Rat (Fischer- 344)	18 wks (F)	Hepatic		150 M (inflammation and increased serum enzyme activity in adult rats)	Fuentelba et al. 2000 copper sulfate
					120 M (inflammation, necrosis, and increases serum enzyme levels in young rats)	
28	Rat (NS)	3-15 wk (F)	Hepatic		180 M (necrosis)	Haywood 1980 copper sulfate
			Renal		180 M (cytoplasmic droplets and desquamation of epithelial cells in proximal tubules)	

Table 3-2 Levels of Significant Exposure to Copper - Oral

(continued)

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL		Reference Chemical Form	
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)		Serious (mg/kg/day)
29	Rat (Wistar)	2-15 wk (F)	Hepatic		280 M (inflammation, necrosis)	550 M (chronic hepatitis)	Haywood 1985 NS
			Renal		280 M (degeneration of proximal tubule cells)		
			Bd Wt			550 M (weight loss) 280 M (50% decrease in body weight gain)	
30	Rat (NS)	3-15 wk (F)	Hepatic		180 M (increased alanine aminotransferase activity)		Haywood and Comerford 1980 copper sulfate
31	Rat (Wistar)	15 wk (F)	Hepatic		320 M (necrosis)	640 M (chronic hepatitis)	Haywood and Loughran 1985 copper sulfate
			Bd Wt			640 M (weight loss) 320 M (50% decrease in body weight gain)	

Table 3-2 Levels of Significant Exposure to Copper - Oral

(continued)

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL		Reference Chemical Form
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)	
32	Rat (Wistar)	4-14 wks (F)	Hepatic		280 M (hepatocellular necrosis)	Haywood et al. 1985a NS
			Renal		280 M (tubular cell necrosis)	
33	Rat (Wistar)	4-15 wk (F)	Renal		200 M (reversible degeneration and necrosis of tubule cells)	Haywood et al. 1985b NS
34	Rat (NS)	30 d (G)	Hemato		100 M (decreased erythrocyte and hemoglobin levels)	Kumar and Sharma 1987 copper sulfate
			Hepatic		100 M (increased glucose, cholesterol, bilirubin, serum enzymes, and decreased total protein levels)	
			Renal		100 M (increased BUN levels)	
35	Rat (Wistar)	15 wks (F)	Cardio		14 M (increased blood pressure)	Liu and Medeiros 1986 copper carbonate

Table 3-2 Levels of Significant Exposure to Copper - Oral

(continued)

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL		Reference Chemical Form
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)	
36	Rat (Holtzman)	21 wks (F)	Musc/skel	120 M		Llewellyn et al. 1985 copper acetate
			Bd Wt		120 (23% decrease in body weight gain)	

Table 3-2 Levels of Significant Exposure to Copper - Oral

(continued)

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL			Reference Chemical Form
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)	Serious (mg/kg/day)	
37	Rat (Fischer- 344) (F)	13 wk	Resp	134 F			NTP 1993 copper sulfate
			Cardio	134 F			
			Gastro	16 M	33 M (squamous mucosa hyperplasia of forestomach)		
			Hemato	33 M	66 M (decreases in hematocrit, hemoglobin, reticulocytes, mean cell volume, and mean cell hemoglobin levels and increases in platelet levels)		
			Hepatic	8 M	66 M (chronic active inflammation with focal necrosis)		
					16 M (increases serum alanine aminotransferase)		
			Renal	9 F	17 F (increased BUN)	134 F (tubular degeneration)	
Bd Wt	66 M	140 M (24% decrease in body weight gain)					

Table 3-2 Levels of Significant Exposure to Copper - Oral

(continued)

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL		Reference Chemical Form
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)	
38	Rat (NS)	20 d (G)	Hemato		100 M (decreases in erythrocyte, hemoglobin, and hematocrit levels)	Rana and Kumar 1980 copper sulfate
			Hepatic		100 M (hepatocellular necrosis)	
			Renal		100 M (tubular cell necrosis)	
39	Mouse (B6C3F1)	13 wk (F)	Resp	814 M		NTP 1993 copper sulfate
			Cardio	814 M		
			Gastro	126 F	267 F (hyperplasia of forestomach mucosa)	
			Hepatic	814 M		
			Renal	814 M		
		Bd Wt	187 M	398 M (12% decrease in body weight gain)		

Table 3-2 Levels of Significant Exposure to Copper - Oral

(continued)

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL		Reference Chemical Form	
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)		Serious (mg/kg/day)
40	Pig (Hampshire)	54 d (F)	Hemato	11	24	(decreased hemoglobin levels)	Kline et al. 1971 copper sulfate
			Bd Wt	11	24	(decreased body weight gain)	
41	Pig (NS)	49 d (F)	Hemato		36 F	(decreased hemoglobin levels)	Suttle and Mills 1966a copper carbonate
			Hepatic		36 F	(increased aspartate aminotransferase activity)	
42	Pig (NS)	6 wks (F)	Hemato		35 F	(decreased hemoglobin level)	Suttle and Mills 1966a copper carbonate
			Hepatic		35 F	(increased aspartate aminotransferase activity)	
43	Mouse (C57BL/6N)	8 wks (W)	Immuno/ Lymphoret		24	(impaired immune function)	Pocino et al. 1990 copper sulfate
44	Mouse (C57BL/6N)	3-5 or 8-10 wks (W)			13	(altered cell-mediated and humoral immunity)	Pocino et al. 1991 copper sulfate

Table 3-2 Levels of Significant Exposure to Copper - Oral

(continued)

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL		Reference Chemical Form
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)	
Neurological						
45	Rat (Sprague- Dawley)	11 mo (W)			36 F (decreased 3,4-dihydroxyphenylacetic acid levels in corpus striatum)	DeVries et al. 1986 copper sulfate
46	Rat (NS)	30 d (F)		23		Murthy et al. 1981 copper sulfate
Reproductive						
47	Rat (Fischer- 344)	13 wk (F)		66 M 68 F		NTP 1993 copper sulfate
48	Mouse (B6C3F1)	13 wk (F)		398 M 536 F		NTP 1993 copper sulfate
49	Mink (dark mink)	153 or 367 d (F)		12		Aulerich et al. 1982 copper sulfate

Table 3-2 Levels of Significant Exposure to Copper - Oral

(continued)

Key to figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL			Reference Chemical Form
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)	Serious (mg/kg/day)	
Developmental							
50	Rat (Wistar)	60-73 d (W)			130 (delayed growth and development)		Haddad et al. 1991 copper acetate
51	Mouse (C57BL/6N)	1 mo + gd 0-19 (F)		138 F	208 (decreased mean litter size and fetal body weights)		Lecyk 1980 copper sulfate
52	Other (dark mink)	153 or 367 d (F)		13			Aulerich et al. 1982 copper sulfate
CHRONIC EXPOSURE							
Death							
53	Mouse (C57BL/6N)	850 d (W)				4.2 (14.7% decrease in lifespan)	Massie and Aiello 1984 copper gluconate
Systemic							
54	Mouse (C57BL/6N)	850 d (W)	Bd Wt	42 M			Massie and Aiello 1984 copper gluconate

a The number corresponds to entries in Figure 3-2.

b Used to derive an acute-duration oral minimal risk level (MRL) of 0.01 mg Cu/kg/day; the NOAEL was divided by an uncertainty factor of 3 to account for human variability.

c Used to derive an intermediate-duration minimal risk level (MRL) of 0.01 mg Cu/kg/day; the NOAEL divided by an uncertainty factor of 3 to account for human variability.

Bd Wt = body weight; Cardio = cardiovascular; CEL = cancer effect level; d = day(s); Endocr = endocrine; (F) = feed; F = Female; G = gavage; Gastro = gastrointestinal; gd = gestational day; Gn pig = guinea pig; hemato = hematological; hr = hour(s); LOAEL = lowest-observed-adverse-effect level; M = male; min = minute(s); mo = month(s); Musc/skel = musculoskeletal; NOAEL = no-observed-adverse-effect level; occup = occupational; NS = not specified; Resp = respiratory; (W) = drinking water; wk = week(s)

Figure 3-2. Levels of Significant Exposure to Copper - Oral
Acute (≤ 14 days)




Figure 3-2. Levels of Significant Exposure to Copper - Oral (Continued)
Intermediate (15-364 days)




Figure 3-2. Levels of Significant Exposure to Copper - Oral (Continued)

Intermediate (15-364 days)





Figure 3-2. Levels of Significant Exposure to Copper - Oral (*Continued*)
 Chronic (≥ 365 days)



3. HEALTH EFFECTS

Cardiovascular Effects. Several human studies have examined the possible relationship between increased serum copper levels and an increased risk of coronary heart disease. Although a number of studies have found increased risk of coronary heart disease deaths with increasing serum copper levels (Ford 2000), a number of studies have not found a relationship. However, whether copper directly affects atherosclerosis or is a marker of inflammation associated with atherosclerosis remains to be established.

There are limited data on the toxicity of copper to the cardiovascular system. A significant increase in systolic blood pressure was observed in rats exposed to 14 mg Cu/kg/day as copper carbonate in the diet for 15 weeks (Liu and Mederios 1986). No histological alterations were observed in the hearts of rats or mice exposed to 285 or 717 mg Cu/kg/day, respectively, for 14 days or 134 or 814 mg Cu/kg/day for 90 days (NTP 1993).

Gastrointestinal Effects. There are numerous reports of acute gastrointestinal effects in humans after ingestion of large amounts of copper in drinking water or beverages. The most prevalent effects are nausea and vomiting, which typically occur shortly after ingestion and are not persistent (Araya et al. 2001, 2003a, 2003b, 2003c; Chuttani et al. 1965; Eife et al. 1999; Gill and Bhagat 1999; Gotteland et al. 2001; Holleran 1981; Jantsch et al. 1984, 1985; Karlsson and Noren 1965; Knobeloch et al. 1994, 1998; Nicholas and Brist 1968; Olivares et al. 2001; Pizarro et al. 1999, 2001; Semple et al. 1960; Spitalny et al. 1984; Walsh et al. 1977). Abdominal pain and diarrhea have also been reported, but their incidence is typically much lower than nausea and vomiting. Although most of the data on gastrointestinal effects in humans come from case reports of accidental exposure from contaminated beverages with limited information on exposure levels, several recently conducted studies were designed to identify the threshold for gastrointestinal effects. These experiments typically involve adults ingesting a single dose of copper sulfate following an overnight fast (Araya et al. 2001, 2003a, 2003b; Gotteland et al. 2001; Olivares et al. 2001). Olivares et al. (2001) identified the lowest LOAEL for gastrointestinal effects; a significant increase in the incidence of nausea was observed at 4 ppm copper (0.01 mg Cu/kg) and higher. At 6 ppm, a significant increase in the incidence of vomiting was also observed. Administering the copper sulfate in an orange-flavored drink increased the threshold for nausea to 8 ppm (0.022 mg Cu/kg) (Olivares et al. 2001). In two multinational studies conducted by Araya and associates (Araya et al. 2001, 2003a), NOAEL and LOAEL values of 4 and 6 ppm (0.042 and 0.091 mg Cu/kg), respectively, were identified for nausea. Araya et al. (2003a) determined that both the copper concentration and the total copper dose are important variables in predicting a response; as the concentration and dose increase, the probability of eliciting nausea increases.

3. HEALTH EFFECTS

Similar thresholds for effects were observed in repeated exposure studies (Araya et al. 2003c; Pizarro et al. 1999, 2001). Abdominal pain, nausea, and/or vomiting were observed in women drinking water containing 5 ppm (0.096 mg Cu/kg) copper sulfate or copper oxide for 1 week (Pizarro et al. 2001). The occurrence of gastrointestinal effects (excluding diarrhea) was not significantly different in subjects ingesting copper sulfate and those ingesting copper oxide. In a 2-week exposure study by Pizarro et al. (1999), significant increases in the incidence of gastrointestinal symptoms were observed in subjects exposed to 3 or 5 ppm (0.0731 and 0.124 mg Cu/kg/day), but not to 1 ppm (0.0272 mg Cu/kg/day). The incidences of nausea, vomiting, and/or abdominal pain were 5, 2, 17, and 15% in the control, 1, 3, and 5 ppm groups, respectively. In a similarly designed study, Araya et al. (2003b) examined the occurrence of gastrointestinal symptoms in adults exposed to copper sulfate for 2 months. The incidences of gastrointestinal symptoms were 11.7, 15.3, 18.3, and 19.7% in the control, 2, 4, and 6 ppm groups, respectively. As analyzed using the chi-square test with Bonferroni correction, the incidence was significantly elevated in the 6 ppm (0.17 mg Cu/kg/day) group; if the Bonferroni correction was not used, the incidence was significantly elevated in the 4 ppm (0.091 mg Cu/kg/day) group. A case report by Spitalny et al. (1984) also examined the effects of repeated exposure to copper. Recurrent, acute symptoms, including nausea, vomiting, and abdominal pain, were reported by three of four family members shortly after drinking juice, coffee, or water in the morning. The effects disappeared when the family switched to bottled water. An early morning water sample contained 7.8 ppm copper. A study by Buchanan et al. (1991) also examined individuals with elevated levels of copper in household water. The occurrence of vomiting and nausea with abdominal pain was not significantly different among residents with a first-draw water sample of 3 ppm or higher, as compared to controls with less than 1.3 ppm copper in first-draw sample. The investigators noted that in a case-control study of this population, all of the cases reported that none of the subjects obtained their water immediately from the tap, but most (70%) only let it run for less than 1 minute. The study found that copper content in the tap water used for drinking averaged 14% of first draw samples.

Most of the available human studies examined the relationship between copper exposure and the manifestation of symptoms of gastrointestinal irritation; Gotteland et al. (2001) and Araya et al. (2003c) also looked at physiological alterations. Gotteland et al. (2001) found significant increases in gastric permeability to sucrose following the bolus ingestion of 10 ppm copper as copper sulfate (0.03 mg Cu/kg); no alterations in intestinal permeability to lactulose/mannitol were found. The increased gastric permeability was independent of gastrointestinal symptoms. A significant delay in decreasing the stomach's antral area was found during the first hour after bolus ingestion of 10 ppm copper as copper

3. HEALTH EFFECTS

sulfate (0.046 mg Cu/kg) (Arayaet et al. 2003c). This change in antral area is suggestive of a delay in gastric emptying. As with gastric permeability, this effect was independent of gastrointestinal symptoms.

Gastrointestinal effects have also been reported in animal studies. Hyperplasia with hyperkeratosis of the squamous mucosa on the limiting ridge separating the forestomach from the glandular stomach was observed in rats and mice exposed to 44 and 197 mg Cu/kg/day, respectively, as copper sulfate in the diet for 14 days or 33 and 267 mg Cu/kg/day, respectively, as copper sulfate in the diet for 13 weeks (NTP 1990a). No gastrointestinal effects were observed in rats and mice exposed to 23 or 92 mg Cu/kg/day for 14 days or in rats and mice exposed to 16 or 126 mg Cu/kg/day 13 weeks. Additionally, no gastrointestinal effects were observed in rats and mice exposed to 29 or 24 mg Cu/kg/day as copper sulfate in drinking water (NTP 1990a).

Hematological Effects. There are limited data on the effect of copper on the human hematological system. Acute hemolytic anemia was observed in an 18-month-old child 2 days after he drank a solution containing approximately 3 g of copper sulfate (Walsh et al. 1977). Acute intravascular hemolysis was also reported in 5 of 125 individuals intentionally ingesting a large dose of copper sulfate (Ahasan et al. 1994). No alterations in hematocrit level or mean corpuscular volume were observed in individuals ingesting 0.14 mg Cu/kg/day as copper gluconate in a capsule for 12 weeks (Pratt et al. 1985).

Information on the hematological effects in animals associated with exposure to high levels of copper is also limited to several studies that measured hemoglobin and hematocrit values. Decreased hemoglobin and hematocrit values were observed in rats exposed to ≥ 66 mg Cu/kg/day (Kumar and Sharma 1987; NTP 1993; Rana and Kumar 1980) for 20–90 days and in pigs exposed to ≥ 24 mg Cu/kg/day for 48–54 days (Kline et al. 1971; Suttle and Mills 1966a, 1966b). Depletion of hematopoietic cells in the bone marrow was observed in rats exposed to 196 mg Cu/kg/day as copper sulfate in the diet for 14 days (NTP 1993). Contrary to these findings, Liu and Medeiros (1986) observed an increase in hemoglobin levels and no change in hematocrit levels in rats fed a diet containing 14 mg Cu/kg/day as copper carbonate for 20 weeks.

Musculoskeletal Effects. No studies were located regarding musculoskeletal effects in humans following oral exposure to copper.

Equivocal results on the effects of copper on the musculoskeletal system have been found. Depressed skeletal growth has been observed in rats administered 100 mg Cu/kg/day via gavage; tail length was used

3. HEALTH EFFECTS

to assess skeletal growth (Rana and Kumar 1980). Using radiographic data, no qualitative or quantitative differences were observed in bones of rats exposed to 120 mg Cu/kg/day as copper acetate in the diet for 21 weeks (Llewellyn et al. 1985). The different outcomes may reflect the different methods used to assess skeletal growth.

Hepatic Effects. With the exception of several defined syndromes—Wilson’s disease, Indian childhood cirrhosis, and idiopathic copper toxicosis—liver effects are rarely reported in humans, although this has not been extensively investigated. In a compilation of case reports of individuals intentionally ingesting copper sulfate, jaundice was reported in 11 of 53 individuals (Chuttani et al. 1965). Centrilobular necrosis, biliary stasis, elevated serum bilirubin level and aspartate aminotransferase activity, and elevated bile salts in the urine were found in five of the individuals with jaundice. Jaundice (Akintonwa et al. 1989), centrilobular congestion (Lamont and Duflou 1988), and acute hepatotoxicity (Ahasan et al. 1994) have also been reported in case reports of lethal ingestion of copper sulfate. O’Donohue et al. (1993) reported a case of an adult with jaundice and hepatomegaly following a 3-year exposure to copper supplements. For 2 years, the individual ingested 30 mg/day followed by 1 year of 60 mg/day. In a study of seven adults receiving capsules containing 0.14 mg Cu/kg/day as copper gluconate, no significant alterations in serum aspartate aminotransferase, alkaline phosphatase, serum gamma glutamyl transferase, or lactate dehydrogenase activities were found (Pratt et al. 1985). No alterations in biomarkers of liver damage (serum aspartate aminotransferase, alanine aminotransferase, gamma glutamyl transferase) were observed in adults exposed to 0.17 mg Cu/kg/day as copper sulfate in drinking water for 2 months (Araya et al. 2003b).

Several studies have examined liver function in infants exposed to elevated levels of copper in drinking water. A no adverse effect level for liver effects was identified in a study of infants (3 months of age at study initiation) exposed to 0.315 mg Cu/kg/day as copper sulfate in drinking water for 9 months (Olivares et al. 1998). No alterations in total bilirubin levels or serum alanine aminotranferase, aspartate aminotransferase, or gamma-glutamyl transferase activities were found. A higher percentage of copper-exposed infants (30.4%) were withdrawn from the study, as compared to the control group (11.1%). The reasons for being withdrawn from the study were blood sampling refusal (eight infants in the copper group and two infants in the control group), protocol transgression (four infants in the copper group and no infants in the control group), and change of address (five infants in the copper group and one infant in the control group). Two recent surveys of infants exposed to 0.8 mg Cu/L in household water did not find significant alterations in serum parameters of liver function or alterations in liver ultrasound imaging tests (Zietz et al. 2003a, 2003b).

3. HEALTH EFFECTS

There is strong evidence to suggest that Wilson's disease, Indian childhood cirrhosis, and possible idiopathic copper toxicosis are caused by an increased genetic susceptibility to copper toxicity.

Wilson's Disease. Wilson's disease is an autosomal recessive genetic disorder with a worldwide occurrence of 1 in 30,000 to 1 in 100,000 depending on the population (Llanos and Mercer 2002; Scheinberg and Sternlieb 1996). It is characterized by high levels of copper in the liver and low levels of serum ceruloplasmin. The accumulation of copper in the liver is due to a genetic defect in one of the Cu-ATPases (ATP7B), resulting in impaired biliary excretion of copper. One of the early manifestations of the disease, typically at 8–12 years of age, is liver damage. Three types of liver damage are seen—cirrhosis, chronic active hepatitis, and fulminant hepatic failure. It is unlikely that the manifestation of Wilson's disease is related to exposure to high levels of copper; high levels of hepatic copper have been observed in affected individuals consuming average copper intakes (Scheinberg and Sternlieb 1996).

Indian Childhood Cirrhosis (ICC). ICC is a type of cirrhosis typically seen in infants and young children (6 months to 5 years of age with a mean of 18 months) living in rural areas of the Indian subcontinent. Other features include high rates of parental consanguinity and up to 22% of siblings affected (Pandit and Bhave 1996; Tanner 1998). Two of the most discriminatory features of ICC are coarse, dark brown orcein staining (representing copper) and intralobular pericellular fibrosis (Pandit and Bhave 1996). Liver copper levels ranging from 790 to 6,654 $\mu\text{g/g}$ dry weight (mean of 939 $\mu\text{g/g}$) were found in 53 children diagnosed with ICC, as compared to levels of 8–118 $\mu\text{g/g}$ (mean 42–45 $\mu\text{g/g}$) in 12 controls aged 6 months to >1 year (Bhave et al. 1982); interpretation of these study results is limited by the small number of controls and the lack of detail on the control group.

In a study of 100 children with ICC and 100 age-, sex-, and caste-matched controls, it was determined that ICC was attributable to the early introduction of cow or buffalo milk feeds contaminated with copper from brass vessels, which were used to store and heat the milk (Bhave et al. 1987). Although a cause and effect relationship between high copper intake and ICC has not been firmly established, there is strong evidence to support an association. In another study in which the parents of 100 children with ICC were advised to use aluminum or stainless steel vessels for preparing infant milk feeds, only 1 of 86 younger siblings of the children with ICC developed ICC (this child was known to have received copper-contaminated milk) as compared to 30 of 125 older siblings (Tanner 1998).

Idiopathic Copper Toxicosis (ICT). Although there are limited data on ICT, it is also believed to be caused by an autosomal-recessive inherited defect in copper metabolism and excess dietary copper

3. HEALTH EFFECTS

(Müller et al. 1998; Wijmenga 2002). In the literature, ICT is also referred to as Indian childhood cirrhosis-like liver disease, copper-associated liver disease in childhood, and Tyrollean infantile cirrhosis. In the last 25 years, there have been <200 cases of ICT reported in a number of countries including Australia, Austria, Germany, Ireland, Italy, Kuwait, Mexico, United Kingdom, and United States. With the exception of a study of ICT in 138 children living in Tyrol Austria (Müller et al. 1996), most papers describe the clinical course of 1–4 children. Compiling the data from these studies, Müller et al. (1998) found a number of patterns: (1) the age of onset of clinical symptoms occurs before the age of 2 years (infantile onset) or before the age of 5 years (late onset), although onset as late as 10 years has also been observed; (2) rapid progression and death within 2 weeks to 11 months; (3) very high copper levels in the liver, 190–3,360 µg/g dry weight (normal is <50 µg/g); (4) abnormal biochemical markers of liver damage such as aminotransferases, alkaline phosphatase, bilirubin, albumin, and prothrombin time; and (5) marked panlobular and pericellular fibrosis associated with an usually mild inflammatory infiltrate, ballooning degeneration of hepatocytes, and an abundance of Mallory bodies. The high levels of copper in the liver, the identification of environmental copper exposure, and the similarity of the clinical presentation and histopathology with ICC suggest that copper is the causative agent. As with ICC, an increased genetic susceptibility to copper toxicity has been suggested. A genealogic investigation conducted by Müller et al. (1996) provided suggestive evidence that the disease is transmitted in an autosomal recessive mode.

The hepatotoxicity of copper in animals has been described and investigated in a number of acute- and intermediate-duration oral exposure studies. The majority of these studies used rats; a small number of studies used pigs and mice. In addition to these studies, there are a number of studies in animals with similar genetic defects as Wilson's disease, including Long Evans Cinnamon (LEC) rats and Bennington terrier dogs. The results of these studies were not considered relevant to healthy humans and will not be discussed. The earliest symptoms of hepatotoxicity in rats orally exposed to copper are increases in serum chemistry enzymes, particularly alanine aminotransferase and aspartate aminotransferase (Epstein et al. 1982; Fuentealba et al. 2000; Haywood 1980; Haywood and Comerford 1980; Kumar and Sharma 1987; NTP 1993; Sugawara et al. 1995). Continued exposure or exposure to higher concentrations can result in inflammation, parenchymal cell hypertrophy, and hepatocellular necrosis (Aburto et al. 2001b; Fuentealba et al. 2000; Haywood 1980, 1985; Haywood and Loughran 1985; Haywood et al. 1985a; NTP 1993). At very high doses, chronic hepatitis (Haywood 1985; Haywood and Loughran 1985) has also been observed.

3. HEALTH EFFECTS

Studies in rats provide information on the dose-response relationships as a function of exposure duration. The highest NOAEL and lowest LOAEL for liver effects in acutely exposed rats are 92 and 198 mg Cu/kg/day, respectively, administered as copper sulfate in the diet for 14 days (NTP 1993). Chronic inflammation was observed at the LOAEL; a LOAEL for serum chemistry changes was not identified in the available acute exposure studies because the only study testing low doses (NTP 1993) did not assess this parameter. The threshold for hepatotoxicity in rats following intermediate-duration exposure appears to be between 8 - 16 mg Cu/kg/day. NTP (1993) found a significant increase in serum alanine aminotransferase activity in Fischer 344 rats exposed to 16 mg Cu/kg/day as copper sulfate in the diet for 13 weeks; no effects were observed at 8 mg Cu/kg/day. However, Epstein et al. (1982) found a significant increase in aspartate aminotransferase in Sprague-Dawley rats exposed to 8 mg Cu/kg/day as copper sulfate in drinking water for 90 days; differences in the exposure route and rat strain may have contributed to these differences. Histological damage (chronic active inflammation and focal hepatocellular necrosis) has been observed at 66 mg Cu/kg/day (administered as copper sulfate in the diet for 90 days) and higher (NTP 1993). Severe hepatic damage (chronic hepatitis) has been observed in rats exposed to >550 mg Cu/kg/day as copper sulfate in the diet for 15 weeks (Haywood 1985; Haywood and Loughran 1985).

The available rat hepatotoxicity data, along with toxicokinetic data, suggest that there are three phases of copper toxicity in the rat. In the first phase, copper levels increase in the liver, with minimal to no damage to hepatic tissues. As the hepatic copper levels increase, inflammation and necrosis occur. Thereafter, the copper levels in the liver begin to decrease and the parenchymal tissue begins to regenerate. At this point, the animal develops a tolerance to copper. Haywood et al. (1985a) speculated that the tolerance resulted from a shift in the liver from copper storage and biliary excretion to copper transport and renal clearance. Tolerance appears to protect the animals from subsequent liver toxicity. For example, no adverse liver effects were observed in rats exposed to 640 mg Cu/kg/day as copper sulfate in the diet when this exposure was preceded by a 15-week exposure to 320 mg Cu/kg/day as copper sulfate in the diet. This is in contrast to the severe hepatocellular necrosis that was observed in animals exposed to a control diet for 15 weeks followed by a 3-week exposure to 640 mg Cu/kg/day (Haywood and Loughran 1985). The time course of each phase of liver toxicity appears to be dose-related. At higher doses, the onset of the necrosis and regeneration occurred earlier as compared to lower doses. Additionally, there appears to be an upper limit of copper intake, which would induce copper tolerance; doses that exceed this level would result in permanent damage to the liver. Dietary exposure of rats to ≥ 550 mg Cu/kg/day as copper sulfate for 15 weeks resulted in chronic hepatitis with no evidence of regeneration of parenchymal tissue (Haywood and Loughran 1985).

3. HEALTH EFFECTS

There are limited experimental data on the hepatotoxicity of copper in other animal species. Pigs fed a diet providing 35–36 mg Cu/kg/day for 7 weeks had a significant increase in aspartate aminotransferase activities (Suttle and Mills 1966a, 1966b). It appears that rats and pigs are equally sensitive to high levels of copper in the diet or drinking water. In contrast, mice do not appear to be as sensitive to the hepatic toxicity of copper as rats. No hepatic effects were observed in mice exposed to 814 mg Cu/kg/day for 13 weeks as compared to rats, which exhibited an increase in alanine aminotransferase activity at 16 mg Cu/kg/day and chronic active inflammation at 66 mg Cu/kg/day (NTP 1993).

Renal Effects. There is limited information on the renal toxicity of copper in humans. Congestion of the glomeruli and denudation of tubular cells were observed in four individuals consuming a single lethal dose of copper sulfate (Chuttani et al. 1965). Acute renal failure was reported in 5 of 125 individuals intentionally ingesting large doses of copper sulfate (Ahasan et al. 1994). Hematuria, glycosuria, cylindruria, and proteinuria, indicative of renal tubular damage, were observed in a child who drank a solution containing approximately 3 g of copper sulfate (Walsh et al. 1977).

A number of animal studies confirm that the kidney is a target of copper toxicity. Renal toxicity as a result of copper loading follows a specific time course (Haywood 1980, 1985; Haywood et al. 1985a, 1985b). No treatment-related effects were observed in rats exposed to 300 mg Cu/kg/day as copper sulfate in the diet for 1–2 weeks (Haywood 1980). However, eosinophilic droplets were observed in the epithelial cell cytoplasm of the proximal convoluted tubules in rats exposed to 450 mg Cu/kg/day for 2 weeks (Haywood et al. 1985a). The number of eosinophilic droplets increased with increasing duration (Haywood 1980, 1985). Exposure to 100–280 mg Cu/kg/day for 3–5 weeks resulted in necrosis and degeneration of proximal tubule cells (Haywood 1985; Haywood et al. 1985a, 1985b; Rana and Kumar 1980). After 9 weeks, extensive desquamation of the epithelial cells of the proximal convoluted tubules was evident in rats exposed to 180 mg Cu/kg/day (Haywood 1980). Complete regeneration of the proximal tubules was observed after 15 weeks of copper treatment in rats exposed to 180–280 mg Cu/kg/day (Haywood 1980, 1985; Haywood et al. 1985a, 1985b). In contrast to the Haywood and associates studies, a 13-week study by NTP (1993) did not find evidence of regeneration of renal tissue. An increase in protein droplets in epithelial cell cytoplasm and the lumen of the proximal convoluted tubules was observed in rats exposed to 10 or 92 mg Cu/kg/day as copper sulfate in drinking water or diet, respectively, for 2 weeks or to 33 mg Cu/kg/day as copper sulfate in the diet for 13 weeks. At 134 mg Cu/kg/day, karyomegaly and tubule cell degeneration were also observed. Additional renal effects observed in the intermediate-duration study included an increase in serum urea nitrogen levels in females exposed to ≥ 17 mg Cu/kg/day, increased urinary glucose output in males exposed to ≥ 66 mg

3. HEALTH EFFECTS

Cu/kg/day, and increased urinary aspartate aminotransferase and N-acetyl- β -glucosaminidase activities in male and female rats exposed to 140 or 134 mg Cu/kg/day, respectively. The NTP (1993) study identified a NOAEL of 9 mg Cu/kg/day. No effects were observed in mice fed a diet for 13 weeks which provided 814 mg Cu/kg/day as copper sulfate (NTP 1993).

Body Weight Effects. No studies were located regarding body weight effects in humans following oral exposure to copper.

Dietary exposure studies have reported 12–24% decreases in body weight gain in rats following exposure to 120–140 mg Cu/kg/day for 2–15 weeks (Llewellyn 1985; NTP 1993), in mice following exposure to 398 mg Cu/kg/day for 13 weeks (NTP 1993), or in pigs (magnitude of decreased weight gain not reported) following exposure to 24 mg Cu/kg/day for 54 days (Kline et al. 1971). No effect levels of 66 (NTP 1993), 187 (NTP 1993), and 11 mg Cu/kg/day (Kline et al. 1971) have been reported in rats, mice, and pigs, respectively; Epstein et al. (1982) also reported no adverse effects on body weight gain in rats exposed to 8 mg Cu/kg/day in drinking water. More severe decreases in body weight gain and weight loss have also been reported (Haywood 1985; Haywood and Loughran 1985); the weight loss was reported at lethal concentrations. Only one study examined the effect of copper on body weight gain following chronic-duration exposure (lifetime exposure beginning at 58 days of age); this study found no biologically significant effect in mice exposed to 42 mg Cu/kg/day as copper gluconate in drinking water (Massie and Aiello 1984).

3.2.2.3 Immunological and Lymphoreticular Effects

Information on the immunotoxicity of copper following oral exposure is limited to two drinking water studies in which mice were exposed to several concentrations of copper sulfate for 8 weeks (Pocino et al. 1990) or copper chloride for 3–5 or 8–10 weeks (Pocino et al. 1991). In these studies, groups of mice underwent several tests to assess immune function: *in vitro* lymphoproliferative responses to *Escherichia coli* lipopolysaccharide (LPS), and concanavalin A (Con A), induction and evaluation of antibody response to sheep red blood cells, evaluation of autoantibody production, and induction and elicitation of delayed-type hypersensitivity response (only tested in the Pocino et al. 1991 study). At the lowest dose tested (13 mg Cu/kg/day as copper chloride), impaired cellular (proliferative response to LPS) and humoral (autoantibody production) immunity were observed. Impaired performance on the remaining immune function tests were observed at ≥ 26 mg Cu/kg/day as copper chloride (Pocino et al. 1991) or

3. HEALTH EFFECTS

≥ 24 mg Cu/kg/day as copper sulfate (Pocino et al. 1990). The LOAEL values from these studies are presented in Table 3-2 and Figure 3-2.

3.2.2.4 Neurological Effects

No studies were located regarding neurological effects in humans following oral exposure to copper.

No effects on spontaneous motor activity (assessed using an actophometer), learning ability (assessed using a pole climbing chamber), or relearning capacity and memory (assessed using a Y-maze) were observed in rats fed a diet containing 23 mg Cu/kg/day as copper sulfate (Murthy et al. 1981). This study found no alterations in brain dopamine or norepinephrine levels. De Vries et al. (1986) also did not find significant alterations in corpus striatal dopamine levels in rats exposed to 36 mg Cu/kg/day as copper sulfate in drinking water for 11 months. However, a 25% decrease in a dopamine metabolite, 3,4-dihydroxyphenylacetic acid, was found in the corpus striatum.

3.2.2.5 Reproductive Effects

No studies were located regarding reproductive effects in humans following oral exposure to copper.

Reproductive performance, as assessed by the length of gestation, number of kits whelped, and average kit weight, was not adversely affected in minks fed a diet containing 12 mg Cu/kg/day as copper sulfate (Aulerich et al. 1982). No other oral exposure studies examined reproductive function. The intermediate-duration study by NTP (1993) did not find any histological alterations or alterations in sperm morphology or vaginal cytology in male and female rats exposed to 66 and 68 mg Cu/kg/day, respectively, or in male and female mice exposed to 398 and 536 mg Cu/kg/day, respectively. The NOAEL values for reproductive effects are reported in Table 3-2 and plotted in Figure 3-2.

3.2.2.6 Developmental Effects

No studies were located regarding developmental effects of humans following oral exposure to copper.

There are limited data on the developmental toxicity of copper in experimental animals. Delayed growth and development were observed in the offspring of rats exposed to 130 mg Cu/kg/day as copper sulfate in

3. HEALTH EFFECTS

the diet for 7 weeks prior to mating and during gestation (Haddad et al. 1991). In 11.5-day-old embryos, significant decreases in mean somite number, crown-rump length, and yolk sac diameter were observed. In 21.5-day-old fetuses and newborns, delayed ossification was observed in the cervical and cauda vertebrae, sternum, metacarpals, forelimb phalanges, metatarsals, and hindlimb phalanges. Exposure of mouse dams to a higher dose, 208 mg Cu/kg/day as copper sulfate in the diet, resulted in decreased mean litter size and decreased fetal body weights; the statistical significance of these effects is not known (Lecyk 1980). No statistically significant alterations in newborn mortality or body weight were observed in the offspring of mink exposed to 13 mg Cu/kg/day as copper sulfate in the diet (Aulerich et al. 1982). There was a trend toward increased kit mortality between birth and 4 weeks of age in the offspring of mink exposed to 6 or 13 mg Cu/kg/day. The incidences were 12, 9, 19, 38, and 32% in the 1, 6, 3, 6, and 13 mg Cu/kg/day groups, respectively; the statistical significance of this effect was not reported. The NOAEL values and all reliable LOAEL values for developmental effects in each species are recorded in Table 3-2 and plotted in Figure 3-2.

3.2.2.7 Cancer

No studies were located regarding carcinogenic effects in humans following oral exposure to copper.

Several oral studies have examined the carcinogenicity of copper compounds in animals. These studies did not find increases in the occurrence of tumors in mice exposed to 86 mg Cu/kg/day as the pesticide, copper 8-hydroxyquinoline (BRL 1968), liver tumors in rats exposed to 130 mg Cu/kg/day as copper acetate (Kamamoto et al. 1973), or large intestine tumors in rats exposed to 9 mg Cu/kg/day as an unspecified copper compound (Greene et al. 1987). These studies are limited in scope and it can not be determined whether the maximum threshold dose (MTD) was achieved. An increased occurrence of hepatocellular carcinomas has been reported in Long-Evans Cinnamon rats (Sawaki et al. 1994), an animal model for Wilson's disease. However, liver cancer has not been reported in individuals with Wilson's disease; thus the significance of this finding is not known.

3.2.3 Dermal Exposure

3.2.3.1 Death

No studies were located regarding death in humans and animals following dermal exposure to copper.

3. HEALTH EFFECTS

3.2.3.2 Systemic Effects

No studies were located regarding respiratory, cardiovascular, gastrointestinal, musculoskeletal, hepatic, renal, endocrine, dermal, or body weight effects in humans or animals following dermal exposure to copper.

Hematological Effects. Hemolytic anemia was observed in a severely burned and debilitated child in whom copper sulfate crystals were being applied to granulation tissue. Increased serum and urine copper levels were observed (Holtzman et al. 1966). Because the skin was severely damaged, this study cannot be used to predict the dermal toxicity of copper following exposure to intact skin. No studies were located regarding hematological effects in animals following dermal exposure to copper.

Ocular Effects. Eye irritation has been reported by factory workers exposed to copper dust (Askergren and Mellgren 1975). No studies were located regarding ocular effects in animals following exposure to copper.

3.2.3.3 Immunological and Lymphoreticular Effects

In some individuals, exposure to copper metal produces pruritic dermatitis. Saltzer and Wilson (1968) reported a case of a woman who had recurrent pruritus on her ring finger and wrist caused by copper metal in her ring and wristwatch. Allergic contact dermatitis has been observed in individuals following a patch test using a copper penny and/or a copper sulfate solution (Barranco 1972; Saltzer and Wilson 1968).

No studies were located regarding the following health effects in humans and/or animals after dermal exposure to copper:

3.2.3.4 Neurological Effects

3.2.3.5 Reproductive Effects

3.2.3.6 Developmental Effects

3.2.3.7 Cancer

3. HEALTH EFFECTS

3.2.4 Other Routes of Exposure

Cardiovascular Effects. A dramatic decrease in pulse pressure and heart rate was observed in New Zealand white rabbits infused with 2.5 mg Cu/kg as copper sulfate with an infusion pump in the femoral vein for 10–15 minutes (Rhee and Dunlap 1990). Systolic and diastolic pressure initially increased, then rapidly decreased.

Reproductive Effects. Intraperitoneal exposure to 0.95 or 1.4 mg Cu/kg/day for 26 days resulted in significant decreases in testes, seminal vesicle, and ventral prostate weights and in plasma testosterone levels in Wistar rats (Chattopadhyay et al. 1999); decreases in testicular $\Delta 5$ - 3β -hydroxysteroid dehydrogenases and 17β -hydroxysteroid dehydrogenase activities were also observed at 1.4 mg Cu/kg/day. An *in vitro* study (Holland and White 1988) demonstrated that cupric ions and cuprous ions decrease human spermatozoa motility.

Cancer. Several studies have examined the carcinogenicity of copper compounds following parenteral administration. No significant alterations in tumor incidence were observed in male Wistar rats receiving subcutaneous injections of 2 mg Cu/kg/day as copper acetate (Yamane et al. 1984), in male and female F344 rats receiving intramuscular injections of 0.25 or 0.41 mg Cu/kg/day as finely ground copper (Furst 1971), or in Wistar rats receiving 150 mg Cu/kg as copper oxide, 150 mg Cu/kg as copper sulfide, or 70 mg Cu/kg as copper sulfate (Gilman 1962). An increase in the occurrence of renal cell carcinoma was observed in male Wistar rats receiving 3–5 mg Cu/kg as cupric nitrilotriacetate 5 days/week for 12 weeks (Toyokuni et al. 1996); cupric nitrilotriacetate is a chelated compound of copper that is water soluble. A study by BRL (1968) found a slight, but statistically significant, increase in the incidence of reticulum cell sarcomas in mice 18 months after receiving a single subcutaneous injection of copper 8-hydroxyquinoline; the significance of this finding is not known.

3.3 GENOTOXICITY

No studies were located regarding genotoxicity in humans after inhalation, oral, or dermal exposure to copper or its compounds. Several studies have assessed the genotoxicity of copper sulfate following oral or parenteral exposure; the results of these *in vivo* genotoxicity studies are summarized in Table 3-3. Significant increases in the occurrence of micronuclei and chromosomal aberrations have been observed in chick bone marrow cells and erythrocytes (Bhunya and Jena 1996) and mouse bone marrow cells (Agarwal et al. 1990; Bhunya and Pati 1987). A study by Tinswell and Ashby (1990) did not find

3. HEALTH EFFECTS

Table 3-3. Genotoxicity of Copper *In Vivo*

Species (test system)	End point	Results	Reference	Compound
<i>Drosophila melanogaster</i> (injection into larvae)	Recessive lethals	+	Law 1938	Copper sulfate
White Leghorn chick bone marrow cells (intraperitoneal injection and oral exposure)	Chromosomal aberrations	+	Bhunya and Jena 1996	Copper sulfate
White Leghorn chick bone marrow cells (intraperitoneal injection and oral exposure)	Micronuclei	+	Bhunya and Jena 1996	Copper sulfate
White Leghorn chick erythrocytes (intraperitoneal injection and oral exposure)	Micronuclei	+	Bhunya and Jena 1996	Copper sulfate
Inbred Swiss mice bone marrow cells (intraperitoneal and/or subcutaneous injection)	Chromosomal aberrations	+	Bhunya and Pati 1987	Copper sulfate
Inbred Swiss mice bone marrow cells (intraperitoneal and/or subcutaneous injection)	Micronuclei	+	Bhunya and Pati 1987	Copper sulfate
Inbred Swiss mice (intraperitoneal injection)	Sperm abnormalities	+	Bhunya and Pati 1987	Copper sulfate
CBA mice bone marrow cells (intraperitoneal injection)	Micronuclei	-	Tinwell and Ashby 1990	Copper sulfate
Swiss mice (intraperitoneal injection)	Chromosomal aberrations	+	Agarwal et al. 1990	Copper sulfate

+ = positive results; - = negative results

3. HEALTH EFFECTS

increases in the number of micronuclei in mouse bone marrow cells. Increases in the occurrence of recessive lethals (Law 1938) and sperm abnormalities (Bhunya and Pati 1987) have also been observed in *Drosophila* and mice, respectively.

Several studies copper sulfate and copper chloride genotoxicity did not find significant increases in the occurrence of reverse mutations in *Salmonella typhimurium* (Marzin and Phi 1985; Tso and Fung 1981; Wong 1988) or *Saccharomyces cerevisiae* (Singh 1983). In contrast, Demerec et al. (1951) found an increased occurrence of reverse mutations in *Escherichia coli*. Positive results have been found in studies testing for DNA damage. Errors in DNA synthesis by viral DNA polymerase (Sirover and Loeb 1976), a reduction in DNA synthesis (Garrett and Lewtas 1983; Sirover and Loeb 1976), and an increase in the occurrence of DNA strand breaks (Sideris et al. 1988; Sina et al. 1983) have been observed. The increase in sister chromatid exchange in Chinese hamster cells (Sideris et al. 1988) is consistent with the clastogenic effects observed in *in vivo* assays. The results of these studies are summarized and are presented in Table 3-4.

3.4 TOXICOKINETICS

Physiologically normal levels of copper in the body are held constant by alterations in the rate and amount of copper absorption, compartmental distribution, and excretion.

3.4.1 Absorption

3.4.1.1 Inhalation Exposure

No studies were located regarding the rate and extent of absorption following inhalation exposure of humans to copper. There are limited data on copper absorption in animals. Copper oxide was observed in alveolar capillaries 3 hours after albino rats were exposed to a welding dust aerosol generated from pure copper wires (no additional exposure information was provided) (Batsura 1969). The half-time of copper sulfate in the lungs was estimated to be 7.5 hours after intratracheal instillation of 20 µg copper per Wistar rat (Hirano et al. 1990).

3. HEALTH EFFECTS

Table 3-4. Genotoxicity of Copper *In Vitro*

Species (test system)	End point	Results		Reference	Compound
		With activation	Without activation		
Prokaryotic organisms:					
<i>Salmonella typhimurium</i> TA102	Reverse mutation	NT	–	Marzin and Phi 1985	Copper sulfate
<i>S. typhimurium</i> TA98, TA102, TA1535, TA1537	Reverse mutation	–	–	Wong 1988	Copper chloride
<i>S. typhimurium</i> TA100	Reverse mutation	NT	–	Tso and Fung 1981	Copper chloride
<i>Escherichia coli</i>	Reverse mutation	NT	+	Demerec et al. 1951	Copper sulfate
Avian myeloblastosis virus, DNA polymerase	Errors in DNA synthesis	NT	+	Sirover and Loeb 1976	Copper chloride
<i>Bacillus subtilis</i>	rec assay	NT	–	Nishioka 1975	Copper chloride
Eukaryotic organisms:					
Fungi:					
<i>Saccharomyces cerevisiae</i>	Reverse mutation	NT	–	Singh 1983	Copper sulfate
<i>S. cerevisiae</i>	Recombination	NT	–	Sora et al. 1986	Copper sulfate
Mammalian cells:					
Chinese hamster ovary cells	DNA synthesis	NT	+	Garrett and Lewtas 1983	Copper chloride
Rat hepatocytes	DNA strand breaks	NT	+	Sina et al. 1983	Copper sulfate
Chinese hamster V79 cells	DNA strand breaks	NT	+	Sideris et al. 1988	Copper nitrate
Chinese hamster V79 cells	Sister chromatid exchange	NT	+	Sideris et al. 1988	Copper nitrate

+ = positive results; – = negative results; DNA = deoxyribonucleic acid; NT = not tested

3. HEALTH EFFECTS

3.4.1.2 Oral Exposure

Copper is absorbed from the stomach and small intestine; there appear to be species differences in the site of maximal absorption. The site of maximal copper absorption is not known for humans, but it is assumed to be the stomach and duodenum because of the rapid appearance of ^{64}Cu in the plasma after oral administration (Bearn and Kunkel 1955). In rats, copper is primarily absorbed from the duodenum, and to a lesser extent from the stomach (Van Campen and Mitchell 1965); in Golden hamsters, copper is primarily absorbed from the lower small intestine (25–35 cm from the pylorus) (Crampton et al. 1965).

Copper is absorbed from the gastrointestinal tract as ionic copper or bound to amino acids. Absorption of the latter apparently involves at least two kinetically distinguishable processes. The first mechanism transports copper from the mucosal side of the intestine to the serosal side. Only a small fraction of the ingested copper is transported via this mechanism (Crampton et al. 1965; Gitlan et al. 1960). The second mechanism of copper absorption involves the delivery of copper to the absorptive surface, mucosal uptake and binding to metallothionein or another intestinal binding protein (Evans and LeBlanc 1976). The copper bound to metallothionein can be slowly released to the blood (Marceau et al. 1970) or is excreted when the mucosal cell is sloughed off.

A number of human studies have examined the oral absorption of ^{64}Cu ; the average absorption efficiencies ranged from 24 to 60% in presumably healthy adults (Jacob et al. 1987; Johnson et al. 1988b; Strickland et al. 1972; Turnlund et al. 1982, 1983, 1985, 1988a; 1988b; 1989; Weber et al. 1969).

Numerous factors may affect copper absorption. These factors include: the amount of copper in the diet (Farrer and Mistilis 1967; Strickland et al. 1972; Turnland et al. 1989), competition with other metals, including zinc, iron, and cadmium (Davies and Campbell 1977; Hall et al. 1979; Haschke et al. 1986; Hoogenraad et al. 1979; Prasad et al. 1978; Turnland et al. 1988a) and age (Varada et al. 1993). The absorption of copper appears to be inversely related to the amount of copper in the gastrointestinal tract (Strickland et al. 1972; Turnland et al. 1989). In a study of 11 young men administered various copper doses in food over a period of 42–98 days, absorption efficiencies of 55–56, 36, and 12% were found at doses of 0.785, 1.68, and 7.53 mg/day, respectively (Turnland et al. 1989). In humans, the amount of stored copper does not appear to influence copper absorption (Strickland et al. 1972). In rats, the absorption of copper appears to be inversely related to the amount of cadmium in the diet (Davies and Campbell 1977). A significant decrease in copper absorption was observed when the copper:cadmium

3. HEALTH EFFECTS

ratio was 1:4. The amount of copper retained in the intestinal mucosal cells was also inversely related to cadmium dietary concentration. In addition, increased levels of zinc in the diet result in decreased in copper absorption in humans and rats (Hall et al. 1979; Hoogenraad et al. 1979; Prasad et al. 1978). Turnland et al. (1988a) found that diets low in zinc (below the dietary requirement) decreased copper absorption in humans; 48.1% of radiolabeled copper was absorbed when the diet contained 1.3 mg copper and 16.5 mg zinc (dietary requirement is 15 mg zinc), and 37.2–38.5% of radiolabelled copper was absorbed when the diet contained 1.3 mg copper and 5.5 mg zinc. A decrease in copper absorption has been observed in infants with high intakes of iron (Haschke et al. 1986). Apparently conflicting results have been reported on the effect of ascorbic acid on copper absorption in humans. Based on a decrease in serum ceruloplasmin levels, Finley and Cerklewski (1983) concluded that a diet high in ascorbic acid resulted in a decrease in copper status. In a study by Jacob et al. (1987), copper absorption was not affected by a high ascorbic acid intake. A decrease in serum ceruloplasmin activity was also found; however, the amount of ceruloplasmin protein was not affected.

Studies in humans and animals provide suggestive evidence of age-related changes in copper absorption. Varada et al. (1993) found that copper absorption was linear and nonsaturable in suckling (16 days of age) and weanling (21–22 days of age) rats. In contrast, copper absorption was saturable in adolescent rats (6 weeks of age). The levels of copper retained in the intestine were greater in the suckling rats than in the weanling or adolescent rats. However, the increased retention was not related to increased metallothionein levels; the levels of metallothionein (after zinc induction) were higher in the adolescent rats compared to the younger rats. A linear relationship between copper intake and retention was also found in a balance study of infants (aged 2–16 weeks) (Dörner et al. 1989). Olivares et al. (2002) did not find significant differences in copper absorption between 1-month-old and 3-month-old infants. The relatively small range of doses used in this study does not allow for a determination of whether copper absorption is saturable in infants. Several studies of adults did not find differences in copper absorption between male and female adults aged 20–83 years (Johnson et al. 1992) or between elderly men (65–74 years) and young men (22–30 years) (Turnland et al. 1982, 1988b).

Human studies did not find that increased levels of fiber (α -cellulose or phytate) (Turnland et al. 1985) or ascorbic acid (Turnland et al. 1987) significantly altered copper absorption. However, a study in rats found an increase in fecal excretion of copper (and a decrease in apparent absorption) in rats fed a high fiber (potato fiber or sugar beet pulp) diet (Gralak et al. 1996). The administration of copper in infant formula or in a solution high in fulvic acid did not appear to influence copper uptake from the intestinal lumen into the intestinal mucosa of suckling rats, as compared to copper in drinking water (Lind and

3. HEALTH EFFECTS

Glynn 1999). However, the absorption rate of copper into the circulatory system was decreased when administered in the infant formula or fulvic acid solutions. Gender does not appear to influence copper absorption. Johnson et al. (1992) found that women aged 20–59 years absorbed more copper (66.1–74.1%) than similarly aged men (62.0–69.2%); however, when net copper absorption was normalized by body weight, no sex-related differences in absorption were found. No sex-related differences in net copper absorption were found in older (60–83 years) men and women.

3.4.1.3 Dermal Exposure

The available *in vivo* data do not provide information on the rate and extent of absorption through intact skin following dermal exposure of humans or animals to copper. Following a copper azide explosion that yielded metallic copper and nitrogen fumes, a small increase in serum copper levels was found in the affected worker (Bentur et al. 1988). Similarly, animal studies demonstrate that copper can pass through dermal barriers when applied with an appropriate vehicle, (e.g., salicylic acid or phenylbutazone) (Beveridge et al. 1984; Walker et al. 1977). *In vitro* studies suggest that copper is poorly absorbed through intact skin. Less than 6% of copper deposited on *ex vivo* human skin samples was absorbed (Pirot et al. 1996a, 1996b); copper chloride was absorbed to a higher extent than copper sulfate (Pirot et al. 1996a).

3.4.2 Distribution

3.4.2.1 Inhalation Exposure

No studies were located regarding the rate and extent of distribution of copper following inhalation exposure of humans or animals.

3.4.2.2 Oral Exposure

Following ingestion of copper, copper levels in the blood rapidly rise. The copper is predominantly bound to albumin. There is some evidence that albumin plays a passive role in copper transport, carrying a large portion of the exchangeable copper in the circulation and releasing this to other carriers for actual cell-specific uptake. There is also evidence that transcuprein is another plasma protein carrier (Weiss and Linder 1985). Thus, dietary copper is transported to, and enters, the liver and kidney. Copper then reemerges into the plasma bound to the ceruloplasmin. Ceruloplasmin, which tightly binds six or seven

3. HEALTH EFFECTS

copper atoms (Musci et al. 1993; Saenko et al. 1994), is the most abundant copper protein in the plasma; 60–95% of the plasma copper is bound to ceruloplasmin (Harris 1993). Copper is transported from the liver to other tissues via ceruloplasmin. Ceruloplasmin does not enter the cell (Percival and Harris 1990). Copper, probably as Cu(I) rather than Cu(II) (Dameron and Harris 1989; Percival and Harris 1989), enters the cell via a carrier-mediated process. The membrane-bound copper transporting adenosine triphosphatase (Cu-ATPase), which selectively binds copper ions, transports copper ions into and out of cells (Harris et al. 1998). In most organs and tissues, copper turnover is biphasic (Levenson and Janghorbani 1994). In the plasma, the half-lives of the first and second components were 2.5 and 69 days, respectively. It is likely that the first order component is ceruloplasmin associated copper. The respective calculated copper half-lives for other tissues are 3.9 and 21 days for the liver, 5.4 and 35 days for the kidney, and 23 and 662 days for the heart; copper turnover in the brain appears to be monophasic with a half-life of 457 days.

3.4.2.3 Dermal Exposure

No studies were located regarding the rate and extent of distribution of copper following dermal exposure of humans or animals to copper.

3.4.3 Metabolism

The metabolism of copper consists mainly of its transfer to and from various organic ligands, most notably sulfhydryl and imidazole groups on amino acids and proteins. Several specific binding proteins for copper have been identified that are important in the uptake, storage, and release of copper from tissues.

In the liver and other tissues, copper is stored bound to metallothionein and amino acids and in association with copper-dependent enzymes. Several studies have shown that copper exposure induces metallothionein synthesis (Mercer et al. 1981; Wake and Mercer 1985). Increased levels of metallothionein may be associated with resistance to copper toxicity in pigs (Mehra and Bremner 1984). Ceruloplasmin is synthesized in the liver. Copper is incorporated into the molecule, and it is released from the liver. Copper exposure has also been shown to induce ceruloplasmin biosynthesis (Evans et al. 1970b; Haywood and Comerford 1980).

3. HEALTH EFFECTS

3.4.4 Elimination and Excretion

3.4.4.1 Inhalation Exposure

No studies were located regarding the rate and extent of excretion of copper following inhalation exposure of humans or animals.

3.4.4.2 Oral Exposure

Bile is the major pathway for the excretion of copper. After the oral administration of radioactive copper as copper acetate in healthy humans, 72% was excreted in the feces (Bush et al. 1955). A considerable fraction of fecal copper is of endogenous biliary origin. The remainder of the fecal copper is derived from unabsorbed copper and copper from desquamated mucosal cells. Copper in bile is associated with low molecular weight copper binding components as well as macromolecular binding species (Gollan and Dellar 1973). Reabsorption of biliary copper is negligible (Farrer and Mistilis 1967).

Normally, 0.5–3.0% of daily copper intake is excreted into the urine (Cartwright and Wintrobe 1964).

3.4.4.3 Dermal Exposure

No studies were located regarding the rate and extent of excretion of copper following dermal exposure of humans or animals to copper.

3.4.4.4 Other Routes of Exposure

Biliary excretion of copper following intravenous administration does not increase proportionally with dosage, suggesting that the hepatobiliary transport of copper is saturable (Gregus and Klaassen 1986). Thus, at high copper intakes, urinary copper excretion increases (Gitlan et al. 1960).

3.4.5 Physiologically Based Pharmacokinetic (PBPK)/Pharmacodynamic (PD) Models

Physiologically based pharmacokinetic (PBPK) models use mathematical descriptions of the uptake and disposition of chemical substances to quantitatively describe the relationships among critical biological

3. HEALTH EFFECTS

processes (Krishnan et al. 1994). PBPK models are also called biologically based tissue dosimetry models. PBPK models are increasingly used in risk assessments, primarily to predict the concentration of potentially toxic moieties of a chemical that will be delivered to any given target tissue following various combinations of route, dose level, and test species (Clewell and Andersen 1985). Physiologically based pharmacodynamic (PBPD) models use mathematical descriptions of the dose-response function to quantitatively describe the relationship between target tissue dose and toxic end points.

PBPK/PD models refine our understanding of complex quantitative dose behaviors by helping to delineate and characterize the relationships between: (1) the external/exposure concentration and target tissue dose of the toxic moiety, and (2) the target tissue dose and observed responses (Andersen et al. 1987; Andersen and Krishnan 1994). These models are biologically and mechanistically based and can be used to extrapolate the pharmacokinetic behavior of chemical substances from high to low dose, from route to route, between species, and between subpopulations within a species. The biological basis of PBPK models results in more meaningful extrapolations than those generated with the more conventional use of uncertainty factors.

The PBPK model for a chemical substance is developed in four interconnected steps: (1) model representation, (2) model parametrization, (3) model simulation, and (4) model validation (Krishnan and Andersen 1994). In the early 1990s, validated PBPK models were developed for a number of toxicologically important chemical substances, both volatile and nonvolatile (Krishnan and Andersen 1994; Leung 1993). PBPK models for a particular substance require estimates of the chemical substance-specific physicochemical parameters, and species-specific physiological and biological parameters. The numerical estimates of these model parameters are incorporated within a set of differential and algebraic equations that describe the pharmacokinetic processes. Solving these differential and algebraic equations provides the predictions of tissue dose. Computers then provide process simulations based on these solutions.

The structure and mathematical expressions used in PBPK models significantly simplify the true complexities of biological systems. If the uptake and disposition of the chemical substance(s) are adequately described, however, this simplification is desirable because data are often unavailable for many biological processes. A simplified scheme reduces the magnitude of cumulative uncertainty. The adequacy of the model is, therefore, of great importance, and model validation is essential to the use of PBPK models in risk assessment.

3. HEALTH EFFECTS

PBPK models improve the pharmacokinetic extrapolations used in risk assessments that identify the maximal (i.e., the safe) levels for human exposure to chemical substances (Andersen and Krishnan 1994). PBPK models provide a scientifically sound means to predict the target tissue dose of chemicals in humans who are exposed to environmental levels (for example, levels that might occur at hazardous waste sites) based on the results of studies where doses were higher or were administered in different species. Figure 3-3 shows a conceptualized representation of a PBPK model.

A PBPK model for copper has not been identified.


3.5 MECHANISMS OF ACTION

3.5.1 Pharmacokinetic Mechanisms

Copper is an essential element required for the normal functioning of more than 30 enzymes. The ability of copper to cycle between an oxidized state, Cu(II), and reduced state, Cu(I), is used by cuproenzymes involved in redox reactions. However, it is this property of copper that is also potentially toxic because the transitions between Cu(II) and Cu(I) can result in the generation of superoxide radicals and hydroxyl radicals (Camakaris et al. 1999). Under most circumstances, a number of homeostatic mechanisms maintain a physiologically essential concentration of copper. Copper homeostasis involves regulation of absorption, cellular uptake, intracellular transport, sequestration/storage, cellular efflux, and excretion from the body. Turnland et al. (1989) demonstrated that copper absorption from the gastrointestinal tract is inversely proportional to dietary intake; as dietary copper increases, absorption efficiency decreases. At dietary concentrations of 0.785, 1.68, and 7.53 mg/day (the recommended dietary allowance [RDA] for copper is 0.900 mg/day), 56, 36, and 12%, respectively, of the radiolabelled copper was absorbed. How the absorption of copper is regulated is not fully understood. *In vitro* studies provide evidence that copper uptake into intestinal cells appears to be saturable (Arredondo et al. 2000). This study also provides suggestive evidence that copper uptake into the intestinal cell and efflux are influenced by intracellular copper concentrations. There is evidence that copper diffuses across the intestinal cell membrane; however, it is unlikely that this is the only absorption mechanism. It is possible that recently identified copper transporters (hCtr1 and hCtr2) play a role in the regulation of copper uptake. The Menkes protein (MNK), a copper-translocating P-type ATPase, may be involved in the transport of copper across the basolateral membrane of intestinal cells (Pena et al. 1999). MNK protein is involved the delivery of copper to copper-dependent enzymes and the efflux of copper from the cell. The export of

3. HEALTH EFFECTS

Figure 3-3. Conceptual Representation of a Physiologically Based Pharmacokinetic (PBPK) Model for a Hypothetical Chemical Substance



Note: This is a conceptual representation of a physiologically based pharmacokinetic (PBPK) model for a hypothetical chemical substance. The chemical substance is shown to be absorbed via the skin, by inhalation, or by ingestion, metabolized in the liver, and excreted in the urine or by exhalation.

Source: adapted from Krishnan et al. 1994

3. HEALTH EFFECTS

copper via the MNK protein appears to be regulated by intracellular copper concentration. Exposure to copper produces a conformational change in the MNK protein resulting in the formation of a copper cluster, which allows access to the phosphorylation site that upon phosphorylation, initiates copper translocation (Dameron and Harrison 1998). Once copper is released from the intestinal cells, it is transported bound to albumin and histidine to the liver via the portal circulation. Once in the hepatic cells, copper complexes with small cytoplasmic proteins known as copper chaperones. These copper chaperones are involved in intracellular distribution of copper ions. In the liver, another P-type ATPase, Wilson protein (WND), delivers copper to ceruloplasmin, which is then released to the blood for distribution to other tissues and organs. Under conditions of elevated copper, WND is involved in the release of copper at the canalicular membrane with ensuing biliary excretion of copper. The liver plays a critical role in copper homeostasis as both the storage site for this metal and as part of the physiologic route for excretion through the biliary system. The molecular mechanisms determining biliary copper excretion are becoming clearer due to the better understanding of genetic defects, such as Wilson's disease. Specifically, the Wilson protein localized to the trans-Golgi network of hepatocytes not only delivers copper to ceruloplasmin, but also is essential for biliary copper excretion. Recently, other proteins have also been identified that interact with Wilson protein and appear to be equally important in the process of biliary copper excretion (Tao et al. 2003).

3.5.2 Mechanisms of Toxicity

Although a number of studies have investigated the mechanisms of copper hepatotoxicity in rats, it is not known whether rats would be a good model for human liver toxicity unrelated to a genetic defect in copper metabolism. Lysosomes serve an important role in hepatic copper metabolism. Excess copper is sequestered within hepatocyte lysosomes where it is complexed with metallothionein. However, this protective mechanism is saturable and liver lesions can develop above the saturation limit. In copper loaded rats, lysosomes become enlarged and more fragile with decreased membrane fluidity (Myers et al. 1993). The results of the Haywood et al. (1985a) study do not suggest that liver damage is due to rupturing of lysosomes because lysosomal instability precedes and is not synchronous with liver damage. It is speculated that saturation of the lysosomes results in an accumulation of copper in the nucleus and subsequent nuclear damage (Fuentelba and Haywood 1988; Fuentelba et al. 1989; Haywood et al. 1985a). The mechanism by which copper accumulates in the nucleus and the mechanisms by which it provokes injury are not clear. It has been suggested that excess copper results in oxidative damage, including lipid peroxidation. Increases in the level of thiobarbituric acid reactive substance (TBARS), a measure of lipid peroxidation, have been found in copper-loaded rats (Myers et al. 1993; Sokol et al.

3. HEALTH EFFECTS

1993). However, a study by Aburto et al. (2001b) did not find significant alterations in the levels of malondialdehyde, a lipid peroxidation by product, prompting the study authors to postulate that lipid peroxidation does not play a major role in copper toxicity although it may occur as a terminal event as a consequence of cell injury. Sokol et al. (1990, 1993) suggested that oxidant injury to hepatocyte mitochondria may be one of the initiating factors in hepatocellular damage. Numerous studies have shown that rats can develop tolerance to high levels of copper. After 3–5 weeks of copper loading resulting in tissue damage, the copper levels in the liver begin to decline and the tissue begins to regenerate (Haywood and Loughran 1985). It is believed that the mechanism involved in tolerance development is the increased synthesis of metallothionein (Evering et al. 1991a, 1991b; Freedman and Peisach 1989).

Studies in monkeys, dogs, and ferrets provide strong evidence that copper-induced emesis results from stimulation of the vagus nerve. Abdominal vagotomy resulted in a dramatic decrease in the occurrence of emesis in dogs (Fukui et al. 1994) and ferrets (Makale and King 1992) orally exposed to copper sulfate and in monkeys receiving oral or intravenous injections of copper sulfate (Fukui et al. 1993). In monkeys, administration of compounds that block 5-HT₃ receptors also resulted in a decrease in emesis following oral or intravenous administration of copper sulfate (Fukui et al. 1993). In contrast, 5-HT₃ blockers did not affect the occurrence of emesis in dogs (Fukui et al. 1994) or ferrets (Bhandari and Andrew 1991) receiving an oral dose of copper sulfate, but compounds that block 5-HT₄ receptors did inhibit copper-induced vomiting. Fukui et al. (1994) suggested that copper sulfate caused gastrointestinal irritation that resulted in the release of 5-HT and evoked emesis by activation of abdominal visceral afferents through 5-HT₄ receptors.

3.5.3 Animal-to-Human Extrapolations

The toxicity of copper has been assessed in a number of experimental animal species, and sensitivity to copper toxicity is highly species dependent. Ruminants are more susceptible than nonruminant species. NTP (1993) demonstrated that rats are much more sensitive than mice to the hepatotoxicity of copper. In rats, dietary exposure to 16 mg Cu/kg/day for 13 weeks resulted in an increase in alanine aminotransferase activity; chronic active liver inflammation was observed at 66 mg Cu/kg/day. In contrast, no evidence of liver damage was observed in mice exposed to 814 mg Cu/kg/day for 13 weeks.

Most of the experimental data on the toxicity of copper come from studies in which rats were used; however, the relevance of this species to human toxicity has not been fully evaluated. The dietary

3. HEALTH EFFECTS

requirement for copper in rats is 5 mg Cu/kg diet (NRC 1995); a commonly used diet for rats (AIN76, AIN 93G, AIN93M) has a cupric carbonate concentration of 300 mg/kg diet (160 mg Cu/kg diet). An intermediate-duration exposure to approximately 250 mg Cu/kg diet resulted in mild liver effects (increased serum alanine aminotransferase) (NTP 1993). It is unlikely that humans would tolerate prolonged exposure to a copper dose that is 50 times higher than the dietary requirement (0.65 mg Cu/kg/day); gastrointestinal disturbances were observed in women ingesting 0.0731 mg Cu/kg/day in drinking water (Pizarro et al. 1999). Thus, the applicability of these animal data to humans is not known.

The Long-Evans Cinnamon rat is often used as a model for Wilson's disease. This rat strain shares many characteristics with Wilson's disease: accumulation of liver copper, decreased serum copper and ceruloplasmin levels, and impaired biliary excretion of copper (Sugawara et al. 1991, 1992, 1994; Suzuki et al. 1995).

3.6 TOXICITIES MEDIATED THROUGH THE NEUROENDOCRINE AXIS

Recently, attention has focused on the potential hazardous effects of certain chemicals on the endocrine system because of the ability of these chemicals to mimic or block endogenous hormones. Chemicals with this type of activity are most commonly referred to as *endocrine disruptors*. However, appropriate terminology to describe such effects remains controversial. The terminology *endocrine disruptors*, initially used by Colborn and Clement (1992), was also used in 1996 when Congress mandated the EPA to develop a screening program for "...certain substances [which] may have an effect produced by a naturally occurring estrogen, or other such endocrine effect[s]...". To meet this mandate, EPA convened a panel called the Endocrine Disruptors Screening and Testing Advisory Committee (EDSTAC), and in 1998, the EDSTAC completed deliberations and made recommendations to EPA concerning *endocrine disruptors*. In 1999, the National Academy of Sciences released a report that referred to these same types of chemicals as *hormonally active agents*. The terminology *endocrine modulators* has also been used to convey the fact that effects caused by such chemicals may not necessarily be adverse. Many scientists agree that chemicals with the ability to disrupt or modulate the endocrine system are a potential threat to the health of humans, aquatic animals, and wildlife. However, others believe that endocrine-active chemicals do not pose a significant health risk, particularly in view of the fact that hormone mimics exist in the natural environment. Examples of natural hormone mimics include the isoflavonoid phytoestrogens (Adlercreutz 1995; Livingston 1978; Mayr et al. 1992). These chemicals are derived from plants and are similar in structure and action to endogenous estrogen. Although the public health significance and descriptive terminology of substances capable of affecting the endocrine system remains controversial,

3. HEALTH EFFECTS

scientists agree that these chemicals may affect the synthesis, secretion, transport, binding, action, or elimination of natural hormones in the body responsible for maintaining homeostasis, reproduction, development, and/or behavior (EPA 1997). Stated differently, such compounds may cause toxicities that are mediated through the neuroendocrine axis. As a result, these chemicals may play a role in altering, for example, metabolic, sexual, immune, and neurobehavioral function. Such chemicals are also thought to be involved in inducing breast, testicular, and prostate cancers, as well as endometriosis (Berger 1994; Giwercman et al. 1993; Hoel et al. 1992). However, in the case of ambient human exposures, the validity of these possibilities has yet to be established conclusively.

There is no evidence that copper interferes with the normal function of the neuroendocrine axis.

3.7 CHILDREN'S SUSCEPTIBILITY

This section discusses potential health effects to humans from exposures during the period from conception to maturity at 18 years of age, when all biological systems will have fully developed. Potential effects on offspring resulting from exposures of parental germ cells are considered, as well as any direct or indirect effects on the fetus and neonate resulting from maternal exposure during gestation and lactation. Potentially relevant animal and *in vitro* models are also discussed.

Children should not be considered small adults. They may differ from adults in their exposures and in their susceptibility to hazardous chemicals. Children's unique physiology and behavior can influence the extent of their exposure. Exposures of children are discussed in Section 6.6 Exposures of Children.

Children may differ from adults in their susceptibility to hazardous chemicals, but whether there is a difference depends on the chemical (Guzelian et al. 1992; NRC 1993). Children may be more or less susceptible than adults to health effects, and the relationship may change with developmental age (Guzelian et al. 1992; NRC 1993). Vulnerability often depends on developmental stage. There are critical periods of structural and functional development during both prenatal and postnatal life at which particular structures and/or functions will be most sensitive to perturbation. Damage may not be evident until a later stage of development. There are often differences in pharmacokinetics and metabolism between children and adults. For example, absorption may be different in neonates because of the immaturity of their gastrointestinal tract and their larger skin surface area in proportion to body weight (Morselli et al. 1980; NRC 1993); the gastrointestinal absorption of lead is greatest in infants and young children (Ziegler et al. 1978). Also, the distribution of xenobiotics may be different; for example, infants

3. HEALTH EFFECTS

have a larger proportion of their bodies as extracellular water, their brains and livers are proportionately larger, and the composition and quality of their lipid depots differ from those of adults (Altman and Dittmer 1974; Fomon 1966; Fomon et al. 1982; Owen and Brozek 1966; Widdowson and Dickerson 1964). The infant also has an immature blood-brain barrier (Adinolfi 1985; Johanson 1980) and probably an immature blood-testis barrier (Setchell and Waites 1975). Many xenobiotic metabolizing enzymes have distinctive developmental patterns. At various stages of growth and development, levels of particular enzymes may be higher or lower than those of adults, and sometimes unique enzymes may exist at particular developmental stages (Komori et al. 1990; Leeder and Kearns 1997; NRC 1993; Vieira et al. 1996). Whether differences in xenobiotic metabolism make the child more or less susceptible also depends on whether the relevant enzymes are involved in activation of the parent compound to its toxic form and/or in detoxification. There may also be differences in excretion, particularly in newborns who have a low glomerular filtration rate and have not developed efficient tubular secretion and resorption capacities (Altman and Dittmer 1974; NRC 1993; West et al. 1948). In addition, children and adults may differ in their capacity to repair damage from chemical insults. In general, children also have a longer remaining lifetime in which to express damage from chemicals; this potential is particularly relevant to the development of cancer.

Certain characteristics of the developing human may increase exposure or susceptibility to certain toxicants, whereas others may decrease susceptibility to the same toxicant. For example, although infants breathe more air per kilogram of body weight than adults, this difference might be somewhat counterbalanced by their alveoli being less developed, which results in a disproportionately smaller surface area for alveolar absorption (NRC 1993).

Copper is an essential element required for normal growth and development and for a variety of metabolic functions including iron metabolism, cross-linking of connective tissue, and lipid metabolism. Signs of copper deficiency in infants and children include anemia that is unresponsive to iron supplementation, neutropenia, bone abnormalities, and hypopigmentation of the hair (Cordano 1998; Danks 1988).

Exposure to excess levels of copper has been associated with adverse health effects in infants and children. There is an extensive body of literature on two syndromes that have been associated with exposure to high levels of copper, Indian childhood cirrhosis and idiopathic copper toxicosis. Both are characterized by severe liver damage in infants and children (<5 years of age). In the case of Indian childhood cirrhosis, excessive copper exposure has been traced to the use of brass or copper containers for storage and heating of milk. Doses as high as 0.930 mg/kg/day have been estimated; this dose is

3. HEALTH EFFECTS

approximately 30 times higher than the dietary requirement for copper (Tanner 1998). Idiopathic copper toxicosis (also referred to as non-Indian childhood cirrhosis) has also been linked to exposure to high levels of copper in drinking water and/or the use of copper utensils (Wijemenga 2002). A common finding in both syndromes is the early dietary introduction of non-mother's milk and/or formula. Genealogical investigations provide suggestive evidence that both syndromes are transmitted in an autosomal recessive mode. However, the mechanism of action has not been identified. It is possible that the genetic defect results in reduced copper efflux from the liver. Very high levels of copper have been detected in the livers of affected infants; copper levels ranging from 790 to 6,654 $\mu\text{g/g}$ dry weight (mean of 939 $\mu\text{g/g}$) have been reported in infants diagnosed with Indian childhood cirrhosis (levels in control infants ranged from 8 to 118 $\mu\text{g/g}$) (Bhave et al. 1982). Support for the genetic component comes from the finding that decreasing copper exposure levels dramatically decreases the occurrence of Indian childhood cirrhosis (Tanner 1998). Additionally, no alterations in serum biomarkers of liver damage (alanine aminotransferase activity, aspartate aminotransferase activity, gamma glutamyl transferase activity, and total bilirubin levels) were observed in infants ingesting water containing 2 mg/L copper (0.319 mg/kg/day) (Olivares et al. 1998) or infants living in households with tap water copper levels of 0.8 mg/L (Zietz et al. 2003a, 2003b). Together, these data suggest that exposure to copper levels exceeding the copper metabolic capacity of certain individuals with a genetic defect is the causative agent for severe liver damage.

Another adverse health effect that has been reported in infants and children is gastrointestinal upset. This effect, which is one of the most commonly reported adverse health effect in adults, is manifested in nausea, vomiting, abdominal pain, and/or diarrhea. Symptoms usually occur shortly after ingesting a copper-contaminated beverage or drinking water containing a high level of copper. In most of the reports of gastrointestinal upset in children (Gill and Bhagat 1999; Karlsson and Noren 1965; Knobloch et al. 1994; Spitalny et al. 1984; Walsh et al. 1977), no reliable information on copper concentration or dose was reported. In one report where school-age children ingested a beverage stored in an old urn, the concentration of copper in the beverage was estimated to be 300 mg/L (Gill and Bhagat 1999). Another study reported vomiting in infants ingesting a single dose of 7.5 mg/L copper sulfate (Karlsson and Noren 1965). Knobloch et al. (1994) noted that children appear to be more sensitive to the gastrointestinal effects of copper than adults. This statement was based on two surveys of residents with elevated copper levels in the drinking water. In the first survey, it appears that children who were described as "unusually irritable" or had recurrent headaches were categorized as having gastrointestinal upset. In the second survey, mothers were asked to recall the frequency of gastrointestinal effects for all family members. A significantly higher percentage of children, as compared to adults, were reported to have gastrointestinal

3. HEALTH EFFECTS

effects. It is difficult to determine what role recall bias had in the results and how well the mothers knew of symptoms of gastrointestinal upset in the adult members of the household. The available data are inconclusive to assess accurately whether there is an age-related difference in the gastrointestinal toxicity of copper.

The potential age-related differences in the toxicity of copper has been assessed in rats exposed to 120 mg Cu/kg/day as copper sulfate in the diet for 12 weeks (Fuentelba et al. 2000). The observed liver effects were more severe in young rats (exposed *in utero*, during lactation, and for 12 weeks post weaning) as compared to the effects observed in adult rats. The copper levels in the liver were also higher in the young rats (1,553–1,635 versus 472–534 µg/g). The doses used in this study are very high, 1,000 times higher than the rat dietary requirement of 0.15–0.30 mg/kg/day (AIN 1977). It is not known if increased liver sensitivity would also occur at lower copper doses. Although these data are suggestive that children may be more sensitive to the hepatotoxicity of high doses of copper, uncertainty in the use of rats as a model for human toxicity limits the extrapolation of these study results to humans.

Several studies have investigated the potential developmental toxicity of dietary copper sulfate; the results suggest that *in utero* exposure to copper can result in delays in growth and development in the offspring of rats exposed to 130 mg Cu/kg/day (Haddad et al. 1991) and mice exposed to 208 mg Cu/kg/day (Lecyk 1980). No developmental effects were observed in the offspring of mink exposed to 13 mg Cu/kg/day (Aulerich et al. 1982).

There is concern that toxicokinetic differences between infants and adults may result in increased sensitivity in infants. During the second half of pregnancy, particularly in the third trimester, the fetus accumulates copper at a rate of 50 g/kg/day (Widdowson et al. 1974). Approximately half of the copper in the fetus is stored in the liver, mostly bound to metallothionein. Additionally, the rate of transfer of copper from the liver to the bile or blood is decreased due to the immaturity of the liver. The magnitude of the amount of copper in the fetal liver is similar to levels observed in Wilson's disease; however, the fetal/neonatal liver tolerates these high concentrations (Olivares et al. 2000). After birth, the copper levels in the liver steadily decrease from about 51 µg/g at birth to 5.7 µg/g at 6–14 months of age (Klein et al. 1991; Olivares et al. 2000).

3. HEALTH EFFECTS

3.8 BIOMARKERS OF EXPOSURE AND EFFECT

Biomarkers are broadly defined as indicators signaling events in biologic systems or samples. They have been classified as markers of exposure, markers of effect, and markers of susceptibility (NAS/NRC 1989).

Due to a nascent understanding of the use and interpretation of biomarkers, implementation of biomarkers as tools of exposure in the general population is very limited. A biomarker of exposure is a xenobiotic substance or its metabolite(s) or the product of an interaction between a xenobiotic agent and some target molecule(s) or cell(s) that is measured within a compartment of an organism (NAS/NRC 1989). The preferred biomarkers of exposure are generally the substance itself or substance-specific metabolites in readily obtainable body fluid(s), or excreta. However, several factors can confound the use and interpretation of biomarkers of exposure. The body burden of a substance may be the result of exposures from more than one source. The substance being measured may be a metabolite of another xenobiotic substance (e.g., high urinary levels of phenol can result from exposure to several different aromatic compounds). Depending on the properties of the substance (e.g., biologic half-life) and environmental conditions (e.g., duration and route of exposure), the substance and all of its metabolites may have left the body by the time samples can be taken. It may be difficult to identify individuals exposed to hazardous substances that are commonly found in body tissues and fluids (e.g., essential mineral nutrients such as copper, zinc, and selenium). Biomarkers of exposure to copper are discussed in Section 3.8.1.

Biomarkers of effect are defined as any measurable biochemical, physiologic, or other alteration within an organism that, depending on magnitude, can be recognized as an established or potential health impairment or disease (NAS/NRC 1989). This definition encompasses biochemical or cellular signals of tissue dysfunction (e.g., increased liver enzyme activity or pathologic changes in female genital epithelial cells), as well as physiologic signs of dysfunction such as increased blood pressure or decreased lung capacity. Note that these markers are not often substance specific. They also may not be directly adverse, but can indicate potential health impairment (e.g., DNA adducts). Biomarkers of effects caused by copper are discussed in Section 3.8.2.

A biomarker of susceptibility is an indicator of an inherent or acquired limitation of an organism's ability to respond to the challenge of exposure to a specific xenobiotic substance. It can be an intrinsic genetic or other characteristic or a preexisting disease that may result in an increase in absorbed dose, a decrease in the dose-level required for biological effectiveness, or a target tissue response. Biomarkers of susceptibility are discussed in Section 3.10 "Populations that are Unusually Susceptible."

3. HEALTH EFFECTS

3.8.1 Biomarkers Used to Identify or Quantify Exposure to Copper

Copper levels can readily be measured in tissues, body fluids, and excreta. Depending on the dose and exposure duration, inhalation and/or oral exposure to copper can result in increased levels of copper in serum, urine, hair, and liver. Increased whole blood and serum copper levels have been reported in humans following intentional ingestion of a single dose of 1–30 g of copper as copper sulfate (Chuttani et al. 1965). The serum and whole blood levels of copper ranged from 239 to 346 and from 383 to 684 $\mu\text{g}/100\text{ mL}$, respectively; the serum and whole blood levels in non-exposed individuals were 151.6 and 217 $\mu\text{g}/100\text{ mL}$, respectively. Following chronic inhalation exposure to 111–464 $\text{mg Cu}/\text{m}^3$ copper dust, plasma serum levels of $>200\text{ }\mu\text{g}/100\text{ mL}$ were observed in 16% of factory workers exposed to copper dust (Suciu et al. 1981). However, increased serum copper levels may only be reflective of recent exposure. Chuttani et al. (1965) observed that serum ionic copper rapidly diminished to normal levels following an acute bolus dose.

A relationship between blood copper levels and the severity of symptoms has not been established. Among individuals intentionally ingesting a single dose of copper sulfate (1–30 g), Chuttani et al. (1965) noted that there did not appear to be any difference between serum copper levels in individuals only exhibiting gastrointestinal effects and those with more severe symptoms (jaundice, renal manifestations, or shock). In contrast, whole blood copper levels were much higher in the individuals with severe symptoms (798 $\mu\text{g}/100\text{ mL}$) compared to those with mild symptoms (287 $\mu\text{g}/100\text{ mL}$).

Copper levels in hair and nails can also be used to assess copper exposure. In a study of preschool children, the levels of copper in hair and toenail samples were log-normally distributed (Wilhelm et al. 1991). The geometric mean concentrations of copper in hair and toenails were 10.6 $\mu\text{g}/\text{g}$ (range of 5.4–20.7 $\mu\text{g}/\text{g}$) and 7.5 $\mu\text{g}/\text{g}$ (range of 3.0–18.6 $\mu\text{g}/\text{g}$), respectively. Based on a hair growth rate of 10 mm per month, the copper levels in the first 2 cm proximal to the scalp would represent copper intake over 2 months (Hopps 1977). In contrast, toenail samples would represent copper intake over 12–18 months, based on a toenail growth rate of 1 mm/month (Fleckman 1985). Increased hair copper levels have been reported in workers exposed to 0.64–1.05 mg/m^3 of an unspecified copper compound; the concentration of copper in the hair was 705.7 $\mu\text{g}/\text{g}$, as compared to a 8.9 $\mu\text{g}/\text{g}$ concentration in non-exposed workers (Finelli et al. 1981), and increased hair and fingernail copper levels were observed in children with Indian childhood cirrhosis (Sharda and Bhandari 1984).

3. HEALTH EFFECTS

3.8.2 Biomarkers Used to Characterize Effects Caused by Copper

The harmful health effects of copper occur over a wide range of copper intakes from too little copper in the diet to excessive copper exposure.

Low Intakes of Copper. The nutritional requirements of copper and the health effects associated with copper deficiency have been reviewed by numerous authors (Gallagher 1979; Mason 1979; O'Dell 1984). Copper deficiency is rarely observed in humans; the existence of covert copper deficiency among segments of the population is unknown. The limited data available on human health effects of inadequate copper intakes are derived mostly from case reports of severely malnourished children, patients maintained by total parenteral nutrition without copper, and children with Menkes' disease (a genetic disorder resulting in impaired copper absorption). Copper deficiency is characterized by hypochromic anemia, abnormalities of connective tissues, and central nervous system disorders. Sudden death associated with spontaneous rupture of a major blood vessel or the heart itself has been observed in some animal species.

The manifestations of copper deficiency are related to a decrease in several of the copper-containing metalloenzymes. The most severe biochemical alteration is decreased cytochrome oxidase activity; this is manifested as poor growth, anemia, and central nervous system effects. The decreased oxidative metabolism associated with decreased cytochrome oxidase results in poor growth in infants, weight loss, and emaciation. The hypochromic anemia observed during copper deficiency is not distinguishable from iron deficiency anemia; however, it is not responsive to iron administration. A decrease in protoheme synthesis, a result of decreased cytochrome oxidase, has also been observed. As with anemia, the central nervous system effects, primarily the result of hypomyelination, are associated with low activity levels of cytochrome oxidase; the decreased synthesis of phospholipids observed in copper deficiency may also contribute to the development of central nervous system effects. In addition to the decrease in cytochrome oxidase, a decrease in lysyl oxidase is also observed. Lysyl oxidase is involved in the formation of cross-links in collagen and elastin. Depending on the species, this impairment results in bone disorders, a defective cardiovascular system, or abnormal lung structure.

Exposure to Excess Levels of Copper. No copper-specific biomarkers of effects have yet been identified. The most notable sign of toxicity in humans ingesting a beverage or water containing copper is gastrointestinal distress. Symptoms (typically nausea, vomiting, and abdominal pain) usually occur shortly after ingesting the contaminated beverage. The liver is another sensitive target of copper toxicity. Alterations in a number of serum enzymes have been observed in humans and animals with copper-

3. HEALTH EFFECTS

induced liver damage (Chuttani et al. 1965; Epstein et al. 1982; Haywood 1980; Haywood and Comerford 1980; Müller et al. 1998; NTP 1993; Sugawara et al. 1995). The affected serum enzymes include serum aspartate aminotransferase, alanine aminotransferase, and alkaline phosphatase. Increases in serum bilirubin levels have also been observed in humans. Animal studies demonstrate that the rise in serum enzyme activities are the first evidence of liver damage. However, alterations in serum enzyme levels are not unique to copper-induced liver damage.

3.9 INTERACTIONS WITH OTHER CHEMICALS

Numerous studies have demonstrated the interaction between copper and several other metals. Dietary zinc strongly affects copper absorption. A diet high in zinc can result in copper deficiency. Reductions in erythrocyte superoxide dismutase, indicative of marginal copper deficiency, have been found in studies of women ingesting zinc supplements (50 mg zinc/day) for 10 weeks (Yadrick et al. 1989) and men ingesting 50 mg zinc/day for 6 weeks (Fisher et al. 1984). The exact mechanism of the zinc-copper interaction is not known. However, increased dietary zinc results in induction of metallothionein synthesis in the intestine and metallothionein has a greater binding capacity for copper than for zinc. Thus, the dietary copper is sequestered in the intestinal mucosal cell and eventually is excreted in the feces when the mucosal cell is sloughed off (Hall et al. 1979; Whanger and Weswig 1971). Because exposure to excess dietary zinc results in decreased copper absorption, it is often used as a treatment for Wilson's disease (Brewer et al. 1993). An oral/intraperitoneal study in mice provides some evidence that zinc and copper may interact at sites other than the intestine. In this study on the influence of zinc on mitigating the immunotoxicity of copper, mice were exposed to copper sulfate in the drinking water for 8 weeks and received an intraperitoneal injection of zinc sulfate once a week (Pocino et al. 1990). Decreases in the magnitude of the proliferative response to con A or LPS and the antibody response to sheep red blood cells were observed in the copper-exposed mice, but not the mice receiving copper and zinc. However, zinc did not modify the increased production of auto-antibodies reactive with bromelain-treated mouse red blood cells.

Several other divalent cations compete with copper for intestinal absorption. Exposure to dietary cadmium (Evans et al. 1970a), ferrous iron (Wapnir et al. 1993; Yu et al. 1994), and stannous tin (Pekelharing et al. 1994; Wapnir et al. 1993) can result in decreased copper absorption. In the case of cadmium, the decrease is related cadmium induction of metallothionein and the binding of copper to it. Tetrathiomolybdate is used for the treatment of Wilson's disease (Brewer 1995) and excessive dietary molybdenum can also result in decreased uptakes and, therefore, copper utilization and toxicity. Two

3. HEALTH EFFECTS

mechanisms of action of tetrathiomolybdate have been proposed: it reacts with copper-metallothionein to form a soluble complex which is excreted (Ogra et al. 1996) and it can complex with nonceruloplasmin plasma copper, preventing its cellular absorption (Brewer 1995).

Because selenide is a strong reducing agent (Frost 1972), it has been postulated that selenium may play a role in detoxifying copper. Aburto et al. (2001a, 2001b) examined the possible interaction between copper and selenium. Selenium did not influence the hepatotoxicity of copper in rats fed diets with excess levels of copper and inadequate, adequate, or excess levels of dietary selenium. Hepatic copper levels and histological alterations were not significantly different in rats receiving a high copper/high selenium diet as compared to rats receiving a high copper/adequate selenium diet.

3.10 POPULATIONS THAT ARE UNUSUALLY SUSCEPTIBLE

A susceptible population will exhibit a different or enhanced response to copper than will most persons exposed to the same level of copper in the environment. Reasons may include genetic makeup, age, health and nutritional status, and exposure to other toxic substances (e.g., cigarette smoke). Whatever the basis of the increased susceptibility may be, the result is reduced detoxification or excretion of copper or compromised functioning of organs affected by copper. Populations at greater risk due to unusually high exposure to copper are discussed in Section 6.7, Populations with Potentially High Exposures.

A number of populations of individuals unusually susceptible to copper toxicity have been identified. The increased susceptibility to copper toxicity is associated with genetic defects that impair copper homeostatic mechanisms. Wilson's disease, also referred to as hepatolenticular degeneration, is an autosomal recessive disorder with a worldwide incidence of 1 in 30,000 (Scheinberg and Sternlieb 1996). The primary genetic defect in Wilson's disease is in ATP7B, which encodes a P-type ATPase (Wilson protein), which delivers copper to ceruloplasmin. The genetic defect results in impaired biliary excretion of copper and an accumulation of copper in the liver. As described by Brewer and Yuzbasiyan-Gurkan (1992), the progression of the disease begins with an accumulation of copper in the liver, damage to the liver, and subclinical liver cirrhosis. Over time, the individual will develop hepatic, neurological, and psychiatric symptoms. The hepatic effects are characterized by jaundice, hypoalbuminemia, ascites, coagulation defects, hyperammonemia, hepatic encephalopathy, and/or liver failure; in the cases of massive liver failure, large amounts of copper are released from the liver resulting in hemolytic anemia. Neurological symptoms include tremors and other movement disorders and speech abnormalities. Psychiatric and behavioral symptoms are often found in individuals also manifesting neurological other

3. HEALTH EFFECTS

symptoms. The psychiatric symptoms include reduced performance in school or work, inability to cope, depression, very labile moods ranging from mania to depression, sexual exhibitionism, and frank psychosis. Individuals with Wilson's disease have low serum ceruloplasmin levels, elevated urinary copper levels, and elevated liver copper levels; Kayser-Fleischer rings, which result from corneal copper deposits, are also detected in individuals with Wilson's disease. Individuals who are heterozygotes for Wilson's disease may also be unusually susceptible to the toxicity of copper. Increases in urinary copper and hepatic concentrations and decreased copper incorporation into ceruloplasmin have been observed in heterozygotes. These findings may suggest that long-term exposure to elevated levels of copper may result in copper overload. Although the incidence of heterozygotes is not known, NAS (2000) estimates that 1 in 40,000 individuals (approximately 1% of the U.S. population) may be heterozygotes for Wilson's disease.

Indian childhood cirrhosis (ICC) and idiopathic copper toxicosis (ICT) are two syndromes that result in severe, often fatal, liver cirrhosis in infants and young children. Although the basis of the defect has not been firmly established, it is believed to be due to an inherited autosomal recessive defect in copper metabolism aggravated by high copper intake (Bhave et al. 1982, 1987; Müller et al. 1996, 1998). ICC occurs in infants and children living in rural areas of the Indian subcontinent who are introduced early to cow or buffalo milk that is stored or heated in brass or copper vessels. Copper is believed to be the causative agent because the milk has very high copper levels, very high copper levels are found in the liver, and replacing the brass or copper vessels with aluminum or stainless steel vessels eliminates the occurrence of ICC in siblings of ICC affected children (Bhave et al. 1982; Tanner 1998). A high degree of parental consanguinity, the occurrence of ICC in children, but not the parents, and the fact that 22% of siblings affected suggest an autosomal recessive component to the disease (Pandit and Bhave 1996; Tanner 1998). For ICT, which includes Tyrolean infantile cirrhosis, sources of high copper exposure have been identified. For the 138 cases of ICT in children living in the Tyrolean region of Austria, the source of the copper was the use of a water/unpasteurized cow's milk mixture that was heated in a copper pot (Müller et al. 1996). For the other cases of ICT that have been identified in a number of countries, the source of the excess copper intake was drinking water (Müller et al. 1998). The similarity of ICT to ICC has prompted investigators to suggest that ICT may also be due to an autosomal recessive genetic defect in copper metabolism and excessive copper intake at a very young age. A genealogical investigation by Müller et al. (1996) provides supportive evidence for a genetic basis of the disease.

It has been postulated that individuals with a deficiency of the enzyme glucose-6-phosphate dehydrogenase would be susceptible to the toxic effects of oxidative stressors such as copper (Calabrese

3. HEALTH EFFECTS

and Moore 1979; Chugh and Sakhuja 1979). This has not been supported by epidemiological or experimental data. In the blood, most of the copper is bound to ceruloplasmin. With the exception of ingestion of a very large dose of copper salts, the levels of nonceruloplasmin bound copper remain low following copper exposure. Thus, it is unlikely that this relatively small change in free copper would alter the survival of glucose-6-phosphate dehydrogenase deficient red cells.

3.11 METHODS FOR REDUCING TOXIC EFFECTS

This section will describe clinical practice and research concerning methods for reducing toxic effects of exposure to copper. However, because some of the treatments discussed may be experimental and unproven, this section should not be used as a guide for treatment of exposures to copper. When specific exposures have occurred, poison control centers and medical toxicologists should be consulted for medical advice. The following texts provide specific information about treatment following exposures to copper:

Ellenhorn MJ, Schonwald S, Ordog G, et al., eds. 1997. *Medical toxicology: Diagnosis and treatment of human poisoning*. Second edition. Baltimore, MD: Williams & Wilkins, 1554-1556.

Goldfrank LR, Flomenbaum FE, Lewin NA, et al., eds. 1998. *Goldfrank's toxicologic emergencies*. Sixth edition. Stamford, CT: Appleton & Lange, 1339-1340.

Haddad LM, Shannon MW, Winchester JF, eds. 1998. *Clinical management of poisoning and drug overdose*. Third edition. Philadelphia, PA: WB Saunders, 165.

3.11.1 Reducing Peak Absorption Following Exposure

Following ingestion of copper or copper compounds, milk or water should be given immediately after ingestion and/or prior to vomiting. Because of the strong emetic properties of copper and copper compounds, vomiting usually occurs shortly after ingestion. Induction of vomiting and gastric lavage are contraindicated following ingestion of caustic copper salts, such as copper sulfate. Gastric lavage may be indicated after ingestion of noncorrosive copper compounds (HSDB 2002).

For individuals with Wilson's disease, the administration of a diet high in zinc is used as a maintenance treatment (Brewer et al. 1989). The zinc interferes with copper absorption by inducing intestinal metallothionein resulting in increased copper sequestration (Brewer et al. 1992).

3. HEALTH EFFECTS

3.11.2 Reducing Body Burden

A number of methods have been employed to reduce copper body burden. These methods range from the use of chelating agents to increases in dietary levels of zinc and molybdenum. Gao et al. (1989) tested the *in vitro* effectiveness of five chelating agents using human serum albumin. The agents (in order of decreasing effectiveness) were ethylenediaminetetraacetate (EDTA), diethylene triaminopentaacetate, ethylene glycol-*bis*-(aminoethylether)-tetraacetate, nitrilotriacetate, and iminodiacetate. The *in vivo* effectiveness of these agents has not been established. D-penicillamine is often used to decrease the elevated levels of hepatic copper in individuals with Wilson's disease (Walshe 1996; Walshe and Yealland 1993) and idiopathic childhood cirrhosis (Rodeck et al. 1999). However, a number of potential side effects have been associated with penicillamine treatment (Brewer and Yuzbasiyan-Gurkan 1992). A variety of other chelating agents have been tested in copper loaded rats. Tetraethylenepentamine pentahydrochloride (TETREN) was more effective in increasing urinary excretion of copper than 1,4,7,11-tetraazaundecane tetrahydrochloride (TAUD) or penicillamine, which were equally effective (Domingo et al. 2000). TETREN did not result in a decrease in copper levels in the liver, although a significant decrease in kidney copper levels was observed. In contrast, TAUD and penicillamine reduced the levels of copper in the liver. None of the three chelating agents affected the amount of copper excreted into the feces.

The known interaction between copper and molybdenum have been used to treat individuals with Wilson's disease. The administration of tetrathiomolybdate to individuals with neurological or psychiatric symptoms associated with Wilson's disease has resulted in an improvement or reversal of symptoms (Brewer 1995). In blood plasma, tetrathiomolybdate complexes with nonceruloplasmin plasma copper, preventing its cellular absorption. Studies in Long-Evans Cinnamon rats, a model for Wilson's disease, and sheep have found that administration of tetrathiomolybdate results in a dramatic decrease in the levels of copper in the liver (Humphries et al. 1988; Kumaratilake and Howell 1989; Ogra et al. 1996) and decreased liver damage (Humphries et al. 1988). Tetrathiomolybdate also reacts with copper bound to metallothionein resulting in a soluble copper-tetrathiomolybdate complex (Ogra et al. 1996). The addition of molybdenum to a high sulfur, low copper diet can result in a decrease in liver and plasma copper levels in copper loaded sheep (van Ryssen 1994).

Although zinc is used in the treatment of Wilson's disease to decrease the absorption of copper, zinc does not appear to be effective in reducing the copper body burden. No alterations in hepatic copper levels were observed in sheep administered a low copper, high zinc diet (van Ryssen 1994). A reduction in hepatic copper levels has been observed in dogs administered a high zinc diet (Brewer et al. 1992);

3. HEALTH EFFECTS

however, it is believed that the reduction was secondary to the induction of copper deficiency and the mobilization of copper from the liver (van Ryssen 1994).

3.11.3 Interfering with the Mechanism of Action for Toxic Effects

There are limited data on methods for interfering with the mechanisms of action of copper. An *in vitro* study suggested that lazaroids (21-aminosteroids) may have a protective effect against copper-induced erythrocyte lipid peroxidation (Fernandes et al. 1992). Oxidative damage to the erythrocyte membrane may be the cause of the hemolysis observed following exposure to very high doses of copper.

3.12 ADEQUACY OF THE DATABASE

Section 104(i)(5) of CERCLA, as amended, directs the Administrator of ATSDR (in consultation with the Administrator of EPA and agencies and programs of the Public Health Service) to assess whether adequate information on the health effects of copper is available. Where adequate information is not available, ATSDR, in conjunction with the National Toxicology Program (NTP), is required to assure the initiation of a program of research designed to determine the health effects (and techniques for developing methods to determine such health effects) of copper.

The following categories of possible data needs have been identified by a joint team of scientists from ATSDR, NTP, and EPA. They are defined as substance-specific informational needs that if met would reduce the uncertainties of human health assessment. This definition should not be interpreted to mean that all data needs discussed in this section must be filled. In the future, the identified data needs will be evaluated and prioritized, and a substance-specific research agenda will be proposed.

3.12.1 Existing Information on Health Effects of Copper

The existing data on health effects of inhalation, oral, and dermal exposure of humans and animals to copper are summarized in Figure 3-4. The purpose of this figure is to illustrate the existing information concerning the health effects of copper. Each dot in the figure indicates that one or more studies provide information associated with that particular effect. The dot does not necessarily imply anything about the quality of the study or studies, nor should missing information in this figure be interpreted as a “data

3. HEALTH EFFECTS

Figure 3-4. Existing Information on Health Effects of Copper

	Systemic									
	Death	Acute	Intermediate	Chronic	Immunologic/Lymphoretic	Neurologic	Reproductive	Developmental	Genotoxic	Cancer
Inhalation				●						●
Oral	●	●	●	●						
Dermal		●		●	●					

Human

	Systemic									
	Death	Acute	Intermediate	Chronic	Immunologic/Lymphoretic	Neurologic	Reproductive	Developmental	Genotoxic	Cancer
Inhalation		●	●		●					
Oral	●	●	●	●	●		●	●	●	●
Dermal										

Animal

● Existing Studies

3. HEALTH EFFECTS

need.” A data need, as defined in ATSDR’s Decision Guide for Identifying Substance-Specific Data Needs Related to Toxicological Profiles (Agency for Toxic Substances and Disease Registry 1989), is substance-specific information necessary to conduct comprehensive public health assessments.

Generally, ATSDR defines a data gap more broadly as any substance-specific information missing from the scientific literature.

The toxicity of inhaled copper has been investigated in a couple of occupational exposure studies. These studies examined a limited number of systemic end points, and exposure is poorly characterized. There are numerous reports and studies on the toxicity of ingested copper in humans. Most of the reports and studies focused on the gastrointestinal effects following acute exposure to copper in drinking water or other beverages. Data on other health effects in humans comes from individuals with Wilson’s disease, Indian childhood cirrhosis, and idiopathic copper toxicosis. These diseases/syndromes are the result of genetic defect(s) resulting in impaired copper kinetics; the latter two syndromes are also associated with exposure to high levels of copper in drinking water or milk (due to storage of milk in brass vessels). These studies provide information on potential targets of toxicity, primarily the liver.

Information on the dermal toxicity of copper is limited to reports of contact dermatitis in individuals and eye irritation in workers exposed to copper dust.

As with the human database, there are limited data on the toxicity of inhaled copper in animals. The available studies have primarily focused on potential respiratory effects. There is a more extensive database on the toxicity of ingested copper in animals. These studies have found a number of systemic effects, including gastrointestinal, hepatic, and renal effects following acute, intermediate, and chronic exposure. Immunological and developmental effects have also been reported in animal studies. Several studies have also examined potential neurological and reproductive targets, but have not found effects. Carcinogenic effects were not found in several animal studies; however, the studies are limited in scope and tested low doses. No animal studies examining the dermal toxicity of copper were identified.

3.12.2 Identification of Data Needs

Acute-Duration Exposure. No data were located regarding health effects after acute inhalation exposure to copper in humans. Animal data are limited to information from studies in mice and hamsters conducted by Drummond et al. (1986). Respiratory tract irritation and impaired immune function were

3. HEALTH EFFECTS

observed. This study was not selected as the basis for an acute-duration inhalation MRL because it only examined a limited number of end points, and the liver and kidney, which are targets following oral exposure, were not examined; in addition, the animals were only exposed for 3 hours/day. Additional inhalation studies are needed to identify the critical targets of toxicity and to establish concentration-response relationships for copper. The most commonly reported effect in humans acutely exposed to copper is gastrointestinal upset. The reported symptoms include nausea, vomiting, abdominal pain, and diarrhea (Chutanni et al. 1965; Gill and Bhagat 1999; Gotteland et al. 2001; Nicholas and Brist 1968; Olivares et al. 2001; Pizarro et al. 1999, 2001; Semple et al. 1960; Walsh et al. 1977). Hepatic and renal effects have also been seen in individuals ingesting lethal doses of copper sulfate (Chuttani et al. 1965). Animal studies support the identification of the gastrointestinal tract, liver, and kidneys as sensitive targets of copper toxicity. Hyperplasia of the forestomach has been observed in rats and mice exposed to copper sulfate in the diet for 14 days (NTP 1993). Hepatic effects ranging from increases in alanine aminotransferase activity to hepatocellular necrosis and renal effects (protein droplets in proximal tubules) have been observed in rats exposed to fairly high doses of copper sulfate in the diet (Haywood 1980; Haywood and Comerford 1980; Haywood et al. 1985b; NTP 1993). Decreases in body weight gain have also been observed in rats (NTP 1993). The acute-duration oral database was considered adequate for derivation of an MRL. The MRL was based on gastrointestinal upset in women ingesting drinking water containing copper sulfate for 2 weeks (Pizarro et al. 1999). There are limited data on the dermal toxicity of copper. Pruritic dermatitis and allergic contact dermatitis have been reported in humans exposed to copper. No animal studies were identified. These data provide suggestive evidence that copper may be irritative to the skin; additional dermal studies are needed to determine whether copper exposure will also result in systemic effects.

Intermediate-Duration Exposure. No studies were located regarding health effects in humans after intermediate-duration inhalation. Only one animal inhalation exposure study was located. This study did not find any adverse histological alterations in the lungs or functional alterations in alveolar macrophages of rabbits exposed to copper chloride (Johansson et al. 1983, 1984). Because the lungs were the only tissues examined, these studies were not considered suitable for derivation of an intermediate-duration inhalation MRL for copper. Additional studies are needed to identify the critical targets of toxicity and establish concentration-response relationships for inhaled copper. Three experimental human studies and two community-based studies have examined the oral toxicity of copper in healthy humans. The primary focus of these studies was examination of the potential of low doses of copper to induce hepatic effects in adults (Araya et al. 2003b; Pratt et al. 1985) or infants (Olivares et al. 1998; Zietz et al. 2003a, 2003b); no adverse effects were found. The Araya et al. (2003b) study also assessed the potential for gastrointestinal

3. HEALTH EFFECTS

effects in adults and found significant increases in the incidence of effects as a function of dose/duration. A number of animal studies have reported adverse liver and kidney effects following intermediate-duration oral exposure to copper compounds (Epstein et al. 1982; Fuentealba et al. 2000; Haywood 1980, 1985; Haywood and Comerford 1980; Haywood and Loughran 1985; Haywood et al. 1985a, 1985b; Kumar and Sharma 1987; NTP 1993). The observed liver and kidney effects demonstrated dose- and duration-response relationships. The studies by Haywood and associates demonstrate that rats can develop a tolerance to copper following repeated oral exposure. Studies in other animal species are needed to determine if this phenomenon is unique to rats or is observed in other species as well. Other systemic effects that have been reported in animals include hyperplasia of the forestomach mucosa (NTP 1993), decreased erythrocyte and hemoglobin levels (Kumar and Sharma 1987; Rana and Kumar 1980; Suttle and Mills 1966a), and decreased body weight gain or weight loss (Haywood 1985; Haywood and Loughran 1985; Kline et al. 1971; Llewellyn 1985; NTP 1993). For the most part, these studies involved dietary exposure of rats to copper sulfate; additional studies in other species would be useful for identifying a model for human toxicity. The Araya et al. (2003b) human study was used as the basis of an intermediate-duration oral MRL for copper. No data on the dermal toxicity of copper following intermediate-duration exposure were identified. Studies are needed to identify the critical targets of copper toxicity following dermal exposure.

Chronic-Duration Exposure and Cancer. Systemic effects such as nausea (Suciu et al. 1981), hepatomegaly (Suciu et al. 1981), decreased hemoglobin and erythrocyte levels (Finelli et al. 1981), and respiratory irritation (Askergren and Mellgren 1975; Suciu et al. 1981) have been observed in workers exposed to copper dust. The mild gastrointestinal effects observed in some workers were attributed to swallowing airborne copper dust (Suciu et al. 1981). The poor characterization and/or the lack of controls preclude deriving a chronic-duration inhalation MRL based on the occupational exposure studies. Additional studies are needed to identify the critical targets of toxicity of inhaled copper. There are numerous reports of severe health effects in infants and children ingesting copper-contaminated milk or water containing high levels of copper (Müller et al. 1996, 1998; Pandit and Bhave 1996; Tanner 1998). Indian childhood cirrhosis and idiopathic copper toxicosis are characterized by severe liver cirrhosis occurring before the age of 5 years. There is suggestive evidence that both of these syndromes are related to increased dietary intake of copper in conjunction with increased genetic susceptibility. Nausea, vomiting, and abdominal pain were reported by members of a family with very high levels of copper in the drinking water (Spitalny et al. 1984). The animal database on the oral toxicity of copper following chronic-duration exposure is limited to one study (Massie and Aiello 1984) that found a decrease in lifespan and no effect on body weight gain in mice exposed to copper gluconate for 850 days. No other

3. HEALTH EFFECTS

end points of toxicity were examined in this study. The database was considered inadequate for derivation of a chronic-duration oral MRL. Additionally, studies that examine a variety of end points are needed to identify the critical targets of toxicity and establish dose-response relationships. Information on the dermal toxicity of copper is limited to a report of ocular irritation in workers exposed to copper dust (Askergren and Mellgren 1975). Additional dermal toxicity studies are needed to identify the critical targets of toxicity following dermal exposure.

Data on the carcinogenicity of copper in humans are limited to a study of copper miners (Chen et al. 1993) and a follow-up to this study (Chen et al. 1995). Increased risk of cancer, stomach cancer, and lung cancer were observed. Because the workers were also exposed to radon and radon daughters, silica, iron, titanium, sulfur, and arsenic, a causal relationship between copper and increased cancer risk can not be established. No studies examining the association between copper ingestion and cancer risk in humans were identified. Several animal studies have examined the carcinogenic potential of ingested copper (BRL 1968; Greene et al. 1987; Kamamoto et al. 1973). These studies are limited in scope, the studies by Green et al. (1987) and Kamamoto et al. (1973) only examined one potential target, and tested fairly low doses of copper. No dermal carcinogenicity studies in humans or animals were identified. Additional studies by the inhalation, oral, and dermal routes are needed to assess the carcinogenic potential of copper in humans.

Genotoxicity. No data on the genotoxicity of copper in humans were located; studies of workers or individuals accidentally exposed to high levels of copper would provide value information on its genotoxic potential in humans. The available genotoxicity data suggest that copper is a clastogenic agent (Agarwal et al. 1990; Bhunya and Jena 1996; Bhunya and Pati 1987; Sideris et al. 1988). However, mixed results have been found in point mutation assays (Demerec et al. 1951; Marzin and Phi 1985; Singh 1983; Tso and Fung 1981; Wong 1988). Additional studies are needed to assess copper's potential to induce point mutations. Several studies have also shown that exposure to copper can result in DNA damage (Garrett and Lewtas 1983; Sideris et al. 1988; Sina et al. 1983).

Reproductive Toxicity. There are no human studies and two animal studies that examined the potential of copper to induce reproductive effects. These studies did not find any adverse alterations in reproductive performance in mink (Aulerich et al. 1982), sperm morphology in rats and mice (NTP 1993), or vaginal cytology in rats or mice (NTP 1993). The NTP (1993) study also did not find histological alterations in reproductive tissues. Multigeneration or continuous breeding studies would provide information on the reproductive effects of copper in animals, which may be used to assess possible reproductive effects in humans exposed to high levels of copper.

3. HEALTH EFFECTS

Developmental Toxicity. Developmental studies by the oral route in rats (Haddad et al. 1991) and mice (Lecyk 1980) have shown that high copper intakes can result in impaired growth. The developmental toxicity of copper in humans has not been adequately investigated. No data were located regarding developmental effects of copper after inhalation or dermal exposures in humans or animals. Further studies in other animal species would provide valuable information on the potential of copper to adversely affect development. Such information might be relevant to humans.

Immunotoxicity. There are limited data on the immunotoxic potential of copper and its compounds. Reports on humans developing dermatitis after dermal exposure to copper (Barranco 1972; Saltzer and Wilson 1968) suggest that copper is an allergen. This is supported by a report of a woman developing dermatitis after insertion of a copper IUD (Barranco 1972). Immunological effects also have been observed in mice (Drummond et al. 1986) following acute inhalation exposure to copper sulfate. In addition, impaired immune function has been observed in mice exposed to copper chloride (Pocino et al. 1991) or copper sulfate (Pocino et al. 1990) in drinking water. Intermediate-duration studies concentrating on immunologic effects in different species would be useful for establishing dose-response relationships and assessing whether there are species differences. More studies in humans and animals that examine the immune response to copper exposure and the mechanisms involved therein would be useful.

Neurotoxicity. Neurological impairment has been observed in factory workers exposed to copper dust. No effects on neurobehavioral performance were observed in rats exposed to copper in the diet (Murthy et al. 1981). However, this study did find alterations in the levels of a dopamine metabolite, suggesting that copper may adversely affect the nervous system. Additional studies are needed to further investigate the neurotoxic potential of copper; these studies should assess the potential of copper to perturb dopaminergic pathways and related functions.

Epidemiological and Human Dosimetry Studies. Several studies have examined the toxicity of inhaled copper in workers (Askergren and Mellgren 1975; Finelli et al. 1981; Suciú et al. 1981). These studies have primarily focused on the respiratory tract, although health examinations revealed other adverse effects (e.g., hepatomegaly). Chen et al. (1993, 1995) examined the carcinogenic potential of inhaled copper. In general, these studies are limited by poor exposure characterization, co-exposure to several toxic and/or carcinogenic compounds (e.g., arsenic, cadmium, radon, lead), and limited number of end points examined. Occupational exposure studies examining populations of workers exposed to

3. HEALTH EFFECTS

copper and with minimal exposure to other metals would be useful in assessing the toxicity of inhaled copper. These studies should examine a wide variety of end points, particularly the gastrointestinal tract, liver, and kidneys, which are targets of toxicity following oral exposure.

There are numerous reports of accidental or intentional ingestion of copper. The most commonly reported effect in these studies is gastrointestinal upset. There have also been several experimental studies designed to identify a no effect level for gastrointestinal upset following short-term (2 weeks or less) exposure to copper in drinking water (Olivares et al. 2001; Pizarro et al. 1999, 2001). There are several subpopulations of individuals exposed to higher than normal levels of copper; these groups include communities with higher than normal levels of copper in drinking water and individuals ingesting higher than normal levels of copper in the form of supplements. Studies of these groups that involved examination for a variety of potential effects (including gastrointestinal, hepatic, and renal effects, which have been shown to be sensitive end points in animal studies) could provide useful information on the toxicity of copper in otherwise healthy humans. In addition, if the study group included both children and adults, these data could address the issue of age-related differences in toxicity.

Biomarkers of Exposure and Effect.

Exposure. Copper levels can be measured in tissues, body fluids, and excreta. Whole blood, serum, and urine copper levels have been established in healthy individuals. It has been demonstrated that copper levels in the body increase with increased exposure after acute poisoning. Similarly, increased copper levels were observed in workers after occupational exposure. Serum and urine copper levels, plasma ceruloplasmin levels, and clinical manifestations are specific indicators of copper status. It is doubtful that a single “specific” biomarker of intoxication resulting from exposure to a specific metal will be found. In any case, elevated tissue copper levels should be a sufficient indicator of exposure and the possibility of intoxication.

Effect. There are no specific biomarkers for copper toxicity. Individuals with Wilson's disease are usually diagnosed by examining serum and urine copper levels, plasma ceruloplasmin levels, and clinical manifestations. However, the relationship between serum and urine levels of copper and health effects are not known. Studies examining the possible correlation between blood levels or excreta levels of copper with effects would facilitate medical surveillance leading to early detection and possible treatment.

Absorption, Distribution, Metabolism, and Excretion. The absorption, distribution, metabolism, and excretion of copper administered orally have been studied in animals and, to some

3. HEALTH EFFECTS

extent, in humans. Furthermore, alterations in copper absorption, distribution, and excretion have been studied in deficiency and toxicity states. Despite the information on copper absorption, there is very little information on differences between absorption rates of the various Cu(II) compounds and differences between the bioavailability of copper from food and water.

Several studies have shown that ingested or implanted metallic copper results in increased serum copper levels and liver toxicity (Keller and Kaminski 1984; Yelin et al. 1987). Studies on the release of copper ions from both ingested and implanted metallic copper would be useful.

There is very limited information on copper absorption following inhalation exposure, and data on the absorption of copper through the skin are limited. Further studies in animals on the rate and extent of copper absorption following exposure from both the inhalation route and the dermal route would more fully characterize copper toxicokinetics in animals and by extrapolation in humans.

There is evidence that animals develop a tolerance to continued high doses of copper; more information on the mechanism(s) involved might be useful to establish if humans also could develop tolerance, as well as to provide insight for the development of more effective and efficient treatment of copper toxicity.

Comparative Toxicokinetics. The metabolism of copper has been studied in rats, pigs, hamsters, and humans. However, there are no comparative studies on the effects of high copper intakes on the distribution of copper in the body or the development of tolerance to continued high intakes of copper. Furthermore, the animal species that might serve as the best model for extrapolating results to humans is not known.

Methods for Reducing Toxic Effects. Methods for reducing the toxic effects of copper have primarily focused on reducing body burdens. Many of these methods have been designed for individuals with Wilson's disease; however, it is likely that these would also be effective in other instances of copper intoxication. D-penicillamine (Rodeck et al. 1999; Walshe 1996; Walshe and Yealland 1993) is the most commonly used palliative agent for Wilson's disease; however, it has a number of potentially deleterious side effects. Studies in animals suggest that TETREN and TAUD may also be effective chelating agents (Domingo et al. 2000). Other treatment methods include administration of tetrathiomolybdate (Humphries et al. 1988; Kumaratilake and Howell 1989; Ogra et al. 1996), diets high in molybdenum and sulfur (van Ryssen 1994), and diets high in zinc (Brewer et al. 1992; van Ryssen 1994). Further studies

3. HEALTH EFFECTS

are needed to identify treatments that would interfere with copper's mechanism of toxicity and reduce body burden with minimal side effects.

Children's Susceptibility. There are some data on the toxicity of copper in infants and children. Severe liver damage has been reported in infants and children. These effects are typically clustered in geographically regions and have been grouped into two syndromes: Indian childhood cirrhosis and idiopathic copper toxicosis. Both of these syndromes are associated with elevated copper intakes and early dietary introduction of milk and/or formula, and are believed to have a genetic component. Very high levels of copper are found in the livers of affected children, suggesting that the mechanism of action is related to impaired copper efflux. Additional studies are needed to determine the mechanism of toxicity and to ascertain copper's role in the observed effects. Information that would provide a better understanding of copper absorption and excretion in early infancy and homeostatic mechanisms in infants would also provide valuable documentation on these syndromes and their relationship to copper.

Child health data needs relating to exposure are discussed in Section 6.8.1 Identification of Data Needs: Exposures of Children.

3.12.3 Ongoing Studies

Ongoing studies pertaining to copper have been identified and are shown in Table 3-5 (FEDRIP 2003).

3. HEALTH EFFECTS

Table 3-5. Ongoing Studies on Copper

Investigator	Affiliation	Research description	Sponsor
Turnland JR	Agricultural Research Service, Davis, California	Influence of high copper intake on copper homeostasis and mineral metabolism	USDA
Kelvey LM	Agricultural Research Service, Grand Forks, North Dakota	Determination of a no effect level for copper	USDA
Reeves PG	Agricultural Research Service, Grand Forks, North Dakota	Correlation between sperm motility and copper status in humans and animals	USDA
Harris ED	Texas A & M University	Copper metabolism and homeostasis in humans and animals	CSREES TEX
Thiele DJ	University of Michigan at Ann Arbor	Copper homeostasis	NIGMS
Culotta VC	John Hopkins University	Intracellular pathways of copper trafficking	NIEHS
Gitlin JD	Washington University	Copper chaperones	NIDDKD

CSREES TEX = Cooperative State Research Education and Extension Service, Texas; NIDDKD = National Institute of Diabetes and Digestive and Kidney Disease; NIEHS = National Institute of Environmental Health and Science; NIGMS = National Institute of General Medical Sciences; USDA = U.S. Department of Agriculture

4. CHEMICAL AND PHYSICAL INFORMATION

4.1 CHEMICAL IDENTITY

Copper is the first element of Group IB of the periodic table and displays four oxidation states: Cu(0), Cu(I), Cu(II), and Cu(III). Along with silver and gold, it is classified as a noble metal and, like them, can be found in nature in the elemental form. Copper's unique chemical and physical properties have made it one of the most important metals. These properties include high thermal conductivity, high electrical conductivity, malleability, low corrosion, alloying ability, and pleasing appearance. Properties of metallic copper such as electrical conductivity and fabricability vary markedly with purity. Standard classifications have been defined according to processing method. For example, ASTM B5-74 is >99.90% pure and is the accepted basic standard for electrolyte copper wire bars, etc. (Tuddenham and Dougall 1978). Data on the chemical identity of copper are shown in Table 4-1. Data on the chemical identity of copper sulfate, the most important commercial compound of copper, are shown in Table 4-2.

4.2 PHYSICAL AND CHEMICAL PROPERTIES

Copper is positioned below hydrogen in the electromotive-force series, so it will not displace hydrogen ions from dilute acid. Accordingly, copper will not dissolve in acid unless an oxidizing agent is present. Therefore, while it readily dissolves in nitric and hot concentrated sulfuric acid, it only dissolves slowly in hydrochloric and dilute sulfuric acid, and then only when exposed to the atmosphere (Hawley 1981). It is also attacked by acetic acid and other organic acids. When exposed to moist air, a characteristic green layer of the basic copper carbonate slowly forms (Windholz 1983). This tightly adherent coating protects the underlying metal from further attack and is also prized for its appearance. Copper dissolves in ammonia in the presence of air, forming the cupric ammonium complex ion $\text{Cu}(\text{NH}_3)_4^{2+}$ (Cotton and Wilkinson 1980).

Cu(I) or the cuprous ion disproportionates rapidly (<1 second) in aqueous solution to form Cu(II) and Cu(0) (Cotton and Wilkinson 1980). The only Cu(I) compounds that are stable in water are extremely insoluble ones such as CuCl. It has been shown that Cu(I) complexes may be formed in seawater by photochemical processes and may persist for several hours (Moffett and Zika 1987). Cuprous compounds are generally colorless.

4. CHEMICAL AND PHYSICAL INFORMATION

Table 4-1. Chemical Identity of Copper

Characteristic	Information	Reference
Chemical name	Copper	
Synonym(s)	Not reported	
Registered trade names(s)	Not reported	
Chemical formula	Cu	HSDB 2004
Chemical structure	Face-centered cubic	Budavari 2001
Identification numbers:		
CAS registry	7440-50-8	HSDB 2004
NIOSH RTECS	GL5324000	HSDB 2004
EPA hazardous waste	Not reported	
OHM/TADS	Not reported	
DOT/UN/NA/IMCO shipping	Not reported	
HSDB	1622	HSDB 2004
NCI	Not reported	

CAS = Chemical Abstracts Services; DOT/UN/NA/IMCO = Department of Transportation/United Nations/North America/International Maritime Dangerous Goods Code; EPA = Environmental Protection Agency; HSDB = Hazardous Substances Data Bank; NCI = National Cancer Institute; NIOSH = National Institute for Occupational Safety and Health; OHM/TADS = Oil and Hazardous Materials/Technical Assistance Data System; RTECS = Registry of Toxic Effects of Chemical Substances

4. CHEMICAL AND PHYSICAL INFORMATION

Table 4-2. Chemical Identity of Copper Sulfate

Characteristic	Information	Reference
Chemical name	Copper sulfate	
Synonym(s)	Cupric sulfate; blue stone; blue vitriol; cupric sulphate; Roman vitriol; Salzburg vitriol; blue copperas; copper(II) sulfate	Budavari 2001; Hawley 1997; HSDB 2004
Registered trade names(s)	Not reported	
Chemical formula	CuO ₄ S	Budavari 2001
Chemical structure	CuSO ₄	Budavari 2001
Identification numbers:		
CAS registry	7758-98-7	HSDB 2004
NIOSH RTECS	GL8800000	HSDB 2004
EPA hazardous waste	Not reported	
OHM/TADS	Not reported	
DOT/UN/NA/IMCO shipping	Not reported	
HSDB	916	HSDB 2004
NCI	Not reported	

CAS = Chemical Abstracts Services; DOT/UN/NA/IMCO = Department of Transportation/United Nations/North America/International Maritime Dangerous Goods Code; EPA = Environmental Protection Agency; HSDB = Hazardous Substances Data Bank; NCI = National Cancer Institute; NIOSH = National Institute for Occupational Safety and Health; OHM/TADS = Oil and Hazardous Materials/Technical Assistance Data System; RTECS = Registry of Toxic Effects of Chemical Substances

4. CHEMICAL AND PHYSICAL INFORMATION

Cu(II) or the cupric ion is the most important oxidation state of copper. Cu(II) is the oxidation state of copper generally encountered in water (Cotton and Wilkinson 1980). Cupric ions are coordinated with six water molecules in solution; the arrangement of the water molecules is distorted in that there are four molecules bound closely to the copper in a planar array while the other two are more loosely bound in polar position (Cotton and Wilkinson 1980). Addition of ligands such as NH_3 to the solution will successively displace only the four planar water molecules. Most cupric compounds and complexes are blue or green in color. They are frequently soluble in water.

When Cu(II) is introduced into the environment, the cupric ion typically binds to inorganic and organic materials contained within water, soil, and sediments. In water, Cu(II) binds to dissolved organics (e.g., humic or fulvic acids). The Cu(II) ion forms stable complexes with $-\text{NH}_2$, $-\text{SH}$ and, to a lesser extent, $-\text{OH}$ groups of these organic acids. Cu(II) will also bind to inorganic and organic components in sediments and soils with varying affinities. For example, Cu(II) binds strongly to hydrous manganese and iron oxides in clay and to humic acids in organic matter, but much less strongly to aluminosilicates in sand. As in water, the binding affinities of Cu(II) with inorganic and organic matter in sediments and soils is dependent on pH, the oxidation-reduction potential in the local environment, and the presence of competing metal ions and inorganic anions.

Cu(III) is strongly oxidizing and only occurs in a few compounds (Kust 1978). At this time, none of these compounds are industrially important or environmentally significant.

Data on the physical and chemical properties of copper and copper sulfate are shown in Table 4-3.

4. CHEMICAL AND PHYSICAL INFORMATION

Table 4-3. Physical and Chemical Properties of Copper and Copper Sulfate

Property	Copper	Copper sulfate
Molecular weight	63.546 ^a	159.61 ^a
Color	Reddish ^b	Blue crystals, white dehydrated ^b
Physical state	Solid ^b	Solid ^b
Melting point	1,083 ^c	Decomposes at 560 ^a
Boiling point	2,595 ^c	No data
Specific gravity (20/4 °C)	8.94 ^c	3.60 ^a 2.286 (pentahydrate) ^a
Odor	No data	None ^d
Odor threshold		
Air	No data	No data
Water	No data	No data
Taste	No data	No data
Taste threshold	No data	No data
pK _a		
Solubility:		
Water	Insoluble ^e	32.0g/100g (20 °C) ^f
Organic solvent(s)		Soluble in methanol, slightly soluble in ethanol ^b
Partition coefficients:		
Log K _{ow}	No data	No data
Log K _{oc}	No data	No data
Vapor pressure:	1 (1,628 °C) ^g	No data
Henry's law constant at 25 °C	No data	No data
Autoignition temperature	No data	No data
Flashpoint	No data	No data
Flammability limits	No data	No data
Conversion factors at 25 °C ppm to mg/m ³	Since these substances exist in the atmosphere in the particulate state, the concentration is expressed as mg/m ³ .	
Explosive limits	No data	No data

^aLide 2000^bLewis 1997^cBudavari et al. 2001^dMeister et al. 2001^eStewart and Lassiter 2001^fDean 1985^gLewis 2000pK_a = The dissociation constant of the conjugate acid

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

5.1 PRODUCTION

Copper occurs naturally in many minerals, such as cuprite (Cu_2O), malachite ($\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$), azurite ($2\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$), chalcopyrite (CuFeS_2), chalcocite (Cu_2S), and bornite (Cu_5FeS_4). It also occurs uncombined as the metal (Tuddenham and Dougall 1979; Weast 1980). The copper content of ore deposits ranges from 0.5 to 5% by weight, whereas in igneous rock copper content ranges from 0.0005 to 0.011% (Duby 1980; Weant 1985). The three most important sources of copper are chalcocite, chalcopyrite, and malachite (Weant 1985). The major U.S. deposits are porphyry, indicating that they are of hydrothermal origin and are uniformly distributed in fractures or veins.

The United States is the world's second leading copper producer. The country produced 148 million metric tons of mined ore in 2001, with an average copper content of 0.48% (USGS 2001). Mine production of recoverable copper in the United States totaled 1,340,000 metric tons in 2001, an estimated 10% of world production behind Chile, which accounted for 35%. Copper was mined in six states in 2001, with Arizona accounting for 67% of U.S. copper production, followed by Utah (13%), New Mexico (13%) and Nevada (1%). There were 23 copper-producing mines in 2001, down from 27 in 2000. Thirteen of these are copper mines accounting for 99% of production in the United States. The remaining mines yielded copper as a by-product of gold, lead, silver, or zinc mining. Of the 13 largest mines, 10 were in Arizona, 2 were in New Mexico, and 1 was in Utah. Production, processing and use of copper and copper compounds in the United States, listed by state, are given in Tables 5-1 and 5-2, respectively.

After mining, most of the ore is crushed and concentrated to a material containing 15–35% copper using flotation. The remaining copper is obtained by first leaching the ore or tailings and then concentrating the leachate by applying solvent extraction or ion exchange (Butterman 1982).

Most primary copper is produced from its sulfide ore by matte smelting, an operation yielding a molten sulfide of copper and iron, called matte, which is further oxidized in a conversion step to yield metallic copper. The conversion operation takes place in two stages. In the first, slag-forming stage, FeS is oxidized to iron oxides, which combine with a silica flux to form a slag. In the second, copper-producing stage, CuS_2 is oxidized to form sulfur dioxide and metallic copper. The product of the conversion

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

Table 5-1. Facilities that Produce, Process, or Use Copper

State ^a	Number of facilities	Minimum amount on site in pounds ^b	Maximum amount on site in pounds ^b	Activities and uses ^c
AK	2	10,000	99,999	12
AL	47	100	49,999,999	1, 2, 3, 5, 7, 8, 9, 11, 12, 13
AR	45	1,000	49,999,999	1, 2, 3, 4, 7, 8, 9, 12, 13, 14
AZ	26	0	999,999,999	1, 3, 4, 5, 7, 8, 9, 11, 12, 13, 14
CA	153	0	99,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14
CO	14	1,000	499,999,999	2, 3, 4, 7, 8, 11, 12, 14
CT	54	100	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 11
FL	28	1,000	9,999,999	2, 3, 4, 6, 7, 8, 9, 10, 11, 12
GA	58	0	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 13, 14
HI	2	1,000	9,999	12
IA	31	1,000	99,999,999	1, 2, 3, 4, 5, 7, 8, 9, 12
ID	3	10,000	999,999	1, 3, 8, 9, 12
IL	150	0	99,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 14
IN	148	0	499,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
KS	23	100	9,999,999	2, 3, 4, 6, 7, 8, 9, 11, 12, 14
KY	69	100	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13
LA	10	100	9,999,999	2, 6, 8, 10, 12, 13, 14
MA	59	1,000	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12
MD	7	1,000	999,999	1, 2, 4, 5, 7, 8, 9
ME	10	10,000	9,999,999	2, 3, 8
MI	133	0	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12
MN	49	100	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
MO	73	100	499,999,999	1, 2, 3, 4, 5, 7, 8, 9, 11, 12
MS	33	100	9,999,999	4, 7, 8, 9, 12
MT	1	1,000	9,999	6, 11
NC	72	0	99,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12
ND	2	10,000	99,999	7, 8
NE	18	1,000	9,999,999	1, 3, 5, 7, 8, 9, 11, 12, 13
NH	19	100	49,999,999	2, 3, 4, 7, 8, 9
NJ	36	1,000	49,999,999	1, 2, 3, 4, 6, 7, 8, 9, 11, 12
NM	7	1,000	9,999,999	2, 3, 8, 12
NV	6	1,000	99,999	7, 8, 11, 12
NY	100	0	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14
OH	204	100	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
OK	51	100	99,999,999	1, 2, 3, 4, 7, 8, 9, 11, 12, 13
OR	23	0	999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12
PA	207	0	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14
PR	22	1,000	9,999,999	1, 2, 3, 8, 11
RI	28	1,000	9,999,999	2, 3, 4, 6, 7, 8, 9, 12

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

Table 5-1. Facilities that Produce, Process, or Use Copper

State ^a	Number of facilities	Minimum amount on site in pounds ^b	Maximum amount on site in pounds ^b	Activities and uses ^c
SC	53	100	9,999,999	1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 14
SD	11	1,000	49,999,999	1, 5, 7, 8, 12, 14
TN	75	0	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14
TX	102	100	99,999,999	1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 12, 14
UT	10	1,000	9,999,999	1, 3, 4, 5, 6, 7, 8, 11, 12
VA	42	1,000	9,999,999	1, 2, 3, 5, 6, 7, 8, 10, 11, 12, 13, 14
VT	4	1,000	99,999	2, 3, 4, 6, 8, 9
WA	24	1,000	999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
WI	148	0	499,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 14
WV	12	1,000	9,999,999	2, 3, 6, 7, 8, 12
WY	4	0	99,999	1, 4, 9, 10, 12

Source: TRI01 2003

^aPost office state abbreviations used^bAmounts on site reported by facilities in each state^cActivities/Uses:

- | | | |
|--------------------------|--------------------------|-----------------------------|
| 1. Produce | 6. Impurity | 11. Chemical processing aid |
| 2. Import | 7. Reactant | 12. Manufacturing aid |
| 3. Onsite use/processing | 8. Formulation component | 13. Ancillary/Other uses |
| 4. Sale/Distribution | 9. Article component | 14. Process impurity |
| 5. Byproduct | 10. Repackaging | |

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

Table 5-2. Facilities that Produce, Process, or Use Copper Compounds

State ^a	Number of facilities	Minimum amount on site in pounds ^b	Maximum amount on site in pounds ^b	Activities and uses ^c
AK	5	10,000	9,999,999	1, 2, 3, 4, 5, 7, 10, 12, 13, 14
AL	37	0	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13
AR	26	100	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14
AZ	27	1,000	10,000,000,000	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14
CA	74	0	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
CO	7	100	99,999	1, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14
CT	24	1,000	999,999,999	1, 3, 5, 6, 7, 8, 9, 10, 11, 12
DC	1	1,000	9,999	12
DE	5	1,000	99,999	1, 2, 3, 5, 7, 9, 12, 13
FL	36	100	999,999	1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 12, 13, 14
GA	36	0	99,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13
IA	26	0	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
ID	8	1,000	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10
IL	92	0	99,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
IN	72	0	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
KS	11	0	999,999	1, 3, 5, 7, 8, 9, 10, 11, 12, 13
KY	39	100	499,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14
LA	26	0	99,999,999	1, 2, 3, 4, 5, 7, 8, 10, 11, 12, 13
MA	25	100	9,999,999	1, 3, 4, 5, 6, 7, 8, 10, 11, 12
MD	10	100	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 13
ME	5	100	999,999	1, 3, 5, 7, 8, 11, 13
MI	52	100	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14
MN	29	100	499,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
MO	34	0	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
MS	14	100	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 13
MT	9	1,000	99,999,999	1, 2, 3, 4, 5, 6, 8, 9, 10, 11, 12, 13, 14
NC	48	100	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
ND	8	100	99,999	1, 5, 7, 9, 12, 13, 14
NE	10	10,000	99,999,999	1, 3, 4, 5, 7, 8, 9, 10, 11, 12, 13, 14
NH	12	100	9,999,999	1, 3, 5, 6, 7, 8, 9, 11, 12, 13
NJ	24	0	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14
NM	8	1,000	99,999,999	1, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13
NV	17	100	10,000,000,000	1, 4, 5, 6, 7, 9, 10, 11, 12, 13, 14
NY	30	100	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
OH	83	0	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
OK	14	100	999,999	1, 3, 4, 5, 6, 8, 9, 10, 11, 12, 13, 14
OR	16	100	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12
PA	92	0	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
PR	3	1,000	99,999	7, 10

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

Table 5-2. Facilities that Produce, Process, or Use Copper Compounds

State ^a	Number of facilities	Minimum amount on site in pounds ^b	Maximum amount on site in pounds ^b	Activities and uses ^c
RI	12	100	999,999	1, 2, 3, 4, 7, 8, 10, 12
SC	31	0	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14
SD	2	1,000	999,999	1, 3, 5, 6, 10, 13
TN	47	100	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12
TX	89	0	99,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
UT	14	1,000	10,000,000,000	1, 3, 4, 5, 6, 7, 8, 9, 12, 13
VA	36	0	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
WA	13	1,000	999,999	1, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13
WI	35	0	999,999	1, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
WV	17	1,000	999,999	1, 3, 4, 5, 7, 8, 9, 12, 13, 14
WY	4	100	999,999	1, 5, 9, 12, 13

Source: TRI01 2003

^aPost office state abbreviations used^bAmounts on site reported by facilities in each state^cActivities/Uses:

- | | | |
|--------------------------|--------------------------|-----------------------------|
| 1. Produce | 6. Impurity | 11. Chemical processing aid |
| 2. Import | 7. Reactant | 12. Manufacturing aid |
| 3. Onsite use/processing | 8. Formulation component | 13. Ancillary/Other uses |
| 4. Sale/Distribution | 9. Article component | 14. Process impurity |
| 5. Byproduct | 10. Repackaging | |

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

operation is blister copper, which is 98.5–99.5% copper. Concentrated leachate from low-grade ore is subject to electrowinning, which electrolyzes aqueous sulfate solutions, or to cementation, which displaces copper from solution by a more active metal such as iron (Duby 1980). Further purification is obtained by electrolytic refining. For more details on copper mining, ore processing, smelting, and refining, see Duby (1980) and EPA (1980b).

Production of copper in the United States includes not only the processing of both domestic and foreign ores, but also the recovery of scrap. Scrap is a significant part of the U.S. copper supply. Scrap refers to both 'old scrap' (metal that has been used) and 'new scrap' (generated during fabrication). In 1999, smelting was performed in the United States by four primary smelters and two secondary smelters with a combined capacity of 1,750,000 metric tons per year (USGS 2000). Together, they produced 1,290,000 metric tons of copper from both domestic and foreign ores and scrap in 1999 (USGS 2001). In 2000, smelting was performed in the United States by four primary smelters and one secondary smelter with a combined capacity of 1,180,000 metric tons per year. Production of copper from U.S. smelters in 2000 was reported to be 1,000,000 metric tons, down from 1,290,000 metric tons in 1999 (USGS 2001). During 2000, 23 refineries operating with a combined capacity of 2,400,000 tons, produced 1,587,000 metric tons of copper from domestic and foreign ores. An additional 208,000 metric tons of copper was produced from new and old scrap for a combined total refinery production in the United States of 1,790,000 tons (USGS 2001). This level of refinery production was down from a level of 2,120,000 metric tons in 1999 (USGS 2001). Production of secondary copper and copper-alloys amounted to 1,490,000 metric tons in both 1999 and 2000 (USGS 2000). Apparent consumption for 2000 was 3,130,000 metric tons (USGS 2001). This includes domestic refined copper production, net imports of refined copper, copper recovered from old scrap, and stock adjustments. These alloys, primarily brass and bronze, contain approximately 60–90% copper.

Most industrially important copper compounds are made starting with copper metal. Copper sulfate, the most commercially important copper compound, was produced by at least six companies in plants in Casa Grande, Arizona; Sewaren and Oak Bridge, New Jersey; El Paso and Garland, Texas; Sante Fe Springs, California; Union, Illinois; Copperhill, Tennessee; and Sumter, South Carolina (Jolly and Edelstein 1987).

Copper sulfate also is produced as a by-product of copper production by ore-leaching with sulfuric acid. Production of copper sulfate increased by 29% from 1996 to 2000, standing at 55,500 metric tons in 2000

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

(USGS 2000). However, in 2001, production of copper sulfate decreased slightly to 55,200 metric tons (USGS 2001). Recent production figures for other copper compounds were not located.

5.2 IMPORT/EXPORT

In 2000, 1,340,000 million tons of unmanufactured copper and 1,060,000 metric tons of refined copper were imported (USGS 2001). Peru, Canada, and Chile were the principal sources of imported refined copper. The quantity of imported unmanufactured copper increased by 93% since 1994; the increase was almost entirely in the importation of refined powder, as opposed to ore concentrate, blister copper, or scrap (USGS 1994, 2001). Imports of copper sulfate amounted to 4,650 metric tons and were primarily obtained from Australia and Mexico (USGS 2001).

In 2000, 483,000 metric tons of copper were exported, of which 19% was refined copper (USGS 2000). In 2001, exports dropped to 379,000 metric tons with a large decrease in exports of refined copper (from 93,600 metric tons in 2000 to 22,500 metric tons in 2001); exports of unalloyed copper scrap increased from 228,000 metric tons in 2000 to 262,000 metric tons in 2001 (USGS 2001).

5.3 USE

Copper is one of the most important metals because of its durability, ductility, malleability, and electrical and thermal conductivity. It is used primarily as the metal or in alloys. Its alloys, including brass, bronze, gun metal, and Monel metal, are important commodities. All current American coins are copper alloys. A small percentage of copper production goes into the manufacture of copper compounds, primarily copper sulfate.

The Copper Development Association's 2000 estimates of the end-use distribution of copper and copper-alloy products by the industrial sector were: construction, 39%; electrical and electrical products, 28%; transportation equipment, 11%; industrial machinery and equipment, 11%; and consumer and general products, 11% (USGS 2002). The top 10 markets for copper and copper-alloy during 1986 were, in order of importance: plumbing, building wire, telecommunications, power utilities, in-plant equipment, air conditioning, automotive electrical, automotive nonelectrical, business electronics, and industrial valves and fittings (Jolly and Edelstein 1987).

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

Copper sulfate was the only copper compound for which end-use distribution data were available; these data addressed only domestic producers. Sixty-five percent of production went into agricultural use, 28% for industrial uses such as metal finishing, mineral froth flotation and wood preservatives, and 7% for water treatment.

In agriculture, copper compounds are used as fungicides and to prepare copper fungicidal products, algicides for reservoirs and streams and nutritional supplements in animal feed and fertilizers. Industrial applications of copper sulfate include use as an activator in froth flotation of sulfide ores, production of chromated copper arsenate wood preservatives, electroplating, azo dye manufacture, mordant for textile dyes, petroleum refining and in the manufacture of other copper compounds such as copper hydroxide and copper carbonate (Mannsville Chemical Products 1984).

Copper compounds are applied as fungicides to foliage, seed, wood, fabric, and leather to protect against blight, downy mildew and rust. The 1982 consumption of copper-containing fungicides was 2.8 million pounds (Mannsville Chemical Products 1984). The major copper compound used for this purpose was the basic copper sulfate (1.8 million pounds). Other important fungicidal compounds were copper hydroxide, copper ammonium carbonate, copper oxychloride and copper oxychloride sulfate. The major target crops of copper-containing fungicides are citrus fruits, peanuts, deciduous fruits (other than apples), potatoes, vegetables and other field crops. Copper compounds are also used as algicides, insecticides and repellents. Products containing copper compounds frequently contain other chemicals and may be sold under various trade names. Formulation may be in wettable powders or aqueous solutions.

5.4 DISPOSAL

It is estimated that 60% of copper in scrap is recycled (Tuddenham and Dougall 1978). In 1986, ~40% of the copper produced came from this source (Jolly and Edelstein 1987). Copper-containing wastes can be concentrated using ion exchange, reverse osmosis, or evaporation, and then reclaimed by electrolysis (HSDB 2002). Copper and copper compounds not recycled are disposed of in landfills or released into waste water. Methods of copper containing sludge disposal from waste water treatment facilities include landfilling, landspreading, incineration or ocean disposal.

In case of a solid copper sulfate spill on land, the solids should be protected from rain and fire-fighting water by covering the material with plastic sheeting (HSDB 2002). In the event of a water spill, the

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

copper sulfate should be neutralized with crushed limestone, slaked lime, or sodium bicarbonate, and the solidified masses should be removed.

6. POTENTIAL FOR HUMAN EXPOSURE


6.1 OVERVIEW

Copper has been identified in at least 906 of the 1,647 hazardous waste sites that have been proposed for inclusion on the EPA National Priorities List (NPL) (HazDat 2004). However, the number of sites evaluated for copper is not known. The frequency of these sites can be seen in Figure 6-1. Of these sites, 895 are located within the United States, 2 are located in the Territory of Guam, 8 are located in the Commonwealth of Puerto Rico, and 1 is located in the U.S. Virgin Islands (the sites in the Territory of Guam, the Commonwealth of Puerto Rico, and the U.S. Virgin Islands are not shown).

Copper and its compounds are naturally present in the earth's crust. Natural discharges to air and water, such as windblown dust, volcanic eruptions, etc., may be significant. Therefore, it is important to consider the copper concentrations within a specific environment, geographical region, or human population study site that has been minimally affected by anthropogenic sources of copper in order to accurately assess the contribution of an anthropogenic activity to human exposures to copper. In air, the mean copper concentrations in the atmosphere range between 5 and 200 ng/m³ in rural and urban locations. Airborne copper is associated with particulates that are obtained from suspended soils, combustion sources, the manufacture or processing of copper-containing materials, or mine tailings. The median concentration of copper in natural water (e.g., rivers, lakes, and oceans) is 4–10 ppb. It is predominantly in the Cu(II) state. Most of it is complexed or tightly bound to organic matter. Little is present in the free (hydrated) or readily exchangeable form. The combined processes of complexation, adsorption, and precipitation control the level of free Cu(II). The chemical conditions in most natural water are such that, even at relatively high copper concentrations, these processes will reduce the free Cu(II) concentration to extremely low values. The mean concentration of copper in soil ranges from 5 to 70 mg/kg and is higher in soils near smelters, mining operations, and combustion sources. Sediment is an important sink and reservoir for copper. In relatively clean sediment such as those found in some of the bays and estuaries along the New England Coast, the copper concentration is <50 ppm; polluted sediment may contain several thousand ppm of copper. The form of copper in the sediment also will be site-specific. In aerobic sediments, copper is bound mainly to organics (humic substances) and iron oxides. However, in some cases, copper is predominantly associated with carbonates. In anaerobic sediments, Cu(II) will be reduced to Cu(I) and insoluble cuprous salts will be formed.

6. POTENTIAL FOR HUMAN EXPOSURE

Figure 6-1. Frequency of NPL Sites with Copper Contamination



Derived from HazDat 2004

6. POTENTIAL FOR HUMAN EXPOSURE

The largest release of copper to the environment by anthropogenic activities is by far to land. The major sources of release are mining operations, agriculture, sludge from publicly-owned treatment works (POTWs) and municipal and industrial solid waste. Mining and milling contribute the most waste. Copper is released to water as a result of natural weathering of soil and discharges from industries and sewage treatment plants. Copper compounds may also be intentionally applied to water to kill algae.

Copper associated with particulate matter is emitted into the air naturally from windblown dust, volcanoes, and anthropogenic sources, the largest of which are being primary copper smelters and ore processing facilities. The concentration of copper in emissions from copper smelters has been found to range between 7 and 137.8 ng/m³ (Hutchinson 1979; Romo-Kröger et al. 1994).

In the general population, the highest exposures to copper come from drinking water and food. Of special concern is copper that gets into drinking water from water distribution systems (both from the water treatment plant and in the home). When a system has not been flushed after a period of disuse, the concentration of copper in tap water may exceed 1.3 ppm, the EPA drinking water limit. The estimated intakes of copper in the general population are 0.15 mg/day from drinking water, and approximately 2 mg/day from food. The dietary intake of copper can be increased from the regular consumption of certain foods, such as shellfish, organ meats (e.g., liver and kidney), legumes, and nuts. However, except for shellfish, where an additional intake of 2–150 mg/day is possible for those individuals who regularly consume shellfish, these other sources of higher copper intake are not expected to increase the total daily intake of copper beyond the recommended limit of 10–12 mg/day for adults (WHO 1996). In comparison to intake of copper through ingestion of water and food, the intake of copper through inhalation of copper in dust is much less significant at an estimated rate of 0.1–4.0 µg copper/day. Contact with available copper also may result from the use of copper fungicides and algicides.

Many workers are exposed to copper in agriculture, industries connected with copper production, metal plating, and other industries. Little information is available concerning the forms of copper to which workers are exposed.

At this time, copper has been identified in 906 out of 1,647 NPL hazardous waste sites in the United States (HazDat 2004). The frequency of these sites within the United States is noted in Figure 6-1. Based on the available data, people living close to NPL sites may be at greater risk for exposure to copper than the general population with respect to inhalation of airborne particulates from the NPL sites, ingestion of

6. POTENTIAL FOR HUMAN EXPOSURE

contaminated water or soil, and/or uptake of copper into fruits and vegetables raised in gardens of residents living near NPL sites. People living near copper smelters and refineries and workers in these and other industries may be exposed to high levels of dust-borne copper by both inhalation and ingestion routes. For example, ingestion of 300 mg of soils near copper smelters by children could result in the intake as high as 0.74–2.1 mg copper per day, based on measurements of copper concentrations in these soils of 2,480–6,912 mg/kg.

6.2 RELEASES TO THE ENVIRONMENT

Industrial manufacturers, processors, and users of copper and copper compounds are required to report the quantities of this substance released to environmental media annually (EPA 1988d). The data compiled in the Toxics Release Inventory (TRI01 2003) are for releases in 2001 to air, water, soil, and transfer of copper and copper compounds for offsite disposal. These data are summarized in Tables 6-1 (copper) and 6-2 (copper compounds). Total releases (rounded to three significant digits) of copper into the environment in 2001 were approximately 11,100,000 pounds (approximately 5,050 metric tons) (TRI01 2003), of which approximately 821,000 pounds (373 metric tons), or 7.4% of the total, were released to air. Another 46,600 pounds (21 metric tons) or approximately 0.4% of the total, were released into water, 0.5% (53,800 pounds, 24 metric tons) was injected underground, and 91.9% (10,200,000 pounds, 4,360 metric tons) was released to land. Total releases (rounded to three significant digits) of copper compounds to the environment in 2001 were approximately 1,000,000,000 pounds (approximately 455,000 metric tons) (TRI01 2003) of which approximately 1,420,000 pounds (645 metric tons), or 0.1% of the total, were released to air. Another 418,000 pounds (190 metric tons), or approximately 0.04% of the total, were released into water, 0.09% (894,000 pounds or 406 metric tons) was injected underground, and 99.8% (998,000,000 pounds or 454,000 metric tons) was released to land. The TRI data should be used with caution because only certain types of facilities are required to report them (i.e., this is not an exhaustive list).

Industrial releases are only a fraction of the total environmental releases of copper and copper compounds. Other sources of copper release into the environment originate from domestic waste water, combustion processes, wood production, phosphate fertilizer production, and natural sources (e.g., wind blown dust, volcanoes, decaying vegetation, forest fires, sea spray, etc.) (Georgopoulos et al. 2001;

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-1. Releases to the Environment from Facilities that Produce, Process, or Use Copper^a

State ^c	Number of facilities	Reported amounts released in pounds per year ^b						
		Air ^d	Water	Under-ground injection	Land	Total on-site release ^e	Total off-site release ^f	Total on and off-site release
AK	2	0	No data	0	27,324	27,324	0	27,324
AL	51	11,825	1,321	0	14,906	28,052	47,708	75,760
AR	48	5,951	573	0	143,970	150,494	231,861	382,355
AZ	30	978	48	0	227,993	229,019	9,094	238,113
CA	168	9,745	908	0	2,608,971	2,619,624	171,791	2,791,415
CO	18	472	16	0	125,387	125,875	15,991	141,866
CT	63	10,662	904	0	5	11,571	47,415	58,985
FL	35	832	303	51,262	158,561	210,958	40,092	251,050
GA	72	4,473	575	0	135,127	140,175	30,052	170,227
HI	2	0	No data	0	117,010	117,010	0	117,010
IA	39	3,050	566	0	250	3,866	35,144	39,010
ID	5	120	5	0	450,820	450,945	4,000	454,945
IL	160	46,380	4,623	0	1,601,420	1,652,423	658,745	2,311,168
IN	161	42,775	1,239	0	369,664	413,678	2,223,563	2,637,241
KS	24	2,535	251	0	297,190	299,976	24,566	324,542
KY	73	18,633	390	0	269,927	288,950	179,301	468,251
LA	15	126,947	710	2,200	2,426	132,283	690	132,973
MA	69	2,530	77	0	0	2,607	64,821	67,427
MD	12	250	10	0	250	510	89,168	89,678
ME	11	106	321	0	500	927	9,034	9,961
MI	142	35,480	685	0	849	37,014	161,721	198,735
MN	50	16,129	10	0	5	16,144	825,881	842,025
MO	79	9,204	671	0	52,581	62,456	409,753	472,209
MS	35	2,350	657	0	520	3,527	56,712	60,239
MT	1	161	No data	0	1,030,000	1,030,161	No data	1,030,161
NC	83	7,965	1,076	0	193,576	202,617	107,899	310,516
ND	3	23	5	0	0	28	339	367
NE	19	3,638	5	0	47,005	50,648	7,628	58,276
NH	22	1,005	25	0	0	1,030	37,543	38,573
NJ	46	15,584	165	5	22,973	38,727	12,291	51,018
NM	7	500	No data	0	118,680	119,180	23,453	142,633
NV	6	905	No data	0	500	1,405	5,201	6,606
NY	104	11,484	14,198	0	114,462	140,144	894,995	1,035,138
OH	251	55,122	5,424	0	606,137	666,683	481,776	1,148,459
OK	64	8,712	303	0	80,952	89,967	29,726	119,693
OR	23	771	9	0	81,984	82,764	1,495	84,259
PA	222	85,511	2,913	0	43,546	131,970	343,353	475,324

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-1. Releases to the Environment from Facilities that Produce, Process, or Use Copper^a

State ^c	Number of facilities	Reported amounts released in pounds per year ^b						Total on and off-site release
		Air ^d	Water	Under-ground injection	Land	Total on-site release ^e	Total off-site release ^f	
PR	21	15,944	5	0	0	15,949	9,247	25,196
RI	33	5,093	5	0	0	5,098	22,759	27,857
SC	55	10,018	875	0	76,761	87,654	59,678	147,332
SD	10	4,885	No data	0	750	5,635	280	5,915
TN	89	165,328	567	0	206	166,101	87,117	253,218
TX	117	26,388	4,006	369	574,322	605,085	178,081	783,166
UT	12	1,179	56	0	46,085	47,320	298	47,618
VA	46	21,649	694	5	239,776	262,124	210,846	472,970
VT	5	0	No data	0	250	250	1,025	1,275
WA	28	474	633	0	166,530	167,637	150,773	318,410
WI	159	25,741	617	0	26,845	53,203	257,886	311,089
WV	12	1,951	116	5	30,759	32,831	8,102	40,933
WY	5	381	0	0	60,882	61,263	75	61,338
Total	2,807	821,838	46,559	53,846	10,168,637	11,090,881	8,268,966	19,359,847

Source: TRI01 2003

^aThe TRI data should be used with caution since only certain types of facilities are required to report. This is not an exhaustive list. Data are rounded to nearest whole number.

^bData in TRI are maximum amounts released by each facility.

^cPost office state abbreviations are used.

^dThe sum of fugitive and stack releases are included in releases to air by a given facility.

^eThe sum of all releases of the chemical to air, land, water, and underground injection wells.

^fTotal amount of chemical transferred off-site, including to publicly owned treatment works (POTW).

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-2. Releases to the Environment from Facilities that Produce, Process, or Use Copper Compounds^a

State ^c	Number of facilities	Reported amounts released in pounds per year ^b						
		Air ^d	Water	Under-ground injection	Land	Total on-site release ^e	Total off-site release ^f	Total on and off-site release
AK	6	475	57	670,000	5,193,578	5,864,110	750	5,864,860
AL	66	27,363	28,896	0	2,315,767	2,372,026	66,409	2,438,435
AR	60	17,082	3,277	0	86,510	106,869	173,666	280,535
AZ	27	135,813	584	0	451,467,272	451,603,669	76,083	451,679,752
CA	92	5,374	878	0	111,029	117,281	166,771	284,053
CO	15	720	15,810	0	115,190	131,720	61,086	192,806
CT	24	2,072	635	0	0	2,707	354,565	357,272
DC	1	0	0	0	4,600	4,600	0	4,600
DE	9	3,203	9,196	0	25,029	37,428	30,509	67,937
FL	54	85,273	22,720	0	648,038	756,031	169,746	925,777
GA	69	14,994	48,943	0	776,472	840,409	369,367	1,209,776
IA	52	15,425	3,386	0	156,575	175,386	152,607	327,993
ID	11	1,305	800	0	424,736	426,841	272	427,113
IL	114	46,746	5,233	0	503,051	555,030	1,021,024	1,576,054
IN	88	52,013	20,411	250	1,098,611	1,171,285	1,044,431	2,215,716
KS	20	3,774	0	0	247,277	251,051	112,520	363,571
KY	49	37,298	40,320	0	889,385	967,003	529,758	1,496,761
LA	33	5,626	17,074	7	274,009	296,716	156,664	453,380
MA	27	720	39	0	3	762	104,579	105,341
MD	19	7,471	8,993	0	20,391	36,855	163,428	200,283
ME	5	2,200	485	0	0	2,685	37,187	39,872
MI	68	60,059	14,850	0	643,734	718,643	511,519	1,230,162
MN	46	9,598	882	0	281,020	291,500	2,115,492	2,406,992
MO	55	20,270	2,900	0	4,658,137	4,681,307	248,752	4,930,059
MS	35	45,892	279	12,000	22,708	80,879	54,015	134,894
MT	10	12,595	10	47,757	3,385,422	3,445,784	32,403	3,478,187
NC	88	21,656	15,299	0	758,429	795,384	124,485	919,869
ND	8	632	11,877	0	138,516	151,025	124,852	275,877
NE	21	1,999	155	0	207,272	209,426	21,672	231,098
NH	13	561	12	0	0	573	28,554	29,127
NJ	30	1,187	5,754	0	18,117	25,058	1,869,339	1,894,397
NM	10	11,839	4,005	0	55,651,587	55,667,431	43,010	55,710,441
NV	18	2,034	160	1	27,004,822	27,007,017	3,067	27,010,084
NY	35	9,816	9,830	1	52,320	71,967	161,344	233,311
OH	105	14,459	17,520	8,100	1,158,856	1,198,935	1,555,185	2,754,120
OK	23	2,466	5,082	662	212,835	221,045	48,189	269,234
OR	25	2,160	1,021	0	216,646	219,827	19,130	238,957

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-2. Releases to the Environment from Facilities that Produce, Process, or Use Copper Compounds^a

State ^c	Number of facilities	Reported amounts released in pounds per year ^b						
		Air ^d	Water	Under-ground injection	Land	Total on-site release ^e	Total off-site release ^f	Total on and off-site release
PA	115	484,099	14,493	0	371,008	869,600	3,121,602	3,991,202
PR	7	44	265	0	0	309	0	309
RI	13	253	608	0	108	969	2,927	3,896
SC	53	32,578	2,664	0	203,101	238,343	309,623	547,966
SD	5	6,150	1,340	0	88,000	95,490	18	95,508
TN	60	17,676	20,876	0	11,178,079	11,216,631	336,292	11,552,923
TX	124	99,287	12,731	155,405	1,302,462	1,569,885	849,639	2,419,524
UT	18	63,960	2,860	0	424,682,491	424,749,311	36,408	424,785,719
VA	50	10,518	19,707	0	390,234	420,459	184,428	604,887
WA	23	6,200	756	0	212,242	219,198	35,097	254,295
WI	56	7,319	13,315	0	59,775	80,409	221,765	302,174
WV	18	4,390	11,557	0	769,830	785,777	199,314	985,091
WY	5	1,470	118	0	235,587	237,175	46,668	283,843
Total	1,978	1,416,114	418,663	894,183	998,260,861	1,000,989,821	17,096,211	1,018,086,032

Source: TRI01 2003

^aThe TRI data should be used with caution since only certain types of facilities are required to report. This is not an exhaustive list. Data are rounded to nearest whole number.

^bData in TRI are maximum amounts released by each facility.

^cPost office state abbreviations are used.

^dThe sum of fugitive and stack releases are included in releases to air by a given facility.

^eThe sum of all releases of the chemical to air, land, water, and underground injection wells.

^fTotal amount of chemical transferred off-site, including to publicly owned treatment works (POTW).

6. POTENTIAL FOR HUMAN EXPOSURE

Harrison 1998). Quantitative information on release of copper to specific environmental media is discussed below. A summary of copper concentrations in environmental media is provided in Table 6-3.

6.2.1 Air

Copper is emitted into the air from both natural and anthropogenic sources. Since copper is a component of the earth's crust, the earth's crust is the primary natural source of copper. Windblown dust has an estimated mean worldwide emission of $0.9\text{--}15 \times 10^6$ kg/year of copper into the atmosphere (WHO 1998). Other natural sources of copper emitted into air (in terms of estimated ranges of worldwide emissions) are forest fires ($0.1\text{--}7.5 \times 10^6$ kg/year), volcanoes ($0.9\text{--}18 \times 10^6$ kg/year), biogenic processes ($0.1\text{--}6.4 \times 10^6$ kg/year), and sea spray ($0.2\text{--}6.9 \times 10^6$ kg/year) (WHO 1998). Based on these data, the mean total non-crustal sources of copper emitted into the atmosphere is $1.3\text{--}38.8 \times 10^6$ kg/year. Anthropogenic emission sources include nonferrous metal production, wood production, iron and steel production, waste incineration, industrial applications, coal combustion, nonferrous metal mining, oil and gasoline combustion, and phosphate fertilizer manufacture. It is estimated that only 0.04% of copper released to the environment is released into the air (Perwak et al. 1980). Global atmospheric anthropogenic and natural emissions of copper have been estimated to be 35×10^6 and 28×10^6 kg/year, respectively (Giusti et al. 1993; Nriagu 1989; Nriagu and Pacyna 1988). The estimates for the anthropogenic and natural emissions are based on the sum of copper emissions from various sources as shown in Tables 6-4 and 6-5, respectively.

The EPA conducted a detailed study of the total amount of copper emitted into the atmosphere (Weant 1985). The sources of emissions and the estimated quantities of copper emitted in 10^6 kg/year are: primary copper smelters, 0.043–6; copper and iron ore processing, 0.480–0.660; iron and steel production, 0.112–0.240; combustion sources, 0.045–0.360; municipal incinerators, 0.0033–0.270; secondary copper smelters, 0.160; copper sulfate production, 0.045; gray iron foundries, 0.079; primary lead smelting, 0.0055–0.065; primary zinc smelting, 0.024–0.340; ferroalloy production, 0.0019–0.0032; brass and bronze production, 0.0018–0.036; and carbon black production, 0.013. Using the ranges of copper emitted from these sources, it is estimated that U.S. copper emissions into air are $0.9424\text{--}7.974(x10^6)$ kg per annum. Daily stack emission rates have been reported for three coal-burning power plants on a kg/day/1,000 megawatt basis (Quee Hee et al. 1982); they are 0.3–0.7 and 2.00 kg/day/1,000 megawatt for those using low-sulfur western coal and high-sulfur eastern coal, respectively. This amounts to annual emission rates of 110–260 and 730 kg/1,000 megawatt, respectively. In another report, emission of copper into air from a 650 megawatt electrical power plant,

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-3. Summary of Copper Concentrations in Environmental Media^a

Environmental media		Concentration	Units
Atmosphere			
Aerosol		0.1–382	ppt
Hydrosphere—water			
Coastal	Dissolved	0.06–4.3	ppb
	Total	0.5–13.8	ppb
	Suspended solids	0.6–370,000	ppm
Estuarine	Dissolved	0.02–4.7	ppb
	Total	1.2–71.6	ppb
	Suspended solids	0.38–72	ppm
Ocean	Dissolved	Not detected–10	ppb
	Total	0.04–10	ppb
	Suspended solids	0.01–2.8	ppm
Lake	Dissolved	0.1–15.6	ppb
	Total	0.1–15.6	ppb
River	Dissolved	0.18–3,000	ppb
	Total	0.5–5,800	ppb
Groundwater	Dissolved	0.003–70	ppb
	Total	1–1,160	ppb
Drinking water	Total	0.3–1,352	ppb
Hydrosphere—sediments			
Coastal	Particulate	0.03–3,789	ppm
	Interstitial water	25.5–32.7	ppb
Estuarine	Particulate	0.3–2,985	ppm
	Interstitial water	0.3–100	ppb
Ocean	Particulate	3.1–648	ppb
	Interstitial water	22–45	ppm
Lake	Particulate	0.4–796	ppm
	Interstitial water	45.6–52	ppb
River	Particulate	5.3–4,570	ppm
Pedosphere			
Soil	Total	0.01–3,138	ppm
	Organic	293–7,634	ppm
Dust	Total	2.9–76	ppm

^aAs reported in the Copper Sourcebook 1998 (Harrison 1998), covering the years 1993–1996.

Source: Georgopoulos et al. 2001

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-4. Global Emissions of Copper from Natural Sources (x10⁶ kg/year)

	Median	Range
Wind-borne particulates	8.0	0.9–15
Marine spray—seasalt and surface organic microlayers	4.0	0.25–7.7
Volcanoes	9.4	0.9–18
Forest fires	3.8	0.1–7.5
Biogenic—continental particulates and volatiles	2.9	0.11–5.6
Total emissions	28	2.3–54

Source: Nriagu 1989

6. POTENTIAL FOR HUMAN EXPOSURE

**Table 6-5. Global Emissions of Copper from Anthropogenic Sources
(x10⁶ kg/year)**

	Median	Range
Coal combustion	5.15	2.3–8.0
Oil combustion	1.86	0.42–3.3
Pyrometallurgical	23.5	15–32
Secondary nonferrous metal production	0.115	0.06–0.17
Steel and iron manufacturing	1.47	0.14–2.8
Refuse incineration	1.5	1.0–2.0
Phosphate fertilizers	0.415	0.14–0.69
Wood combustion	0.9	0.60–1.2
Total emissions	35	20–51

Source: Nriagu and Pacyna 1988

6. POTENTIAL FOR HUMAN EXPOSURE

burning bituminous coal, was estimated at 213 kg/year, based on a summary of reportable TRI releases (Rubin 1999).

Emission factors in grams of copper released to the atmosphere per ton of product have been estimated for various industries (Nriagu and Pacyna 1988). These factors would enable estimation of an industry's copper emissions from its production volume. Missing from these emission estimates is fugitive dust arising from drilling, blasting, loading, and transporting operations associated with copper mining. The only control for reducing fugitive dust is the manual use of water sprays (EPA 1980b). The highest concentrations of copper in atmospheric particulate matter were obtained from mining activities, primary and secondary production, and industrial manufacturing (Table 6-6).

Romo-Kröger et al. (1994) were able to show, through the use of radioactive tracers and cluster analysis of interelemental correlations, that Cu, S, Zn, and As measured near a copper smelter in Chile were derived from the plant and not from the surrounding soil. The concentration of copper in air near the plant decreased from 66 ng/m³ (fine particles) and 131 ng/m³ (coarse particles) to 22 ng/m³ (fine particles) and 50 ng/m³ (coarse particles) during a period of inactivity at the plant, clearly demonstrating the contribution of plant emissions to copper levels in the surrounding area.

The amount of copper and other pollutants in fugitive dust originating from copper production sites, such as from smelter bag houses, or waste sites, is of some concern. In one study, the amount of airborne copper and other heavy metals deposited near a large refuse dump that received municipal and industrial waste and sewage sludge was determined by first measuring the amount of the metal accumulated in moss bags suspended 1–3 meters above the ground. The deposition rate was then determined from the amount of copper in the moss bags accumulated over the summer of 1985 and compared with that for an agricultural control area. The mean copper deposition rates in the two areas were about the same, 0.55 mg/kg-month (range of 0.04–1.6 mg/kg-month) over the refuse dump and 0.51 mg/kg-month (range of 0.26–0.76 mg/kg-month) in the control area (Lodenus and Braunschweiler 1986).

In a study of automobile exhaust emitted from light duty vehicles conducted in Denver, Colorado, it has been shown that this source of copper emission makes a small local contribution to copper in air. The amount of copper emitted in the exhaust from automobiles powered by regular gasoline has been measured to be 0.001–0.003 mg/mile driven using the Urban Dynamometer Driving Schedule (UDDS) of the Federal Test Schedule (FTS) during the summer of 1996 and the winter of 1997 (Cadle et al. 1999).

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-6. Concentrations of Copper in Particulate Matter (<10 µm) Generated from Various Sources^a

Source	Copper concentration (percent, w/w)
Metal mining	6.17 ^b
Secondary metal production	4.6 ^b
Primary metal production	3.50 ^b
Industrial manufacturing	2.16 ^b
Steel production	0.55 ^b
Gray iron foundries	0.19 ^b
Steel foundry, general	0.17 ^b
Solid waste	0.09 ^b
Food and agriculture	0.05 ^b
Chemical manufacturing	0.03 ^b
Petroleum industry	0.03 ^b
Gasoline vehicle exhaust	0.05 ^c
Paved road dust	0.0162 ^c
Construction dust	0.0102 ^c
Landfill dust	0.0102 ^c
Unpaved road dust	0.0087 ^c
Agricultural lands, dust	0.0067 ^c
Diesel vehicle exhaust	0.003 ^c

^aValues obtained from CEIDARS 2000

^bData obtained from USEPA Speciate 3.0; Shareef, G.S; Radian, September, 1987

^cData obtained from KVB Literature Search

6. POTENTIAL FOR HUMAN EXPOSURE

Diesel powered vehicles were also studied and found to emit 0.005–0.039 mg of copper per mile driven for vehicles using #2 diesel fuel.

Only in a few cases has the form of copper released into the air been determined. Copper released into the atmosphere will be in particulate matter in the elemental form or in the form of an oxide, sulfate, or carbonate. Because copper smelters co-emit SO_x gases, copper is expected to be released largely as the sulfate in particulate matter from these facilities. Combustion processes are reported to release copper into the atmosphere as the oxide, elemental copper, and adsorbed copper. Cupric oxide has been identified in emissions from steel manufacturing and in fly ash from oil-fired power plants and open-hearth steel mills (Graedel 1978; Perwak et al. 1980). Copper associated with fine particles (<1 µm) tends to result from combustion, while that associated with large particles (>10 µm) is likely to originate from wind blown soil and dust (Schroeder et al. 1987).

Copper was detected in air at 39 of the 906 NPL hazardous waste sites where copper has been detected in environmental media (HazDat 2004). Copper was detected in offsite air samples at concentrations ranging from 0.02 to 10 µg/m³ (median concentration of 0.38 µg/m³) (HazDat 2002). These copper concentrations in air are generally above the annual atmospheric concentrations of 0.005–0.2 µg/m³ (EPA 1987a).

6.2.2 Water

Much of the copper that enters environmental waters will be associated with particulate matter. Copper is a natural constituent of soil and will be transported into streams and waterways in runoff either due to natural weathering or anthropogenic soil disturbances. Sixty-eight percent of releases of copper to water is estimated to derive from these processes. Copper sulfate use represents 13% of releases to water and urban runoff contributes 2% (Perwak et al. 1980). In the absence of specific industrial sources, runoff is the major factor contributing to elevated copper levels in river water (Nolte 1988). In the EPA-sponsored National Urban Runoff Program, in which 86 samples of runoff from 19 cities throughout the United States were analyzed, copper was found in 96% of samples, at concentrations of 1–100 µg/L (ppb) with a geometric mean of 18.7 µg/L (Cole et al. 1984). This mean concentration of copper in runoff water is higher than the geometric mean concentration of 4.2 ppb for copper in surface water based on measurements in EPA's STORET database (Eckel and Jacob 1988). Of the 71 priority pollutants analyzed, copper, along with lead and zinc, was the most frequently detected.

6. POTENTIAL FOR HUMAN EXPOSURE

Giusti et al. (1993) provided estimates of global anthropogenic and natural copper inputs into oceans that are derived from two sources, atmospheric deposition and riverine input. Atmospheric input has been estimated at $14\text{--}45 \times 10^6$ kg/year for copper in a dissolved form (e.g., rainwater) and $2\text{--}7 \times 10^6$ kg/year for copper in a particulate form (e.g., aerosols). Riverine input is estimated to be 10×10^6 kg/year as dissolved copper and $1,500 \times 10^6$ kg/year as copper bound to particulates.

Domestic waste water is the major anthropogenic source of copper in waterways (Isaac et al. 1997; Nriagu and Pacyna 1988). Studies in Cincinnati and St. Louis showed discharges of copper into sewer systems from residential areas to be significant, with an average loading of 42 mg/person/day (Perwak et al. 1980). In a more comprehensive review, Jenkins and Russell (1994) reported a range of average copper loadings derived from residential and some small industrial contributions of 2.8–83 mg/person/day. Concentrations of copper in influents to 239 waste water treatment plants (12,351 observations) were 0.0001–36.5 ppm, and the median value was ~0.4 ppm (Minear et al. 1981). Copper is not entirely removed in POTWs, and releases from these facilities contribute ~8% of all copper released to water (Perwak et al. 1980). Inputs into the Narraganset Bay, Rhode Island, in decreasing order of importance, are sewage effluent, rivers, urban runoff, and atmospheric fallout (Mills and Quinn 1984; Santschi et al. 1984). Ninety percent of both dissolved and particulate copper was from the effluent of sewage treatment plants that discharged into the Providence River.

While some copper is removed from the waste stream by sewage treatment facilities, considerable copper remains in the effluent and is released into receiving waters (EPA 1981; Perwak et al. 1980). Because removal efficiencies for copper from waste streams tend to remain constant rather than proportional to influent copper concentrations, increases in copper concentrations in POTW influent streams will also result in increased copper concentrations in the effluent streams (Isaac et al. 1997). The copper in domestic waste water has been found to make up a substantial fraction of the copper found in POTW influent in the waste water systems of four Massachusetts municipalities. The range of removal efficiencies reported for pilot and full scale plants suggests that removal depends strongly on plant operation or influent characteristics.

A source of copper released into waterways is from urban storm water runoff. Copper in storm water runoff originates from the sidings and roofs of buildings, various emissions from automobiles, and wet and dry depositional processes (Davis et al. 2001). Concentrations of between 1 and 100 $\mu\text{g/L}$ of copper in storm water runoff have been measured (Georgopoulos et al. 2001). Storm water runoff normally contributes approximately 2% to the total copper released to waterways. In contrast, copper in runoff that

6. POTENTIAL FOR HUMAN EXPOSURE

is obtained from the natural weathering of soil or is release from disturbed soils contributes 68% of the copper released to waterways (Georgopoulos et al. 2001).

The best data on typical POTWs using secondary treatment show that 55–90% of copper is removed in these plants with a median and mean removal efficiency of 82% (Perwak et al. 1980). By contrast, those plants using only primary treatment had a 37% median removal efficiency. A more recent study focused on heavy metal removal in three POTWs that received primarily municipal sewage and used activated sludge as a secondary treatment. The study looked at removals in both the primary and secondary treatment stage. The mean removals of soluble copper and total copper after secondary treatment were 49–82 and 83–90%, respectively. The average copper concentration in the final effluent was 17–102 ppb, which would amount to an output of between 0.58 and 3.47 kg of copper into receiving waters per day, based on an effluent volume of 34,000 cubic meters (9 million gallons) per day (Aulenbach et al. 1987; Stephenson and Lester 1987).

Overflow outfalls within combined sewer systems (e.g., combination of domestic and industrial waste water plus storm water) are the primary sources of copper pollutants entering estuaries and other coastal areas of the United States (Crawford et al. 1995; Georgopoulos et al. 2001; Huh 1996; Iannuzzi et al. 1997). For example, Crawford et al. (1995) compiled a summary of the sources of various metals and other contaminants into the Newark Bay estuary. The mass loadings of copper into the estuary as a function of source are (in kg/day): municipal treatment systems, 103.4; industry direct discharge, 8.82; combined sewer overflows, 48.0; storm-water runoff, 62.2; tributary flow, 39.1 and discharges from the Passaic Valley Commission and Middlesex County Sewerage Authority, 126.5.

Discharges to water from active mining and milling are small and most of the western operations do not release any water because water is a scarce resource and is recycled (Perwak et al. 1980). Discharges from electroplating operations are either made directly to the water environment or indirectly via POTWs. Runoff from abandoned mines is estimated to contribute 314 metric tons annually to surface water (Perwak et al. 1980). These discharges are primarily insoluble silicates and sulfides and readily settle out into stream, river, or lake beds. Releases from manufactured products containing copper may be substantial, but are difficult to predict. Corrosion of copper in plumbing or construction may result in direct discharges or runoff into waterways. Copper and brass production releases relatively little copper to water.

6. POTENTIAL FOR HUMAN EXPOSURE

Waste water generated from copper mining operations comes from seepage, runoff from tailing piles, or utility water used for mine operation. The amount of waste water generated ranges from 0–300 L water/metric ton of ore mined for open pit copper mines and 8–4,000 L water/metric ton of ore mined underground (EPA 1980b). Copper concentrations in waste water from a selected open pit and underground copper mine were 1.05 and 0.87 ppm, respectively. Data regarding copper concentrations in waste water associated with selected concentrating, smelting, and refining operations can be found in EPA (1980b). Drainage from mining operations and abandoned mines has been shown to have an effect on copper content in local surface waters (see Table 6-7) with concentrations as high as 69,000 ppb being measured (Rösner 1998).

Results of an EPA industrial effluent survey show that mean and maximum levels of copper in treated waste water from six industries exceeded 1 and 10 ppm, respectively (EPA 1981). These industries and their mean and maximum discharges in ppm are: inorganic chemicals manufacturing (<1.6, 18); aluminum forming (<160, 2,200); porcelain enameling (1.3, 8.8); gum and wood chemicals (1.4, 3.0); nonferrous metals manufacturing (1.4, 27.0) and paint and ink formulation (<1.0, 60.0). Emission factors in nanograms of copper released per L of water outflow have been estimated for various industries. These factors would enable estimation of an industry's copper releases if the discharge volumes were known (Nriagu and Pacyna 1988).

Effluents from power plants that use copper alloys in the heat exchangers of their cooling systems discharge copper into receiving waters (Harrison and Bishop 1984). The largest discharges occur after start-up and decrease rapidly thereafter. At the Diablo Canyon Nuclear Power Station, a very high start-up discharge containing 7,700 ppb of copper fell to 67 ppb after 24 hours (Harrison et al. 1980). During normal operation at two nuclear power stations 6.5×10^6 cubic meters (1,700 million gallons) of seawater per day is used as cooling water for these facilities and discharged into the ocean with copper levels in the effluent ranging between 0.6 and 3.3 ppb (Harrison et al. 1980). This amounts to a total output of copper in the discharged seawater of 3.9–42 kg per day or 1,400–15,000 kg/annum from these two power plants. Except for after start-up of the cooling system, most of the soluble copper (that which passes through a 0.45 μm filter) discharged was in bound forms (Harrison et al. 1980). During normal operation, <20% of the copper released was in the <1,000 molecular weight fraction, which contains the more available copper species.

Copper sulfate is added directly to lakes, reservoirs, and ponds for controlling algae. However, the copper concentration in the water column generally returns to pretreatment levels within a few days

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-7. Concentrations of Copper in Water

Sample type/ source	Location	Concentration (ppb) Range (mean) [median]	Comments	Reference
Drinking water				
Private wells	Nova Scotia, four communities	40–200 130–2,450, 53% of samples >1,000 ppm	at tap, running water at tap, standing water	Maessen et al. 1985
Private wells	New Bedford, Massachusetts	(330)	at tap, running water	Yannoni and Piorkowski 1995
Not specified	Seattle, Washington	(160) (450), 24% of samples >1,000 ppm	running water standing water	Maessen et al. 1985
River water	Canada (National Survey)	≤5–530 [≤5] ≤5–100 [≤5] ≤5–220 [20]	raw water treated water distributed water	Meranger et al. 1979
Lake water	Canada (National Survey)	≤5–80 [≤5] ≤5–100 [≤5] ≤5–560 [40]	raw water treated water distributed water	Meranger et al. 1979
Well water	Canada (National Survey)	≤5–110 [≤5] ≤5–70 [≤5] 10–260 [75]	raw water treated water distributed water	Meranger et al. 1979
School drinking water	New Jersey	BD–10,200 ^a BD–7,800 BD–8,500	first draw 10-minute flush mid-day, first draw	Murphy 1993
Municipal water supply	Berlin, Germany	0.009–4.2 (0.561)	at tap, running water	Zietz et al. 2003a
Municipal water supply	Lower Saxony, Germany	<0.1–6.40 (0.183) <0.1–3.00 (0.106)	at tap, standing water at tap, running water	Zietz et al. 2003b
Groundwater				
Representative sample	New Jersey	[5.0]	1,063 samples, 90 th percentile 64.0 ppb, maximum 2,783 ppb, groundwater may or may not be used for drinking water	Page 1981
Shallow monitoring well	Denver, Colorado	<1–14 [2]	30 monitoring wells, 22 with PVC casings and 8 with metal casings; samples obtained after purging well 20 minutes	Bruce and McMahon 1996
Surface water				
U.S. Geological Survey stations	United States	(4.2) [4.0]	53,862 occurrences	Eckel and Jacob 1988

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-7. Concentrations of Copper in Water

Sample type/ source	Location	Concentration (ppb) Range (mean) [median]	Comments	Reference
Representative sample	New Jersey	[3.0]	590 samples, 90th percentile 9.0 ppb, maximum 261 ppb	Page 1981
Surface, marine	East Arctic Ocean	(0.126)	26 locations 0.5–1 m depth	Mart and Nurnberg 1984
Surface, marine	Atlantic Ocean	0.0572–0.0210	20 sites, 2 cruises, 0– 1 m depth	Yeats 1988
Pond	Massachusetts	<10–105	Low in summer, high in winter	Kimball 1973
Lakes	Canada	1–8 (2)	Acid sensitive lakes	Reed and Henningson 1984
Lakes	Great Lakes	629–834 (756) 703–1,061 (870) 540–1,098 (830)	Lake Superior Lake Erie Lake Ontario	Nriagu et al. 1996
	Representative samples, nearby to acidic mine drainage	32-1,200 (736)	12 samples taken from streams and ponds near abandoned coal mines in Indiana	Allen et al. 1996
	Representative samples from copper mining areas in Arizona	100–69,000 [1,200]	Samples obtained from the Cerbat Mountains mining area; 15 surface water sites with 14 sites downstream from old tailings and adits	Rösner 1998

^aBD = below detection limit

PVC = Polyvinyl chloride

6. POTENTIAL FOR HUMAN EXPOSURE

(Effler et al. 1980; Perwak et al. 1980). The reduction in dissolved copper during this period was accompanied by an increase in particulate copper (e.g., sorption to algae or other organic matter, which settles into the sediments of these bodies of water). The copper in the settled particulates is in equilibrium with the water column, which greatly favors copper in a bound state.

A potential source of copper release into waterways is leachate from municipal landfills. Copper concentrations in leachate obtained from waste sites have been found to vary widely. For example, copper concentrations in leachate from municipal landfills have been found to range from 0.005 to 1,110 ppm (Christensen et al. 1994; Perwak et al. 1980; Roy 1994). Although copper was measured in these leachates, its origin may not be from copper contained within the waste site, but from the surrounding soils. Cyr et al. (1987) reported that leachate from three municipal landfills in New Brunswick, Canada, did not contain copper concentrations significantly above those in control samples representing the surrounding soil types. Therefore, the emissions of copper from landfills into leachates should be made relative to the contribution of copper from surrounding soils, as determined from appropriately selected control samples.

Copper can enter surface waters as a result of agricultural runoff. For example, estimated loading rates of copper into surface water from irrigation water runoff near the Stillwater National Wildlife Refuge ranged from 0.307 to 8.34 mg/hour, depending on what period of the irrigation season samples were taken (Kilbride et al. 1998). The highest loading rates were obtained during the middle period (August through mid-September) of the irrigation season. The copper in the runoff water was found to be predominantly bound to drift material in the water (e.g., algae, vascular plants, invertebrates, vertebrates, and detrital material).

Copper was detected in groundwater and surface water at 558 and 308 of the 906 NPL hazardous waste sites, respectively, where copper has been detected in environmental media (HazDat 2004). Copper was detected at concentrations ranging from 0.006 to 5.6 ppm (median concentration of 0.103 ppm) in offsite groundwater and 0.00025–590 ppm (median concentration of 0.0282 ppm) in offsite surface water (HazDat 2002).

6.2.3 Soil

An estimated 97% of copper released from all sources into the environment is primarily released to land (Perwak et al. 1980). These include primarily tailings and overburdens from copper mines and tailings

6. POTENTIAL FOR HUMAN EXPOSURE

from mills. The copper in tailings represents the portion of copper that could not be recovered from the ore and is generally in the form of insoluble sulfides or silicates (Perwak et al. 1980). These wastes accumulate in mining states. Other releases to land include sludge from POTWs, municipal refuse, waste from electroplating, iron and steel producers, and discarded copper products (e.g., plumbing, wiring) that are not recycled. The copper content of municipal solid waste is ~0.16%. Much of this waste is landfilled directly or is in the form of residues following incineration. Emission factors in milligrams of copper released per gram of solid waste have been estimated for various industries. These factors would enable estimation of an industry's copper releases in terms of total quantity of solid waste discharged. Sludge from sewage treatment plants is a major source of copper released to land (Nriagu and Pacyna 1988). Agricultural products are believed to constitute 2% of the copper released to soil (Perwak et al. 1980). However, even though the largest releases of copper are to land, uptake of copper in human populations through ingestion of copper in soil are expected to be minimal in comparison to the primary route of exposure through the ingestion of drinking water (see Section 6.5).

Copper was detected in soil and sediment (e.g., lakes, streams, ponds, etc.) at 528 and 338 of the 906 NPL hazardous waste sites, respectively, where copper has been detected in environmental media (HazDat 2004). Copper was detected at concentrations ranging from 0.01 to 182,000 ppm (median concentration of 0.103 ppm) in offsite soils and 0.022–14,000 ppm (median concentration of 43 ppm) in offsite sediments (HazDat 2002).

6.3 ENVIRONMENTAL FATE

When considering the environmental fate of a metal, it is not always possible to clearly separate the processes related to the transport and partitioning of a metal, its compounds, and complexes from those related to transformation and degradation of these metal species. Because of analytical limitations, investigators do not often identify the form of a metal present in the environment. A change in the transport or partitioning of a metal may result from the transformation of the metal from one form to another. For example, complexation of a metal with small organic compounds may result in enhanced mobility, while formation of a less-soluble sulfide would decrease its mobility in water or soil. Adsorption may be the result of strong bonds being formed (transformation) as well as weak ones. Characterizing weak and strong adsorption is dependent on the analytical method that is used and care should be exercised when comparing results from different studies. Deposition and general adsorption of copper are discussed in Section 6.3.1. Speciation, compound formation, and oxidation-reduction are examined in Section 6.3.2.

6. POTENTIAL FOR HUMAN EXPOSURE

6.3.1 Transport and Partitioning**6.3.1.1 Ambient Air**

Copper is released to the atmosphere in the form of particulate matter or adsorbed to particulate matter. It is removed by gravitational settling (bulk deposition), dry deposition (inertial impaction characterized by a deposition velocity), in-cloud scavenging (attachment of particles by droplets within clouds), and washout (collision and capture of particles by falling raindrops below clouds) (Schroeder et al. 1987). The removal rate and distance traveled from the source will depend on a number of factors, including source characteristics, particle size, turbulence, and wind velocity.

Gravitational settling governs the removal of large particles with mass median aerodynamic (MMA) diameters of $>5 \mu\text{m}$, whereas smaller particles are removed by the other forms of dry and wet deposition. The importance of wet to dry deposition generally increases with decreasing particle size. The scavenging ratio (ratio of the copper concentration in precipitation [ppm] to its air concentration [$\mu\text{g}/\text{m}^3$]) for large particles displays a seasonal dependence that reflects more effective scavenging by snow than by rain (Chan et al. 1986). Copper from combustion sources is associated with sub-micron particles. These particles remain in the troposphere for an estimated 7–30 days. In that time, some copper may be carried far from its source (Perwak et al. 1980).

Rates of metal deposition (e.g., depositional fluxes) vary between dry and wet depositional processes and show spatial variability. Dry depositional fluxes of copper tend to decrease between highly urbanized area such as Chicago, Illinois with an average depositional rate of $0.06 \text{ mg}/\text{m}^2/\text{day}$, to less urbanized areas such as South Haven, Michigan with rate of $0.007 \text{ mg}/\text{m}^2/\text{day}$ or areas with minimal anthropogenic activity such as Lake Michigan (between 6 and 10 km off shore) with a rate of $0.01 \text{ mg}/\text{m}^2/\text{day}$ (Paode et al. 1998). Estimated copper deposition rates in urban areas are 0.119 and 0.164 kg per hectare per year ($\text{kg}/\text{ha}/\text{year}$) or 0.0326 and $0.0449 \text{ mg}/\text{m}^2/\text{day}$ for dry and wet deposition, respectively (Schroeder et al. 1987). Bulk deposition reportedly ranges from $0.002\text{--}3.01 \text{ kg}/\text{ha}/\text{year}$ or $0.0005\text{--}0.825 \text{ mg}/\text{m}^2/\text{day}$ (Golomb et al. 1997; Landing et al. 1995; Schroeder et al. 1987). For rural areas, the range of bulk deposition reportedly is $0.018\text{--}0.5 \text{ kg}/\text{ha}/\text{year}$ or $0.0049\text{--}0.14 \text{ mg}/\text{m}^2/\text{day}$, and wet deposition is $0.033 \text{ kg}/\text{ha}/\text{year}$ or $0.009 \text{ mg}/\text{m}^2/\text{day}$. The washout ratio is $114,000\text{--}612,000 (\mu\text{g}/\text{m}^3 \text{ rain})/(\mu\text{g}/\text{m}^3 \text{ air})$ or, expressed on a mass basis, $140\text{--}751 (\mu\text{g}/\text{kg} \text{ rain})/(\mu\text{g}/\text{kg} \text{ air})$. In southern Ontario, Canada, where the average concentration of copper in rain was 1.57 ppb during 1982, 1.36 mg of copper was deposited annually per square meter, or $13.6 \text{ kg}/\text{ha}$, as a result of wet deposition (Chan et al. 1986). For central and

6. POTENTIAL FOR HUMAN EXPOSURE

northern Ontario, the mean concentrations of copper in rain were 1.36 and 1.58 ppb, respectively, and the annual wet depositions averaged in both instances 1.13 mg/m² or 11.3 kg/ha.

For the majority of time, the concentration in air of toxic trace elements, like copper, in a study conducted by Sweet et al. (1993) approached levels measured at a rural site (Bondville, Illinois). These rural levels of airborne copper represented regional background levels in urban study sites with only episodic increases, depending on wind speed and direction and location relative to local point sources. In one urban study site (East St. Louis), smelters are the primary source of copper. Copper depositional fluxes follow an exponential decay as one transitions from urban to rural settings (Sweet et al. 1993). Soil is not the major source of copper in cities or nearby rural soils, but is the predominant source for copper in the atmosphere over more remote areas (Fergusson and Stewart 1992). Sources of copper in urban areas include coal combustion, soil, tire wear, and automobile emissions (Kim and Fergusson 1994). Copper emission from combustion processes is typically associated with fine particles; however, there can be instances where the highest concentrations of copper are measured in coarse particles obtained from paved and unpaved roads and industries (Paode et al. 1998).

Estimated depositional velocities for fine particles (<2.5 µm) and coarse particles (2.5–10 µm) in urban (Chicago) and rural (Kankalee, Illinois) areas have been made (Pirrone and Keeler 1993). These are: urban, 0.25–0.46 cm/second and rural, 0.18–0.25 in (rural) Kankalee, Illinois for fine particles; and urban, 1.47–2.93 cm/second and rural, 0.87–1.71 cm/second for coarse particles. The differences in velocities are due to higher surface roughness and wind velocities in Chicago.

Copper concentrations in particulates formed in a controlled study of waste oil combustion are (in µg/g): 687±11 (10 µm), 575±8 (50 µm), 552±12 (100 µm), 568±9 (300 µm), and 489±8 (500 µm).

Approximately 25% of copper is in the 10 µm fraction and ~18% is in each of the larger fractions (e.g., 50, 100, 300, and 500 µm) (Nerín et al. 1999).

6.3.1.2 Ambient Waters

The average concentrations of copper in Lakes Superior, Erie, and Ontario are 760, 870, and 830 ng/L, respectively (Georgopoulos et al. 2001; Nriagu et al. 1996). These values were derived from measurements taken from 11, 11, and 9 nearshore and offshore sampling sites at different points in the water column up to depths of 251, 55, and 145 meters for Lakes Superior, Erie, and Ontario, respectively (Nriagu et al. 1996). In Lake Ontario, the highest copper concentrations were found at nearshore

6. POTENTIAL FOR HUMAN EXPOSURE

sampling sites neighboring Buffalo, New York (887–1,051 ng/L), Rochester, New York (1,041–1,098 ng/L), and Kingston, Ontario (921–1,026 ng/L). The lowest concentrations of copper in Lake Ontario were measured in an offshore sampling site (540–710 ng/L) that was approximately 40 km from the Buffalo sampling site.

The atmospheric input of copper into the Great Lakes is substantial, 330–1,470 ng/m²/year, which amounts to a total deposition of 8.00–35.6x10¹³ ng/year. This input of copper accounts for 60–80% of the anthropogenic input into Lake Superior and 20–70% into Lakes Erie and Ontario (Georgopoulos et al. 2001; Nriagu et al. 1996). The mean residency times of copper in sediments are estimated to be 15 years in Lake Erie and 101 years in Lake Superior.

Much of the copper discharged into waterways is in particulate matter and settles out. In the water column and in sediments, copper adsorbs to organic matter, hydrous iron and manganese oxides, and clay. In the water column, a significant fraction of the copper is adsorbed within the first hour of introduction, and in most cases, equilibrium is obtained within 24 hours (Harrison and Bishop 1984). In fact, most of the copper in POTW effluent and surface runoff is already in the form of complexes (Sedlak et al. 1997). Copper in waste water discharged into Back River leading into Chesapeake Bay, Maryland, contained 53 ppb of copper, of which 36 ppb (based on weight) were in the form of settleable solids (Helz et al. 1975). The concentration of copper rapidly decreased downstream of the outfall so that 2–3 km from the outfall, the copper concentration had fallen to 7 ppb. The concentration of copper in sediment 2–3 km downstream from the outfall was about a factor of 10 higher than in uncontaminated areas (e.g., Rappahannock River). Based on their data and the results from other studies, Helz et al. (1975) estimated a total of 200 metric tons of copper entered the Chesapeake Bay from the effluent discharged from waste treatment plants.

Copper binds primarily to organic matter in estuarine sediment, unless the sediment is low in organic matter content. A study evaluated the importance of the absorption properties of different nonlithogenic components of aerobic estuarine sediment to copper binding by determining copper's adsorptivity to model components (phases) in artificial seawater (Davies-Colley et al. 1984). The phases included hydrous iron and manganese oxides, clay, aluminosilicates, and organic matter. The binding affinities varied by over a factor of 10,000 and were in the following order: hydrous manganese oxide > organic matter > hydrous iron oxide > aluminosilicates > clay (montmorillonite). The partition coefficients at pH 7 for the more strongly bound phases (manganese oxide, iron oxide, and estuarine humic material), were 6,300, 1,300, and 2,500, respectively. The affinity increased somewhat with pH; but did not vary

6. POTENTIAL FOR HUMAN EXPOSURE

appreciably when the salinity was reduced from 35 to 5%. Considering the compositional characteristics of estuarine sediment in terms of binding capacity, the results indicate that copper binds predominantly to organic matter (humic material) and iron oxides. Manganese oxide contributes only 1% to the binding because of its generally low concentration in sediment; the other phases are usually unimportant. These findings concur with results of selective extraction experiments (Badri and Aston 1983) and studies of the association of copper with humic material (Raspor et al. 1984).

6.3.1.3 Ambient Soils

Most copper deposited on soil from the atmosphere, agricultural use, and solid waste and sludge disposal will be adsorbed with greater concentrations of copper measured in the upper 5–10 centimeters of soil in comparison to lower soil depths, except in sandy soils where the lability of bound copper is greater (Breslin 1999; Giusquiani et al. 1992; Hutchinson 1979; Luncan-Bouché et al. 1997; Keller and Védý 1994; Levy et al. 1992; Perwak et al. 1980). Copper's movement in soil is determined by a host of physical and chemical interactions of copper with the soil components. In general, copper will adsorb to organic matter, carbonate minerals, clay minerals, or hydrous iron and manganese oxides (EPA 1979; Fuhrer 1986; Janssen et al. 1997; Petruzzelli 1997; Tyler and McBride 1982). Sandy soils with low pH have the greatest potential for leaching. In a laboratory study, Luncan-Bouché et al. (1997) have shown that between 55 and 85% of copper bound to sand (with no other soil components added) is remobilized upon reduction of the pH from 9 to 4. In most temperate soils, the pH, organic matter, concentrations of metal oxyhydroxides and ionic strength of the soil solutions are the key factors affecting adsorption (Elliot et al 1986; Fuhrer 1986; Gerritse and Van Driel 1984; Janssen et al. 1997; Rieuwerts et al. 1998; Tyler and McBride 1982). The ionic strength and pH of the soil solution affect the surface charge of soils and thereby influence ionic interaction (Rieuwerts et al. 1998). Soil microorganisms also affect the absorption of copper in soils due to the uptake and assimilation of the metal by these microorganisms (Rieuwerts et al. 1998). However, it is not known how the rate of uptake and absorption capacity of the microorganisms for copper compares with the binding capacity and affinities of copper by organic matter in soils, such as humic and fulvic acids. When the amount of organic matter is low, the mineral content or Fe, Mn, and Al oxides become important in determining the adsorption of copper. Fuhrer (1986) reported that, in oxidized estuarine sediment, adsorption of copper is dominated both by amorphous iron oxide and humic material.

Copper binds strongly to soils with high organic content (14–34% organic matter, dry weight) and the distribution of copper in the soil solution is less affected by changes in pH (within the range of pHs

6. POTENTIAL FOR HUMAN EXPOSURE

normally encountered in the environment) than other metals are (Gerritse and Van Driel 1984). In a laboratory study of competitive adsorption and leaching of metals in soil columns of widely different characteristics, copper eluted in a 0.01 M CaCl₂ leaching solution much more slowly and in much lower quantities than Zn, Cd, and Ni from a low-pH and a high-pH mineral soils and not at all from peat soil, which contained the greatest amount of organic matter (Tyler and McBride 1982). Elliot et al. (1986) investigated at pH-dependent adsorption of the divalent transition metal cations Cd, Cu, Pb, and Zn in two mineral soils (silty clay loam, 0.5 g/kg organic dry weight, and sandy clay, 1.6 g/kg organic) and two soils containing considerable organic matter (loamy sand, 20.5 g/kg organic, and silt loam, 42.5 g/kg organic). Adsorption increased with pH, and Cu and Pb were much more strongly retained than Cd and Zn. Reduction in absorptivity after removal of the organic matter demonstrated the importance of organic matter in binding copper. In a study of clay soils, Wu et al. (1999) observed preferential copper binding to organic matter, but found higher binding affinities to fine (<0.2 μm) clay fractions once the organic matter had been removed.

To determine the factors affecting copper solubility in soil, Hermann and Neumann-Mahlkau (1985) performed a study in the industrial Ruhr district of West Germany, which has a high groundwater table (10–80 cm from the surface) and a history of heavy metal pollution. Groundwater samples were taken from six locations and two soil horizons, an upper oxidizing loam, and a lower reducing loam. Total copper concentrations were high in the upper soil horizons and low in the lower horizons. Copper showed a pronounced solubility only in the oxidizing environment. In the reducing environment, solubility was low, possibly due to the formation of sulfides.

The form of copper at polluted and unpolluted sites may affect its leachability, particularly by acid rain. The leaching of heavy metals by simulated acid rain (pH 2.8–4.2) was measured by applying this rainwater to columns containing humus layers obtained from sites in a Swedish spruce forest both near to and far from a brass mill (Strain et al. 1984). Leaching of copper increased considerably when water with a pH <3.4 was applied to soil from polluted sites. Acid rain produced from SO_x emitted from smelters may increase the leachability of copper in areas affected by smelter stack emissions. The mobility of copper from soils was also found to increase following the introduction of 10–100 mM sodium chloride or calcium magnesium acetate deicing salts into soil (Amrhein et al. 1992). The concentration of sodium chloride or calcium magnesium acetate used in the study approximate those in runoff water produced from the melting of snow along salted roadways.

6. POTENTIAL FOR HUMAN EXPOSURE

Since, in the range of 0.01–1.97 mg of copper per liter of water, 25–75% of copper entering POTWs is removed in sludge, much of which is disposed of by spreading on land, it is important to ascertain whether copper in sludge is apt to leach into soil. Where it stands, this does not appear to be the case and leachate collected from the sludge-amended soil contained <12 ppb of copper (Perwak et al. 1980). In laboratory experiments, three sludges containing 51, 66, and 951 ppm (dry weight) of copper were applied to the top of soil columns containing four coastal plain soils. These soils, Sassafras loamy sand, Woodstown sandy loam, Evesboro loamy sand, and Matapeake silt loam, had similar pHs, 5.1–5.8, and contained 0.9, 1.4, 1.6, and 3.5% organic matter (dry weight), respectively. The sludge containing 51 ppm copper was loaded on the soil columns at amounts that approximated field loadings of between 0 and 90 metric tons per hectare; the sludges containing 66 and 951 ppm copper were loaded in amounts that approximated field loadings of between 0 and 180 metric tons per hectare. The columns were subsequently leached with distilled water at a flow rate of 2.5 cm/day for a total column application of 25.4 cm of water. Only small amounts (<0.01–0.87 ppm) of copper were found in the leachate (Ritter and Eastburn 1978). This suggests that hazardous amounts of copper leach only slowly into groundwater from sludge, even from sandy soils. In another study, soil cores taken after sewage sludge were applied to grassland for 4 years showed that 74 and 80% of copper remained in the top 5 cm of a sandy loam and calcareous loam soil (Davis et al. 1988). Similar studies have also shown that copper is typically confined to the upper 5–10 cm of sludge-amended agricultural soils (Breslin 1999; Giusquiani et al. 1992). In soils receiving long-term, heavy applications of sludge, high copper concentrations (471 mg/kg in comparison to 19.1 mg/kg in unamended control soils) were reported to depths of up to 25 cm (Richards et al. 1998). The mobility of copper into soil from sludge was found to be determined mainly by the amount of soil organic carbon and soil surface area (Domergue and Védy 1992; Gao et al. 1997). In addition, soils amended by sludge with low metal content were found to have increased sorption for copper due to the increased binding capacity provided by the “low metal” organics in the sludge (Petruzzelli et al. 1994).

Similarly, copper remains in the surface layer when it is applied to soil as a liquid. Secondary sewage effluent spiked with 0.83 ppm of copper was applied weekly to four different soils. After 1 year of treatment, the concentration of copper in the surface horizons increased greatly: 50–76% of the applied copper was found in the upper 2.5 cm and 91–138% was found in the upper 12.7 cm (Brown et al. 1983). In a study of accumulation and movement of metal in sludge-amended soils, field plots received massive amounts of sewage over a period of 6 years. Two sludges (one containing industrial waste), with average copper contents of 0.29 and 23 ppm were incorporated into the top 20 cm of soil in the spring. Barley was grown and, after harvest, core samples of soil were taken down to 1 m. Some movement of copper

6. POTENTIAL FOR HUMAN EXPOSURE

into the 22.5–25 cm layer of soil was observed, but little, if any, below this zone. However, at this depth, the copper is still within the root zone of many important food crops and, therefore, is available for uptake into these plants. Also, the availability of the copper in soil, as determined by its extractability with diethylenetriamine pentaacetic acid (DTPA) and nitrate, remained constant over a 4-year period at all depths. From the results of other work, the major portion of the copper (40–74%) is expected to be associated with the organic, Fe-Mn-oxide and carbonate fractions of most soils (Ma and Rao 1997).

6.3.1.4 Bioconcentration and Biomagnification

The bioconcentration factor (BCF) of copper in fish obtained in field studies is 10–667, indicating a low potential for bioconcentration (Perwak et al. 1980). The BCF is higher in mollusks, such as oysters, and squid where it may reach 30,000 and 2.1×10^7 , respectively, (Perwak et al. 1980) and may present a major dietary source of copper that could be of concern for those individuals who regularly consume oysters, clams, or squid. Due to the fact that molluscs are filter feeders and copper concentrations are higher in particulates than in water, this is to be expected. On the other hand, there are limited data suggesting that there is little biomagnification of copper in the aquatic food chain (Perwak et al. 1980). For example, a study was conducted with white suckers and bullheads, both bottom-feeding fish, in two acidic Adirondack, New York, lakes (Heit and Klusek 1985). These lakes were known to have received elevated loadings of copper; but the suckers and bullhead had average copper levels of only 0.85 and 1.2 ppm (dry weight) in their muscle tissue. The biomagnification ratio (the concentration of copper in fish compared to that in their potential food sources on a wet weight/wet weight basis) was <1 , indicating no biomagnification in the food chain. Similarly, the copper content of muscle tissue of fish from copper-contaminated lakes near Sudbury, Ontario, did not differ significantly from that of the same fish species in lakes far from this source (Bradley and Morris 1986).

No evidence of bioaccumulation was obtained from a study of pollutant concentrations in the muscle and livers of 10 mammal species in Donana National Park in Spain (Hernandez et al. 1985). The park is impacted by organochlorine compounds and heavy metals emitted from anthropogenic activities that surround the park. For example, the Guadalquivir River that flows through the park first flows through a major mining region in addition to a large urban area and industrial areas, carrying with it contaminants acquired from these sites. The animal species in the study were classified into three categories (herbivorous, omnivorous, and carnivorous) to ascertain if the pollutants were showing biomagnification in higher trophic levels of animals. No evidence of copper biomagnification in the food chain was observed. Likewise, in a study of a food web in a beech tree forest in Northern Germany, there was no

6. POTENTIAL FOR HUMAN EXPOSURE

evidence of biomagnification in tertiary consumers (e.g., vole, shrew, and mouse) compared to secondary consumers (e.g., earthworm, snail, beetle, and isopod) (Scharenberg and Ebeling 1996). A study of heavy metals in cottontail rabbits on mined land treated with sewage sludge showed that, while the concentration of copper in surface soil was 130% higher than in a control area, the elevation was relatively little in foliar samples and no significant increase in copper was observed in rabbit muscle, femur, kidney, or liver. Apparently, copper was not bioaccumulating in the food chain of the rabbit (Dressler et al. 1986).

At the lowest levels of the food chain, there is little evidence of copper bioaccumulation. In a study of copper uptake in earthworms as a function of copper concentration (6–320 mg/kg dry weight) in sludge amended soils, a bioconcentration factor of <1 (0.67) was obtained (Neuhauser et al. 1995). In another example, a study of earthworms and soil from 20 diverse sites in Maryland, Pennsylvania, and Virginia, copper concentrations in earthworms showed a poor correlation with that in soil (Beyer and Cromartie 1987). These results are consistent with the results of another study that also showed no clear correlation between copper concentrations in earthworm tissues and two soils that were heavily contaminated with heavy metals (copper concentrations of 242 and 815 mg/kg dry weight) (Marinussen et al. 1997).

However, there is some evidence in one study for bioconcentration of copper at low copper concentrations in soil. Even though Scharenberg and Ebeling (1996) showed that there was no evidence for biomagnification of copper in a forest food web, their results did show that the total concentrations of copper in the secondary (18.3–192.0 mg/kg dry weight) and tertiary consumers (9.9–17.4 mg/kg dry weight) were higher than the concentrations of the metal in the dominant vegetation (5.3–10.9 mg/kg dry weight) and soil (1.8–5.8 mg/kg dry weight) in the ecosystem.

Diks and Allen (1983) added copper to four sediment/water systems and studied the distribution of copper among five geochemical phases, namely, absorbed/exchangeable, carbonate, easily reducible (Mn-oxides and amorphous Fe-oxides), organic, and moderately reducible (hydrous Fe-oxides). The investigators then attempted to correlate the concentration in each phase with the copper uptake by tubificid worms. Only copper extracted from the manganese oxide/easily-reducible phase correlated with the copper content of worms at the 95% confidence level. This result suggests that the redox potential and pH in the gut of the worm is such that manganese oxide coatings are dissolved. The copper in the dissolved manganese oxide phase is now soluble and available for uptake by other organisms.

6. POTENTIAL FOR HUMAN EXPOSURE

6.3.2 Transformation and Degradation**6.3.2.1 Air**

Few data are available regarding the chemical forms of copper in the atmosphere and their transformations. In the absence of specific information, it is generally assumed that metals of anthropogenic origin, especially those from combustion sources, exist in the atmosphere as oxides because metallic species are readily attacked by atmospheric oxidants. As these oxides age, sulfatization may occur, but only when SO_x gases are present in the atmosphere in sufficient amount. For example, in Arizona, atmospheric copper oxide levels derived from copper smelters was strongly correlated with co-emitted sulfur (Schroeder et al. 1987).

In fog water, Cu(II) is reduced to Cu(I) by sulfite, which becomes enhanced by the fact that sulfite is also a ligand of and binds to Cu(I) (Xue et al. 1991). Concentrations of Cu(I) in fog water ranged between 0.1 and 1 μM or, respectively, 4 and >90% of copper in the Cu(I) state. The reduction of Cu(II) to Cu(I) is pH dependent and occurs rapidly at $\text{pH} > 6$ (Xue et al. 1991).

6.3.2.2 Water

The Cu(I) ion is unstable in aqueous solution, tending to disproportionate to Cu(II) and copper metal unless a stabilizing ligand is present (EPA 1979; Kust 1978). The only cuprous compounds stable in water are insoluble ones such as Cu_2S , CuCN , and CuF . Therefore, human exposures to copper will predominately be in the form of Cu(II). Copper in its Cu(II) state forms coordination compounds or complexes with both inorganic and organic ligands. Ammonia and chloride ions are examples of species that form stable ligands with copper. Copper also forms stable complexes with organic ligands such as humic acids, binding to $-\text{NH}_2$ and $-\text{SH}$ groups and, to a lesser extent, with $-\text{OH}$ groups. Natural waters contain varying amounts of inorganic and organic species. This affects the complexing and binding capacity of the water and the types of complexes formed. In seawater, organic matter is generally the most important complexing agent (Coale and Bruland 1988). In water, the formation of ligands may affect other physicochemical processes such as adsorption, precipitation, and oxidation-reduction (EPA 1979). More specific information on the transformation and degradation of copper in its cupric [Cu(II)] and cuprous [Cu(I)] states is given below.

At the pH values and carbonate concentrations characteristic of fresh surface waters, most dissolved Cu(II) exists as carbonate complexes rather than as free (hydrated) cupric ions (Stiff 1971).

6. POTENTIAL FOR HUMAN EXPOSURE

Based on the results of a theoretical model, the major species of soluble copper found in freshwater, seawater, and a 50:50 combination of the freshwater and seawater over a pH range of 6.5–7.5 is Cu^{2+} , $\text{Cu}(\text{HCO}_3)^+$, and $\text{Cu}(\text{OH})_2$ (Long and Angino 1977).

The concentration of dissolved copper depends on factors such as pH, the oxidation-reduction potential of the water and the presence of competing cations (Ca^{2+} , Fe^{2+} , Mg^{2+} , etc.), salts (OH^- , S^{2-} , PO_4^{3-} , CO_3^{2-}), and anions of insoluble cupric-organic and -inorganic complexing agents. If the combination of a particular anion with copper forms an insoluble salt, precipitation of that salt will occur. The most significant precipitate formed in fresh surface waters is malachite ($\text{Cu}_2[\text{OH}]_2\text{CO}_3$) (Sylva 1976). Other important precipitates are $\text{Cu}(\text{OH})_2$ (and ultimately CuO) and azurite ($\text{Cu}_3[\text{OH}]_2[\text{CO}_3]_2$). In anaerobic waters, Cu_2S , Cu_2O , and metallic copper forms and settles out (EPA 1979). The combined processes of complexation, adsorption, and precipitation control the level of free $\text{Cu}(\text{II})$ in water. The chemical conditions in most natural water are such that, even at relatively high copper concentrations, these processes will reduce the free $\text{Cu}(\text{II})$ concentration to extremely low values.

As a result of the aforementioned physico-chemical processes, copper in water may be dissolved or associated with colloidal or particulate matter. Copper in particulate form includes precipitates, insoluble organic complexes, and copper adsorbed to clay and other mineral solids. In a survey of nine rivers in the United Kingdom, 43–88% of the copper was in the particulate fraction (Stiff 1971). A study using suspended solids from the Flint River in Michigan found that the fraction of adsorbed copper increased sharply with pH, reaching a maximum at a pH of 5.5–7.5 (McIlroy et al. 1986).

The soluble fraction of copper in water is usually defined as that which will pass through a 0.45 μm filter. It includes free copper and soluble complexes as well as fine particulates and colloids. The soluble fraction may be divided according to the lability (e.g., the relative ability of the copper to dissociate from the bound form to the free ion) of the copper forms in the water. Categories range from the very labile (e.g., free metal ion, ion pairs, inorganic or organic complexes) to slowly or nonlabile (e.g., colloiddally bound to inorganic colloidal phases of other metals such as $\text{Fe}(\text{OH})_3$ or FeOOH , or bound to high molecular weight organic material) metal (Tan et al. 1988). For example, in a typical study, 18–70% of dissolved copper in river water was labile and 13–30% was slowly labile (Tan et al. 1988). Various techniques may be used to classify the lability of different fractions of soluble copper; these techniques include solvent extraction, ion-specific electrodes, ion exchange, ultrafiltration, electrochemical methods such as anodic stripping voltammetry, and gel filtration chromatography (Harrison and Bishop 1984).

6. POTENTIAL FOR HUMAN EXPOSURE

The resulting classification depends on the specific procedure employed. Therefore, a comparison of the results of different researchers should be done in general terms

The nature of copper's association with inorganic and organic ligands will vary depending on the pH, copper concentration, concentration of competing ligands, binding capacity of the ligands, and hardness or salinity of the water (Breault et al. 1996; Cao et al. 1995; Gardner and Ravenscroft 1991; Giusti et al. 1993; Lores and Pennock 1998; Town and Filella 2000). In river water from the northwestern United States that had a relatively high pH (7.0–8.5) and alkalinity (24–219 ppm as CaCO_3), inorganic species like CO_3^{2-} and OH^- were the most important ligands at high copper concentrations (McCrary and Chapman 1979). However, other species such as organic compounds were important at low copper concentrations. On the other hand, copper in samples from surface water of lakes and rivers in southern Maine with a relatively low pH (4.6–6.3) and alkalinity (1–30 ppm as CaCO_3) was largely associated with organic matter (Giesy et al. 1978). The binding of copper to dissolved organics was found to be dependent on the specific organic chemical species (e.g., fulvic acid) and their concentrations in the surface water, the number of available binding sites per fulvic acid carbon, and the hardness of the water (Breault et al. 1996). Increasing water hardness results in decreased fulvic acid binding sites. This effect is due more to the depression of the solubility of high molecular weight fulvic acid in the presence of Ca and Mg ions than to competition of these ions with copper for fulvic acid binding sites. Increasing pH from 8 to 6 resulted in a 7-fold increase in the binding constant for Cu(II) with humic acid (Cao et al. 1995).

The extent to which copper binds to inorganic and organic ligands can be altered by materials carried in runoff. For example, after a period of rain in southeastern New Hampshire, inorganic constituents contributed more to copper binding in lakes and rivers than did dissolved organic matter (Truitt and Weber 1981). A green precipitate, confirmed to be malachite ($\text{Cu}_2[\text{OH}]_2\text{CO}_3$) was formed in river water in Exeter. This water had the highest pH (7.4) and alkalinity (43.5 mg/L as CaCO_3) than six other surface waters (e.g., three rivers, two reservoirs, a pond, and a swamp) with pH and alkalinity values of 5.7–7.4 and 1.7–41 mg/L, respectively. A computer simulation of the copper species in water of a pond and water obtained from an artesian well that fed the pond predicted that 98% of the copper in the artesian well water would exist as the free copper ion (Cu^{+2}), whereas 88 and 63% of the copper in pond water would be bound to organics in the spring and fall, respectively (Giesy et al. 1983). These estimates were based on experimentally determined binding capacities of the organic matter in the two water sources and stability constants for the copper-organic matter complexes.

6. POTENTIAL FOR HUMAN EXPOSURE

Seawater samples obtained in a transect of the uppermost Narragansett Bay in August 1980 were analyzed for dissolved, particulate and organically bound copper to investigate the geochemistry of copper-organic complexes (Mills and Quinn 1984). Narragansett Bay is a partly mixed estuary in Massachusetts and Rhode Island that receives organic matter and metals from rivers, municipal and industrial effluents and from runoff. The Fields Point waste treatment facility accounts for 90% of the copper input into the bay through the Providence River with dissolved copper representing 60% of the total copper input. The concentrations of dissolved and organic copper ranged from 16.4 and 2.3 $\mu\text{g}/\text{kg}$, respectively, in the Providence River to 0.23 and 0.12 $\mu\text{g}/\text{kg}$, respectively, in Rhode Island Sound. Particulate copper concentrations in Narragansett bay ranged from 2.42 to 0.06 $\mu\text{g}/\text{kg}$ and generally comprised 40% of the total copper in the bay. Analysis of the data indicated that ~75% of the dissolved copper that enters the bay from the Providence River is removed within the bay.

Organic ligands may contain a variety of binding sites, and the strength of the resulting copper complexes will vary accordingly. Over 99.7% of the total dissolved copper in ocean surface water from the northeast Pacific was associated with organic ligands (Coale and Bruland 1988). The dominant organic complex, limited to surface water, was a strong ligand of biological origin. A second, weaker class of organic ligand was of geologic origin. An independent study showed the copper binds to humic material at a number of sites. The binding strength of the sites varied by two orders of magnitude (Giesy et al. 1986). The humic material in this study was derived from nine surface waters in the southeastern United States. Soluble copper in water discharged from a nuclear power station was primarily complexed with organic matter in the 1,000–100,000 molecular weight range (Harrison et al. 1980). Ten to 75% of the discharged copper was in particulate form.

The bioavailability of Cu(I) has been difficult to access due to its thermodynamic instability in the environment (Xue et al. 1991). Cu(I) is a reactive reducing agent, and its concentrations in the environment will be determined both by its reaction with oxygen and other oxidants in the aqueous environment to form Cu(II) and its rate of production through the reaction of Cu(II) with reducing agents (Sharma and Millero 1988). Investigators have shown the presence of Cu(I) in seawater, which is thought to occur through the reduction of Cu(II) to Cu(I) by photochemical processes (Moffett and Zika 1987; Xue et al. 1991). The detection of Cu(I) in seawater is likely the result of the stabilization of Cu(I) through complex formation with chloride ions. Cu(II)-organic complexes absorb radiation at wavelengths >290 nm and can undergo charge transfer reactions where the Cu(II) is reduced and a ligand is oxidized. Photochemically-generated reducing agents such as O_2^- and H_2O_2 in the surface water of oceans and

6. POTENTIAL FOR HUMAN EXPOSURE

possibly other natural waters (e.g., lakes) may contribute to the reduction of Cu(II) to Cu(I) in these waters (Moffett and Zika 1987; Sharma and Millero 1988).

Experiments performed in synthetic seawater and water from Biscayne Bay, Florida, showed that in the reduction of Cu(II) to Cu(I), the rate was first-order in Cl^- and second-order in H_2O_2 (Sharma and Zika 1987). The chloride ion is thought to be required for stabilizing Cu(I) by forming the copper complex CuClOH^- , although experimental evidence suggests that the reduction of Cu(II) may also occur through the formation of a complex between Cu(II) and HO_2^- . Experiments have shown that as much as 15% of copper in surface seawater exists as Cu(I). Photochemical reduction by sunlight increases the percentage of free Cu(I). The photochemical reduction mechanism is supported by the observation that the Cu(I) concentration is highest in the surface layer of seawater and that the hydrogen peroxide concentration increases in parallel to that of Cu(I) (Moffett and Zika 1987). In addition, the percentage of free Cu(I) is highest on the surface.

Once Cu(I) is formed, its lifetime is determined by its rate of oxidation to Cu(II). After Biscayne Bay water was exposed to sunlight for 5 hours, the Cu(I) formed was oxidized to Cu(II). The half-life of Cu(I) was 12 hours. Primarily, dissolved oxygen is responsible for this oxidative reaction. Since the oxidation of Cu(I) by O_2 in distilled water occurs in <6 minutes, Cu(I) in seawater apparently is stabilized by the formation of complexes. In the presence of humic acids, the oxidation of Cu(I) occurs very rapidly. In coastal water off the Everglades in Florida, no Cu(I) was detected. This is due to the binding of Cu(II) in organic complexes and the high concentration of radical oxidants in the water. Sharma and Millero (1988) measured the rate of Cu(I) oxidation in seawater as a function of pH, temperature and salinity. The rate of reaction increased with pH and temperature, and decreased with increasing ionic strength (or salinity). The results suggested that the rates are controlled by Mg^{2+} , Ca^{2+} , Cl^- and HCO_3^- through their involvement in complex formation and ligand exchange.

6.3.2.3 Sediment and Soil

The adsorption of copper to soil and sediment was discussed in Section 6.3.1 under transport and partitioning, even though adsorption may really be complexation and transformation. Understanding the transport and fate of copper and its compounds in soils and sediments is important because these compartments tend to be large reservoirs of copper and could have an impact on human exposures to copper. Copper concentrations in drinking water obtained from groundwater can be affected by the leaching of copper from soil. Reservoir sediments have been shown to be sources of copper in drinking

6. POTENTIAL FOR HUMAN EXPOSURE

water (Georgopoulos et al. 2001). Although much of the copper is bound to inorganic or organic matrices in soils and sediments, there is the potential for release of copper into pore water within soils and sediments depending on the extractability of the copper and soil conditions. There is evidence to suggest that copper binding in soil is correlated with pH, cation exchange capacity, the organic content of the soil, the presence of manganese and iron oxides and even the presence of inorganic carbon such as carbonates (Petruzzelli 1997; Rieuwerts et al. 1998). At pHs above 5, absorption of copper from pore water on to soil components becomes a significant process, whereas at pHs below 5, copper largely remains in pore water and is therefore mobile in soil (Perwak et al. 1980). However, broad generalizations about the lability of copper in soils are not possible since the situation will differ among different soil types and environmental conditions. More specific information on the lability (e.g., extractability) of copper from differing soils and conditions is given below.

The form of copper in soil is determined by measuring the extractability of the copper with different solvents. Extractability is a function of the nature of the soil and the form of copper deposited in the soil. If a relatively labile form of copper is applied, binding to inorganic and organic ligands may occur, as well as other transformations. On the other hand, if a mineral form is deposited, it would be unavailable for binding. The capacity of soil to remove copper and the nature of the bound copper were evaluated by incubating 70 ppm of copper with 5 g samples of soil for 6 days (King 1988). Twenty-one samples of soils (10 mineral and 3 organic) from the southeastern United States were included in the study. Some soil samples were taken from the subsoil as well as the surface. The amount of adsorbed copper ranged from 36 to 100%, of which 13–100% was nonexchangeable when extracted with KCl. Removal of copper from solution was much higher with surface soils than with subsurface sandy soils; 95–100% of the copper was removed by five of the mineral surface soils and all three organic soils. The percentage of copper that was nonexchangeable was relatively high in all but some of the acid subsoils. While the fraction of exchangeable copper was not dependent on pH in surface soils, 96% of the variation in exchangeability was correlated with pH in subsoils. The soil/water partition coefficient for copper was >64 for mineral soils and >273 for organic soils. Of the eight heavy metals in the study, only Pb and Sb had higher partition coefficients than copper. Most of the copper in Columbia River estuary sediment and soil was correlated with inorganic carbon (e.g., carbonate), but not with the amount of extractable Fe or the organic carbon content of the sediment (Fuhrer 1986).

The amount of ammonium acetate- and DTPA-extractable copper in wetland soil/sediment resulting from atmospheric deposition from smelters in Sudbury, Ontario, showed the same pattern as total copper, despite random variations in soil pH, redox potential and organic carbon (Taylor and Crowder 1983).

6. POTENTIAL FOR HUMAN EXPOSURE

Therefore, in this case, soil characteristics were not the dominant factors determining extractability and availability, but rather the form of copper that was deposited. The median concentrations of total copper, ammonium acetate-extractable copper, and DTPA-extractable copper at 25 sample sites were 371, 49, and 98 ppm, respectively.

In another study of copper partitioning in nine different contaminated soils, sequential extractions were used to operationally define six soil fractions in decreasing order of copper availability: water soluble > exchangeable > carbonate > Fe-Mn oxide > organic > residual (Ma and Rao 1997). The results of this study showed that the distribution of copper in these six soil fractions differed depending on the total copper concentration in the soil. As the copper concentration increased above 240 mg/kg, between 69 and 74.4% of the total copper was found in the water soluble, carbonate, Fe-Mn oxide, and organic fractions. In relatively uncontaminated soils (<240 mg/kg copper), between 97.6 and 99.6% of the copper was found to be associated with the residual fraction.

Within the estuarine environment, anaerobic sediments are known to be the main reservoir of trace metals. Under anaerobic conditions, cupric salts will reduce to cuprous salts. The precipitation of cupric sulfide and the formation of copper bisulfide and/or polysulfide complexes determine copper's behavior in these sediments (Davies-Colley et al. 1985). In the more common case where the free sulfide concentration is low due to the controlling coexistence of iron oxide and sulfide, anaerobic sediment acts as a sink for copper, that is, the copper is removed from water and held in the sediment as an insoluble cuprous sulfide. However, in the unusual situation where the free sulfide concentration is high, soluble cuprous sulfide complexes may form, and the copper concentration in sediment pore water may then be high.

In sediment, copper is generally associated with mineral matter or tightly bound to organic material (Kennish 1998). As is common when a metal is associated with organic matter, copper generally is associated with fine, as opposed to coarse, sediment. Badri and Aston (1984) studied the association of heavy metals in three estuarine sediments with different geochemical phases. The phases were identified by their extractability with different chemicals and termed easily or freely leachable and exchangeable; oxidizable-organic (bound to organic matter); acid-reducible (Mn and Fe oxides and possibly carbonates); and resistant (lithogenic). In the three sediments, the nonlithogenic fraction accounted for ~14–18% of the total copper and the easily exchangeable component was 5% of the total copper. In addition, the compositional associations of copper in sediment samples taken from western Lake Ontario were analyzed employing a series of sequential extractions (Poulton et al. 1988). The mean and standard

6. POTENTIAL FOR HUMAN EXPOSURE

deviation percentages of copper in the various fractions were: exchangeable, 0 (0); carbonate, 0.1 (0.3); iron or manganese oxide-bound, 0.2 (0.3); organic-bound, 40 (11); and residual, 60 (8). Another study found that 10–20% of the copper in Lake Ontario sediment samples was bound to humic acids, with virtually all of the copper bound to organic matter (Nriagu and Coker 1980). The concentration of copper associated with humic acids was 21–40 times greater than in the sediment as a whole.

6.3.2.4 Other Media

Copper is an essential nutrient in plant metabolism. Therefore, uptake of copper from soil in plants through the roots is a natural and necessary process, a process that is actively regulated by the plant (Clemens 2001). The uptake of copper into plants is dependent on the concentration and bioavailability of copper in soils. The bioavailability of copper is determined largely by the equilibrium between copper bound to soil components and copper in soil solution. As noted in the discussion of copper binding in soils (Section 6.3.1.3), this is determined by copper concentrations in soil, soil type, soil components, pH, oxidation-reduction potential in the soil, and concentrations of other cations and anions in the soil, etc. (Rieuwerts et al. 1998). Other factors include root surface area, plant genotype, stage of plant growth, weather conditions, interaction with other nutrients in the soil and the water table (Gupta 1979). Liming is another factor that affects copper uptake. For example, liming acidic soils has been shown to increase copper uptake in hay, but to decrease copper uptake in wheat (Gupta 1979). However, the effect that liming has on increasing soil pH does not appear to be the overriding mechanism behind the changes in copper uptake in plants, even though there is evidence that the addition of lime to soil to increase the pH to 7 or 8 reduces copper availability to some plants (Perwak 1980). This is evidenced by the fact that changes in pH (5.4–8.0) have been found to have little effect on copper concentrations in plant tissues (Gupta 1979).

6.4 LEVELS MONITORED OR ESTIMATED IN THE ENVIRONMENT

6.4.1 Air

Human exposure to copper in air comes from both natural and anthropogenic sources. For the general population, exposures to copper concentrations in air average between 5 and 200 ng/m³. The concentrations of copper in air can be higher in the proximity of major sources such as smelters, mining operations, and combustion sources (e.g., power plants, incinerators, automobiles, etc.). The results of

6. POTENTIAL FOR HUMAN EXPOSURE

several studies in which concentrations of copper in air were reported and described below and are summarized in Table 6-8.

According to the EPA's National Air Surveillance Network report for the years 1977, 1978, and 1979, median copper concentrations were 133, 138, and 96 ng/m³, respectively, for urban samples and 120, 179, and 76 ng/m³ for nonurban samples, respectively (Evans et al. 1984). In this study, 10,769 urban and 1,402 nonurban air samples collected for 24 hours were analyzed. For 1977, 1978, and 1979, 1% of urban samples exceeded 1,156, 975, and 843 ng/m³, respectively, and 1% of nonurban samples exceeded 1,065, 1,396, and 645 ng/m³, respectively. The maximum urban and nonurban copper concentrations reported were 4,625 and 4,003 ng/m³, respectively. Davies and Bennett (1985) reported average atmospheric copper concentrations of 5–50 ng/m³ in rural areas and 20–200 ng/m³ in urban locations. The concentrations in rural areas are considerably lower than those reported in the EPA survey. Data from many urban locations in the United States show concentrations of copper associated with particulate matter ranging from 3 to 5,140 ng/m³ (Schroeder et al. 1987). Remote and rural areas have concentrations of 0.029–12 and 3–280 ng/m³, respectively. The levels reported by Schroeder et al. (1987) are consistent with those obtained in a study of airborne trace elements in national parks (Davidson et al. 1985). In the Smokey Mountain National Park, the copper concentration in air was 1.6 ng/m³, while in the Olympic National Park, where several locations were monitored, 3.3–6.7 ng/m³ of copper was measured in the atmosphere. The lower copper concentrations found in Smokey Mountain Park compared with those in the Olympic National Park have been attributed to greater vegetative cover and higher moisture in the former and larger amounts of exposed rock and soil in the latter. Average copper crustal enrichment factors (the concentration of copper in air compared with the average concentration in the earth's crust) were 31 and 76, respectively.

As part of the Airborne Toxic Element and Organic Substances (ATEOS) project for determining patterns of toxic elements in different settings, three urban areas (Camden, Elizabeth, and Newark) and one rural site (Ringwood) in New Jersey were studied during two summers and winters between 1981 and 1983 (Lioy et al. 1987). Each site was sampled every 24 hours for 39 consecutive days. As an example, the geometric mean copper concentrations in the summer of 1983 were 16.0, 21.0, 21.0, and 6.0 ng/m³ for Camden, Elizabeth, Newark, and Ringwood, respectively. In the winter of 1983, the mean copper concentrations were slightly higher with values of 21.0, 36.0, 33.0, and 63.0 ng/m³, respectively. The levels of copper measured in these industrial urban areas are considerably higher than the mean values reported in the National Air Surveillance survey where arithmetic means of 0.201 and 0.259 ng/m³ for copper in air were obtained in 1978 and 1979, respectively (Evans et al. 1984). Summer and winter

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-8. Concentrations of Copper in Air

Date/sample	Location	Concentration ^a (ng/m ³) (mean) [median]	Comments	Reference
1977, urban	United States	[133], 433 ₉₀ , 1,156 ₉₉ (207.5), 3,296 _{max}	4,648 samples, National Survey	EPA 1984
1978, urban	United States	[138], 430 ₉₀ , 975 ₉₉ (200.8), 4,625 _{max}	3,615 samples, National Survey	EPA 1984
1979, urban	United States	[96], 363 ₉₀ , 519 ₉₉ (259.3), 1,627 _{max}	2,507 samples, National Survey	EPA 1984
1977, nonurban	United States	[120], 450 ₉₀ , 1,065 ₉₉ (193.2), 16,706 _{max}	709 samples, National Survey	EPA 1984
1978, nonurban	United States	[179], 607 ₉₀ , 1,396 ₉₉ (265.7), 1,396 _{max}	458 samples, National Survey	EPA 1984
1977, nonurban	United States	[76], 322 ₉₀ , 645 ₉₉ (141.7), 4,003 _{max}	235 samples, National Survey	EPA 1984
Urban		20–200, [50]	Representative values	Davies and Bennett 1985
Rural		5–50, [20]		
Remote		0.29–12	Values from literature survey	Schroeder et al. 1987
Rural		3–280		
Urban	Canada	17–500		
Urban	United States	3–5,140		
Urban	Europe	13–2,760		
Urban	Other	2.0–6,810		
1979, remote	Smokey Mountain National Park	(1.6)	Above canopy, crustal enrichment factor 31	Davidson et al. 1985
1980, remote	Olympic National Park	3.3–6.7, (5.6)	Crustal enrichment factor 76	Davidson et al. 1985
1981, 1982, summer	Camden, New Jersey	16.0–18.0 ^b , 100.0 _{max}	Seasonal variations noted; three urban areas and one rural area.	Lioy et al. 1987
	Elizabeth, New Jersey	21.0–29.0, 120.0 _{max}		
	Newark, New Jersey	25.0–33.0, 131.0 _{max}		
	Ringwood, New Jersey	13.0–63.0, 77.0 _{max}		
1982, 1983, winter	Camden, New Jersey	17.0–21.0, 231.0 _{max}		
	Elizabeth, New Jersey	28.0–36.0, 493.0 _{max}		
	Newark, New Jersey	21.0–27.0, 380.0 _{max}		
	Ringwood, New Jersey	6.0–18.0, 29.0 _{max}		

^aPercentile level and maximum indicated as subscripts.

^bConcentrations in Lioy et al. (1987) are geometric means, unless otherwise noted.

6. POTENTIAL FOR HUMAN EXPOSURE

maxima in the four ATEOS study areas were: 100.0 and 131.0 ng/m³ in Camden, 231.0 and 493.0 ng/m³ in Elizabeth, 131.0 and 380.0 ng/m³ in Newark, and 77.0 and 29.0 ng/m³ in Ringwood, respectively. Copper follows the same pattern as other heavy metals, in that increased copper levels are present in winter in urban areas and in summer in rural areas. No explanation for this pattern has been offered.

Anderson et al. (1988) performed a study of the atmospheric aerosols collected at a site in Chandler, Arizona, over a 12-day period in February and March 1982. Several major copper smelters are located ~120 km to the southeast, which were upwind of the sampling site during approximately 50% of the study period. Particles containing >0.5% Cu were termed 'Cu-bearing' particles; 5.6% of the fine (0.4 to ~2 µm) particles collected were in this category. The most abundant type of Cu-bearing particle, representing 74% of the total, was associated with sulfur. However, the analysis was not able to specify the form of sulfur present. These particles were often associated with Zn, Fe, Pb, As, and Ca. Sixteen percent of the Cu-bearing particles were associated with silicon and 4% were associated with chloride. The concentration of Cu-S particles was highest when the surface and upper level winds were from the southeast to the east, and reached a maximum 1–2 days after the winds began to blow. Therefore, the smelters to the southeast appear to be the probable source. The particles associated with silicon and chlorine did not show any apparent correlation with wind and were either from a diffuse regional source or a local source.

Mine waste dump sites are a source of airborne copper carried in dust (Table 6-9). Particle size distribution and the concentration of copper in particle size ranges differ depending on the mine waste site (Mullins and Norman 1994). For example, the mean concentrations (ppm, w/w) of copper in dust (<10 µm particle size range) collected at four mine waste dump sites in Butte, Montana, were 3,370 (Gray Rock), 1,950 (Corra), 1,960 (Late Acquisition), and 2,570 (Railroad Bed).

Mean concentration ranges of copper in remote (any area of lowest copper concentration such as the Antarctic or Arctic) and rural (any site that represents a regional background that is not directly influenced by local anthropogenic emissions) precipitation ranges were 0.013–1.83 and 0.68–1.5 ppb, respectively, based on a weight per unit volume basis (Barrie et al. 1987). Although an earlier survey referred to by these investigators (Galloway et al. 1982) yielded much higher values, 0.060 and 5.4 ppb, these were ascribed to sample contamination. The mean concentration of copper in rain reported in an extensive study in southern Ontario, Canada, was 1.57 (0.36 standard deviation) ppb during 1982 (Chan et al. 1986). These concentrations showed little spatial variability. Concentration of copper

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-9. Particle Size Distributions and Total Copper Concentrations in Dust Collected at Four Mine Waste Pump Sites in Butte, Montana

Site	Particle size (μm)	Percent in total dust collected	Concentration of copper (ppm, w/w)
Corra	4.7–10	76.6 \pm 4.8	1,550
	1.1–4.7	20.9 \pm 0.63	3,110
	<1.1	1.9 \pm 0.14	4,900
Gray rock	4.7–1.0	84.5 \pm 0.93	3,240
	1.1–4.7	13.6 \pm 0.82	4,120
	<1.1	1.9 \pm 0.14	4,370
Railroad bed	4.7–10	61.5 \pm 1.06	2,580
	1.1–4.7	31.3 \pm 0.96	2,850
	<1.1	7.2 \pm 0.26	1,400
Late acquisition	4.7–10	70.3 \pm 1.36	1,560
	1.1–4.7	25.0 \pm 1.18	2,730
	<1.1	4.7 \pm 0.44	3,330

Source: Mullins and Norman 1994

6. POTENTIAL FOR HUMAN EXPOSURE

in cloud water over Olympic Peninsula in Washington State has been measured at $1.7 \pm 1.6 \mu\text{g/L}$ (air-equivalent mean concentration of 0.5 ng/m^3) (Vong et al. 1997).

The concentration of copper in rain samples taken within 2–15 km downwind of the Claremont, New Hampshire, municipal waste incinerator was found to range from 0.11 to $2.12 \mu\text{g/L}$ with a mean concentration of $0.87 \mu\text{g/L}$. The total mean deposition rate of airborne copper from rain was measured to be $4.0 \mu\text{g/m}^2/\text{day}$ for the eight sampling sites used in the study (Feng et al. 2000). However, copper deposition from automobile emissions, as measured by the concentration of copper in snow, did not vary significantly as a function of distance (15–150 meters) from an expressway in Montreal, Canada. Mean concentrations of copper in the snow (expressed as mg/L [and standard deviations]) were measured as 0.051 (0.073), 0.065 (0.127), 0.034 (0.027), and 0.044 (0.051) at 15, 20, 15, and 150 meters, respectively (Loranger et al. 1996).

Airborne concentrations of copper in the indoor atmosphere within homes located in Suffolk and Onondaga counties in New York average between 8 and 12 ng/m^3 (Koutrakis et al. 1992). The concentration was significantly affected by the use of kerosene heaters, which were found to emit copper into the indoor air at a rate of 15,630 ng/hour (Koutrakis et al. 1992).

Elevated levels of copper in fog water have been observed 3 km downwind from a refuse incinerator in Switzerland (Johnson et al. 1987). High concentrations of copper were associated with low pH. The maximum concentration, 673 ppb, occurred at pH 1.94; levels >127 ppb were associated with pH values <3.6 . Copper(II) concentrations in fog water from the central valley of California ranged from 1.7 to 388 ppb (Miller et al. 1987). The source of the copper was not investigated. The highest values were recorded just as the fog was dissipating.

6.4.2 Water

Copper is widely distributed in water since it is a naturally occurring element. Copper levels in surface water range from 0.5–1,000 ppb, with a median of 10 ppb; seawater contains $<1\text{--}5$ ppb (Davies and Bennett 1985; Mart and Nurnberg 1984; Page 1981; Perwak et al. 1980; Yeats 1988). The results of several studies in which copper was detected in drinking water, groundwater, and surface water are described in this section and summarized in Table 6-7. The information in Table 6-7 demonstrates that copper concentrations in drinking water can vary widely ($\leq 5\text{--}10,200$ ppb) and can exceed the action limits of 1,300 ppb that have been set for copper in drinking water (EPA 1991). The table also

6. POTENTIAL FOR HUMAN EXPOSURE

emphasizes the importance of running tap water before using it and the need to control corrosion of piping in water distribution systems.

Copper concentrations in drinking water vary widely as a result of variations in pH, hardness of the water supply and copper released from the water distribution system (Davies and Bennett 1985; Yannoni and Piorkowski 1995). Copper concentrations in drinking water range from a few ppb to 10 ppm. A Canadian national survey of copper and other metals in drinking water was conducted from November 1976 to January 1977 (Meranger et al. 1979). Supplies from 70 municipalities representing 38% of the Canadian population were included in the survey, including 50 derived from river or lake water and 20 derived from groundwater. Unfiltered raw, treated and distributed drinking waters were analyzed. Whether the water was derived from river, lake, or well water did not significantly affect the copper concentration in the raw water. Only in a few supplies did copper levels in raw water exceed 20 ppb and only one of these was derived from groundwater. The results in groundwater contrast with those of Page (1981) in New Jersey, in which over 100 wells contained copper levels in excess of 64 ppb. However, that study included groundwater that was a source of drinking water, in addition to groundwater that was not. The copper concentration in Canadian treated water was generally ~10 ppb (Meranger et al. 1979). In 20% of the samples, the copper level in distributed water was significantly higher than the treated water. The increase was greater in areas where the water was soft and corrosive, thus enhancing leaching of copper from the distribution system.

Elevated concentrations of copper in drinking water can result as a consequence of leaching processes that occur in water distribution systems. A study of 1,000 water samples from random households in Ohio found that ~30% contained copper levels >1 ppm (Strain et al. 1984). The highest copper level in the study was 18 ppm. In a study of private water wells in four communities in Nova Scotia, Maessen et al. (1985) found that the concentrations of copper increased in water that remained in the distribution system overnight, indicating that copper was mobilized from the distribution system. Whereas the level of copper in running water was generally very low, that in the standing water was variable and exceeded 1.0 ppm in 53% of the homes. Similar results were reported for U.S. cities (Maessen et al. 1985; Schock and Neff 1988; Strain et al. 1984). In a study in Seattle, Washington, the mean copper concentrations in running and standing water were 0.16 and 0.45 ppm, respectively, and 24% of the standing water samples exceeded 1.0 ppm (Maessen et al. 1985). The difference in copper level between standing and flushed systems became evident at pH 7 and increased with decreasing pH (Strain et al. 1984). Copper levels in school drinking water were found to differ by 3-fold between first draw and 10-minute flush water samples, irrespective of the corrosiveness of the water (Murphy 1993). However, the concentration of

6. POTENTIAL FOR HUMAN EXPOSURE

copper in both first draw and 10-minute flush samples decreased by approximately 10-fold as the corrosiveness of the water decreased. Increasing pH in water distribution lines has been found to result in an overall decrease in metal concentrations. For example, increasing the pH of water from 7.5 to 8.5 in distribution lines decreased copper concentration by 50% (Yannoni and Piorkowski 1995).

In homes with copper piping, the mean concentration of copper in tap water has been shown to decline with the age of the home. In a sampling of tap water of 2,619 households in Berlin, Germany, that are supplied with municipal drinking water, the mean concentration of copper decreased from 0.77 ppm in homes with stated ages of 0–<5 years to 0.23 ppm in homes with stated ages of 35–<40 years (Zietz et al. 2003a). In another study of 1,619 homes in Lower Saxony, Germany, the mean concentration of copper in first draw tap water decreased from 0.37 ppm in homes with stated ages of 0–<5 years to 0.05 ppm in homes with stated ages of 35–<40 years (Zietz et al. 2003b). These decreases of copper concentration with age were attributed to a buildup of a surface layer on the piping that reduced corrosion. However, in these same two studies, it was found that the concentration of copper in tap water began to increase with increasing age in homes with stated ages of >45 years. This increase in copper concentration was attributed to the increased probability of repair or partial placement (or unknown total replacement) of piping in these homes.

In a study of groundwaters and surface waters throughout New Jersey in which >1,000 wells and 600 surface sites were sampled, the median copper levels in groundwater and surface water were 5.0 and 3.0 ppb, respectively (Page 1981). The respective 90th percentile and maximum levels were 64.0 and 2,783.0 ppb for groundwater and 9.0 and 261.0 ppb for surface water. The pattern of contamination in surface water correlates with light hydrocarbons, while that in groundwater correlates with heavy metals. This suggests that the sources of contamination of surface water and groundwater are different. The nature of the sites with elevated levels of copper was not indicated.

The geometric mean (standard deviation) and median concentration of dissolved copper in surface water based on 53,862 occurrences in EPA's STORET database are 4.2 (2.71) and 4.0 ppb, respectively (Eckel and Jacob 1988). Higher concentrations tend to be found in New England, the western Gulf and the lower Colorado River (Perwak et al. 1980). The finding of high concentrations of copper species in minor river basins reported in EPA's STORET database in 1978 revealed that sources of copper in the Gila, Coeur D'Alene, and Sacramento River Basins appear to be primarily mining activities, especially abandoned sites (Perwak et al. 1980). Generally, the high concentrations (>120 µg/L) were generally observed at localized stations and correlated with low pH of the surface water. However, in another study concerning

6. POTENTIAL FOR HUMAN EXPOSURE

75 Canadian headwater lakes sensitive to acid rain, copper values were relatively low (1–8 ppb range, 2 ppb mean) regardless of pH or alkalinity (Reed and Henningson 1984).

Copper concentrations were measured in surface water obtained from sampling sites in the Spearfish Creek, Whitewood Creek, and Bear Butte Creek watersheds. These watersheds are affected by water leaching from tailings and acid-mine drainage from gold mining operations in the Black Hills of South Dakota. Copper concentrations of <0.24–28 µg/L were measured in surface water, whereas concentrations in sediments were much higher, ranging from 7.8 to 159 mg/kg (May et al. 2001).

In a survey of sources of copper in storm water, measurements of copper concentrations in storm water samples were taken from various urban locations in Birmingham, Alabama. Copper concentrations were generally low in filtered samples (dissolved copper), ranging between 1.4 and 20 µg/L; but were much higher in unfiltered samples (copper bound to particulate matter) with mean values (in µg/L) of 110 (roof areas), 116 (parking areas), 280 (street runoff), 135 (vehicle service areas), 81 (landscaped areas), 50 (urban creeks), and 43 (retention ponds) (Pitt et al. 1995).

As a result of improvements in controlling the quality of discharges from municipal and industrial waste water treatment plants mandated in the Clean Water Act, copper concentrations have been declining in surface waters. For example, median copper concentrations in the Hudson River estuary have fallen 36–56% between the mid-1970s and the mid-1990s (Sañudo-Wilhelmy and Gill 1999).

The copper concentration in some bodies of water evidently varies with season. In a study of a small pond in Massachusetts from April of 1971 to March 1972, the concentration of copper was found to vary, decreasing during the spring and early summer to lows of <10–30 ppm in early August and then increasing when the pond was under the cover of ice to a maximum values of 80–105 ppb in late January and early February (Kimball 1973). Similar seasonal variations were noted in the epilimnion of the offshore waters of the Great Lakes (Nriagu et al. 1996). In both examples, the cycling in copper concentrations is thought to be a response to biological need and copper uptake during the growing season and its subsequent release from seasonal die-off and decay of biota.

Copper concentrations in seawater usually are in the 1–5 ppb range (Perwak et al. 1980). Copper levels are overall lower in the Pacific Ocean than in the Atlantic Ocean and higher near the continental shelf than in the open ocean. Copper concentrations in surface water at a depth of 1 meter transected on a cruise from Nova Scotia to the Sargasso sea ranged from 57.2 to 210 parts per trillion (ppt) (Yeats 1988).

6. POTENTIAL FOR HUMAN EXPOSURE

The mean value in surface water sampled at a depth of 1 meter of the eastern Arctic Ocean was 93 ppt (Mart and Nurnberg 1984). As noted in a review by Kennish (1998), concentrations of copper in estuarine and coastal waters in the United States were 0.3–3.8 and 0.1–2.5 ppb, respectively.

6.4.3 Sediment and Soil

Copper occurs naturally at levels of ~50 ppm in the earth's crust, which includes soil and parent rock (Perwak et al. 1980). In the United States, copper concentrations in differing soil types can vary over a large range (1–300 mg/kg, dry weight); but the mean values are relatively similar (14–41 mg/kg, dry weight) as a function of soil type (Table 6-10) and land resource region (Table 6-11) (Chen et al. 1999; Fuhrer 1986). These copper levels are similar to those given in a review of soil copper concentrations that reported a median concentration of 30 ppm (dry weight) and a range of 2–250 ppm (Davies and Bennett 1985). In other studies of copper concentrations in U.S. soils, the mean copper concentrations are within the range of 14–41 mg/kg. In the work of Ma et al. (1997), the mean concentration of copper in soils of the United States was determined to be 30 mg/kg, whereas the copper concentration in agricultural surface soils in the United States that had not received an application of sludges from municipal waste treatment plants was found to be 18 mg/kg. In Florida surface soils, the geometric mean of copper concentration in 40 different soil types was 4.10 mg/kg, with a range of 1.89–10.7 mg/kg (Ma et al. 1997). Chen et al. (1999) reported copper concentrations in Florida soils ranging from 0.1 to 318 mg/kg with a geometric mean of 2.21 ± 3.15 mg/kg (arithmetic mean of 6.10 ± 22.1 mg/kg). These investigators also reported geometric means of 24.0 mg/kg in California soils and 17 mg/kg in U.S. soils. In agriculturally productive soils, copper ranges from 1 to 50 ppm, while in soil derived from mineralized parent material the copper levels may be much higher (NRC 1977; Perwak et al. 1980).

Copper concentrations in soil may be much higher in the vicinity of a source of copper emissions, such as a mining operation or smelter. Concentrations in the top 5 cm of soil near the boundary of a secondary copper smelter were $2,480 \pm 585$ ppm (Davies and Bennett 1985). Maximum wetland soil/sediment copper concentrations were 6,912 ppm in the immediate vicinity of a Sudbury, Ontario, smelter but the concentration decreased logarithmically with increasing distance from the smelter (Taylor and Crowder 1983). The observation that the copper concentrations were highest in soils within 1–2 km from the smelter and decreased exponentially with increasing distance from the plant suggests that copper in the soil from the study area was primarily derived from particulate emissions from the smelter.

6. POTENTIAL FOR HUMAN EXPOSURE

**Table 6-10. Concentration of Copper in Surface Soils of the United States
(in ppm-Dry Weight [dw], Equivalent to mg/kg-dw)**

Soil	Range	Mean
Sandy soils and lithosols on sandstones	1–70	14
Light loamy soils	3–70	25
Loess and soils on silt deposits	7–100	25
Clay and clay loamy soils	7–70	29
Alluvial soils	5–50	27
Soils over granites and gneisses	7–70	24
Soils over volcanic rocks	10–150	41
Soils over limestones and calcareous rocks	7–70	21
Soils on glacial till and drift	15–50	21
Light desert soils	5–100	24
Silty prairie soils	10–50	20
Chernozems and dark prairie soils	10–70	27
Organic light soils	1–100	15
Forest soils	7–150	17
Various soils	3–300	26

Source: EPA 1995

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-11. Geometric Means of Selected Soil Elements and Associated Soil Parameters in U.S. Surface Soils by Land Resource Regions

Land resource region	mg/kg dry soil
Mineral soils	
Northwestern specialty	34.3
Northwestern wheat	23.2
California subtropical	43.4
Western range and irrigated	26.8
Rocky Mountain	19.1
Northern Great Plains	20.2
Western Great Plains	16.3
Central Great Plains	12.6
Southwest Plateau	10.0
Southwest Prairie	4.9
Northern lake states	15.4
Lake states	18.2
Central feed grains	19.7
East and central farming	8.0
Mississippi Delta	21.1
South Atlantic and Gulf slope	6.3
Northeastern forage	34.0
Northern Atlantic slope	13.5
Atlantic and Gulf coast	7.6
Florida subtropical	31.9
All mineral soils	15.6
Histosols	
Northern lake states	59.6
Lake states	84.7
Northeastern forage	149.0
Florida subtropical	94.3
All histosols	86.9

Source: Holmgren et al. 1993

6. POTENTIAL FOR HUMAN EXPOSURE

Copper and its compounds were found at 906 of 1,647 hazardous waste sites on the NPL of highest priority sites for possible remedial action (HazDat 2002). Since copper is commonly found in soil, it should occur at all sites. In past work, data analysis of metal concentrations measured in soil from hazardous waste sites taken from the 1980–1983 Contract Laboratory Program (CLP) Analytical Results Data Base (CARD) was conducted to ascertain whether elemental concentrations at hazardous waste sites were elevated above that which normally would be expected in soil of similar composition and derivation. Of the 1,307 samples in CARD, 10.5 and 7.3% had copper concentrations exceeding the number normally expected in soil at the 95 and 99% confidence intervals, respectively (Eckel and Langley 1988).

In a study in which the copper concentrations of 340 soil samples collected from diverse land-use situations, the average copper concentrations reported were 25 ppm in agricultural land, 50 ppm in suburban/residential land, 100 ppm in mixed industrial/residential land, and 175 ppm in industrial/inner urban areas (Haines 1984). From an analysis of the spatial distribution of the copper concentrations in soils where lowest copper soil concentrations are observed for rural (agricultural) soils and highest in soils obtained from industrialized urban areas, it was concluded that most of the contamination was a result of airborne deposition from industrial sources.

The concentration of copper in soils and sediments was assessed as part of the National Water-Quality Assessment Program (Rice 1999). The median concentrations of copper at 541 sites throughout the conterminous United States ranged from 5 to 70 $\mu\text{g/g}$ (dry weight). At nonurban indicator sites, the median concentrations ranged from 13 to 47 $\mu\text{g/g}$. The same study derived an average crustal abundance of copper of 60 $\mu\text{g/g}$.

Sediment is an important sink and reservoir for copper. In areas where there is no known input of copper obtained from anthropogenic sources, sediment generally contains <50 ppm copper. The level can reach several thousand ppm in polluted areas (Harrison and Bishop 1984). The mean copper level in surficial sediment of Penobscot Bay, Maine, was 14.1 ppm (dry weight), while that in estuaries or bays in other New England locations ranged from 4.4 to 57.7 ppm (Larsen et al. 1983b). Levels reflect anthropogenic input as well as the mineral content of the regional bedrock. Copper levels in sediment from 24 sites along the New Jersey coast ranged from <1.0 to 202 ppm, with a mean value of 66 ppm (Renwick and Edenborn 1983). The texture of the sediment varied from 94% clay to 100% sand, and the copper level was correlated negatively with the percentage of sand in the sediment.

6. POTENTIAL FOR HUMAN EXPOSURE

Surficial sediment in lakes in the Sudbury region of northeastern Ontario, where several smelters operate, decreased rapidly with increasing distance from the smelters (Bradley and Morris 1986). Three lakes, 10 km from the Sudbury smelters, contained copper concentrations in sediment approaching 2,000 ppm dry weight, over 100 times the concentration in a lake selected as a baseline lake 180 km away.

An analysis of the Coastal Sediment Database (COSED) showed that 73% of coastal waterways had copper concentrations below 42 µg/g; 25% had copper concentrations between 42 and 210 µg/g; and 2% were above 210 µg/g. These higher concentrations are associated with locations of high ship traffic, industrial activity, and relatively poor water flushing (Daskalakis and O'Connor 1995). In coastal areas receiving persistently high influxes of contaminants, high concentrations of copper (151 ppm) have been measured to sediments to depths of 54 cm (Bopp et al. 1993). Combined sewer outflows can also contribute significantly to the copper content in sediments. For example, mean (arithmetic) copper concentrations of 180, 208, 280, and 284 mg/kg were measured in sediment samples obtained near four sewer outflows in the lower Passaic River, New Jersey (Iannuzzi et al. 1997). In Jamaica Bay, New York, copper concentrations in sediments were 151–406 ppm, with a concentration of 151 ppm in sediment core samples obtained at a depth of 52–54 cm (Bopp et al. 1993). The highest concentrations were found in the middle depths (16–44 cm) ranging from 280 to 406 ppm during a period where untreated industrial effluents and sewage outflows were allowed to enter the bay. However, copper concentrations in surface sediments (0–2 cm) were measured at 208 ppm. The decrease in copper concentration in the surface sediments suggests that efforts to reduce metal contaminants from sewage outflows have been making an impact on the copper concentrations in receiving waters and their sediments.

6.4.4 Other Environmental Media

In addition to the ingestion of drinking water, the consumption of food is the other primary route for copper intake in the general population. Several studies of copper content in a variety of foods have been conducted as part of the FDA's Total Diet Survey and are described in this section. These data have been used to estimate the average intakes of copper in the human diet within various age groups. For example, in the 25–30-year-old age group, copper intake has been estimated to be 0.93 mg/day for women and 1.24 mg/day for men (Pennington 1983). The levels of copper in other food sources such as mollusks, fish, and agricultural plants have been measured and the results summarized in this section. One highlight in the data is the potential for high dietary intakes of copper for those individuals who regularly consume of mollusks where the daily intake of copper could increase by 5.7–136 mg/day in comparison to the

6. POTENTIAL FOR HUMAN EXPOSURE

general population (see Section 6.5). Other media covered in this section are human tissues, cigarette smoke, industrial and municipal waste streams, and agricultural products.

The FDA Total Diet Survey has provided copper concentration in various foods, examples of which are given in Table 6-12 (FDA 2001). The highest concentrations of copper were found in liver, in some oat and bran cereals, in some legumes and nuts, and in raw avocados and mushrooms. Coleman et al. (1992) reported copper concentrations in the edible tissues of livestock and poultry with the highest mean concentrations (ppm) found in liver (cow 43.7; lamb 89.8; chicken 4.60; turkey 7.14), followed by kidney (cow 8.15; lamb 5.39; chicken 3.07; turkey 3.68), and muscle (cow 1.41; lamb 1.47; chicken 0.67; turkey 0.83) (Coleman et al. 1992).

More recent measurements of copper concentrations in 265 foods analyzed from 1991 to 1996 and from 1991 to 1999 have been obtained from the FDA Total Diet Study (Capar and Cummingsham 2000; FDA 2000). The copper contents of selected foods provided in the most recent FDA Total Diet Study (FDA 2000) are similar to those obtained from the 1982–1984 FDA study. The contribution of food groups to copper intake varies depending on the age group (Pennington and Schoen 1996). For example, animal flesh only contributes to 18% of the copper intake for a 2-year-old child, but contributes to 38% of the copper intake for a 60–65-year-old male. The results of a 1994–1996 Continuing Survey of Food Intakes (CSFII) found that the daily intakes of copper for men and women ages ≥ 60 years old are 1.3 and 1.0 mg/day, respectively (Ma and Betts 2000). In a separate study by Ellis et al. (1997), copper intake for male and female African-Americans ages 21–65 years old was determined to be 1.0 mg/day for both sexes.

Daily intakes of copper and other essential minerals were estimated for eight age-sex groups of the United States population as part of the FDA's Total Diet Study (Pennington et al. 1986). By analyzing the mean mineral content of samples of 234 foods obtained in 24 cities from mid-1982 to mid-1984 and by using previously determined daily intakes of each food as determined from data obtained from the National Food Consumption Survey (1977–1978) and the Second National Health and Nutrition Examination Survey (1976–1980) (Pennington 1983), the daily mineral intake for the age-sex groups was determined. The copper intakes in mg/day of the eight age-sex groups were: 6–11-month-old infant, 0.47; 2-year-old child, 0.58; 14–16-year-old girl, 0.77; 14–16-year-old boy, 1.18; 25–30-year-old woman, 0.93; 25–30-year-old man, 1.24; 60–65-year-old woman, 0.86; and 60–65-year-old man, 1.17. All values were low in terms of the estimated safe and adequate daily dietary intake of this nutrient. The food item with the highest copper level was beef/calf liver (61 ppm).

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-12. Copper Content of Selected Foods (mg/kg)^a

Food description	Mean	SD ^b	Food description	Mean	SD ^b
Breads			green pepper, raw	0.7	0.3
bagel, plain	1.3	0.2	iceberg lettuce, raw	0.2	0.2
cracked wheat bread	1.8	0.2	lima beans, immature, frozen, boiled	1.5	0.2
English muffin, plain, toasted	1.3	0.1	mixed vegetables, frozen, boiled	0.6	0.2
graham crackers	1.5	0.3	mushrooms, raw	2.4	0.6
rye bread	1.5	0.2	okra, fresh/frozen, boiled	0.8	0.3
saltine crackers	1.4	0.1	onion, mature, raw	0.4	0.1
white bread	1.1	0.2	peas, mature, dry, boiled	2.3	0.3
white roll	1.3	0.2	spinach, fresh/frozen, boiled	0.8	0.3
whole wheat bread	2.3	0.3	summer squash, fresh/frozen, boiled	0.5	0.1
Cereal, rice, and pasta			sweet potato, fresh, baked	1.4	0.4
corn flakes	0.5	0.1	tomato, red, raw	0.5	0.2
crisped rice cereal	2.0	0.2	tomato sauce, plain, bottled	1.2	0.4
egg noodles, boiled	1.0	0.2	tomato, stewed, canned	0.7	0.2
granola cereal	3.0	0.4	turnip, fresh/frozen, boiled	0	0.1
macaroni, boiled	0.9	0.1	white potato, baked with skin	1.0	0.4
oatmeal, quick (1–3 minutes), cooked	0.7	0.1	white potato, boiled without skin	0.6	0.2
oatring cereal	3.3	0.4	winter squash, fresh/frozen, baked, mashed	0.6	0.2
raisin bran cereal	4.4	0.4	Fruits		
shredded wheat cereal	3.7	0.5	apple, red, raw	0.2	0.2
wheat cereal, farina, quick (1–3 minutes), cooked	0.3	0.3	applesauce, bottled	0.2	0.1
white rice, cooked	0.7	0.1	apricot, raw	0.8	0.3
Vegetables			avocado, raw	2.2	0.6
asparagus, fresh/frozen, boiled	1.0	0.2	banana, raw	1.1	0.2
beets, fresh/frozen, boiled	0.7	0.2	cantaloupe, raw	0.3	0.1
black olives	1.4	0.4	fruit cocktail, canned in heavy syrup	0.5	0.1
broccoli, fresh/frozen, boiled	0.2	0.1	grapefruit, raw	0.3	0.1
Brussels sprouts, fresh/frozen, boiled	0.4	0.1	grapes, red/green, seedless, raw	1.1	0.6
cabbage, fresh, boiled	0	0	orange, raw	0.4	0.1
carrot, fresh, boiled	0.3	0.2	peach, canned in light/medium syrup	0.3	0.2
cauliflower, fresh/frozen, boiled	0	0	peach, raw	0.7	0.2
celery	0	0.1	pear, canned in light syrup	0.4	0.1
collards, fresh/frozen, boiled	0.5	0.4	pear, raw	0.8	0.1
corn, fresh/frozen, boiled	0.3	0.2	pineapple, canned in juice	0.5	0.1
cream style corn, canned	0.1	0.2	plums, raw	0.6	0.1
cucumber, raw	0.2	0.2	prunes, dried	2.9	0.3
eggplant, fresh, boiled	0.5	0.2	raisins, dried	3.3	0.4
green beans, fresh/frozen, boiled	0.5	0.3	strawberries, raw	0.5	0.3
green peas, fresh/frozen, boiled	1.0	0.2	watermelon, raw	00.4	0.1

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-12. Copper Content of Selected Foods (mg/kg)^a

Food description	Mean	SD ^b	Food description	Mean	SD ^b
Fruit juices			pork roast, baked	0.8	0.1
apple juice, bottled	0	0.1	pork sausage, pan-cooked	0.8	0.1
grape juice, bottled	0	0.1	quarter-pound hamburger on bun, fast-food	0.9	0.1
grapefruit juice, from frozen concentrate	0.3	0.1	salami, sliced	1.0	0.2
orange juice, from frozen concentrate	0.3	0.1	salmon, steaks or filets, fresh or frozen, baked	0.5	0.1
pineapple juice from frozen concentrate	0.4	0.1	shrimp, boiled	2.3	0.6
prune juice	0.1	0.1	tuna, canned in oil	0.5	0.1
tomato juice, bottle	0.6	0.1	turkey breast, roasted	0.4	0.1
			veal cutlet, pan-cooked	1.0	0.3
Dairy products			Legumes, nuts, and nut products		
American, processed cheese	0.1	0.2	kidney beans, dry, boiled	2.7	0.5
cheddar cheese	0.3	0.2	mixed nuts, no peanuts, dry roasted	15.5	2.6
chocolate milk, fluid	0.3	0.2	peanut butter, smooth	5.2	0.6
cottage cheese, 4% milkfat	0	0	peanuts, dry roasted	5.8	0.6
cream cheese	0	0	pinto beans, dry, boiled	2.4	0.2
eggs, boiled/fried	0.6	0.1	pork and beans, canned	1.8	0.2
eggs, scrambled	0.5	0.1			
half & half	0	0	Fats, oils, condiments, snacks, and sweets		
lowfat (2%) milk, fluid	0	0	butter, regular (salted)	0	0
skim milk	0	0	corn chips	1.0	0.2
sour cream	0	0	fruit flavor sherbet	0	0.1
Swiss cheese	0.4	0.4	gelatin dessert, any flavor	0	0
whole milk	0	0	honey	0	0
			jelly, any flavor	0	0.1
Meat, poultry, and seafood			margarine, stick, regular (salted)	0	0
beef chuck roast, baked	1.0	0.1	mayonnaise, regular, bottled	0	0
beef steak, loin, pan-cooked	1.0	0.2	olive/safflower oil	0	0
bologna, sliced	0.4	0.2	popcorn, popped in oil	1.7	0.4
chicken breast, roasted	0.3	0.1	potato chips	2.8	0.8
chicken, fried (breast, leg, and thigh)	0.7	0.1	pretzels, hard, salted, any shape	1.6	0.2
frankfurters, beef, boiled	0.4	0.1	vanilla ice cream	0.06	0.24
ground beef, pan-cooked	0.8	0.1	white sugar, granulated	0	0
haddock, pan-cooked	0.06	0.13			
ham, baked	0.6	0.2	Beverages		
ham luncheon meat, sliced	0.5	0.1	coffee, from ground	0	0
lamb chop, pan-cooked	1.4	0.2	cola carbonated beverage	0	0
liver, beef, fried	123	57	tea, from tea bag	0	0
pork bacon, pan-cooked	1.2	0.4			
pork chop, pan-cooked	0.8	0.2			

^aSource: FDA 2000^bSD = Standard Deviation

6. POTENTIAL FOR HUMAN EXPOSURE

A baseline value for the copper content of mother's milk was determined by a survey of literature values. Based on the data obtained from 28 study sets, a baseline copper concentration of 331 ppb was determined from a range of values of 197–751 ppb and a median of 290 ppb (Iyengar and Woittiez 1988). In a separate study of 11 lactating women, it was found that the variability in the copper content of mother's milk was primarily subject-related. The copper concentration in milk declined moderately, from 0.43 µg/mL between 1 and 3 months postpartum to 0.24 µg/mL between 10 and 12 months postpartum (Vaughan et al. 1979). In a study of 82 lactating women, the copper concentration in breast milk ranged between 0.8 and 1.1 ppm and remained relatively constant in individual women over the first 7 days postpartum (Arnaud and Favier 1995).

The concentrations of copper in the soft tissue in mussels and oysters collected as part of the U.S. Mussel Watch Program in 1976–1978 were 4–10 ppm (dry weight) for mussels and 25–600 ppm for oysters (Goldberg 1986). Copper concentrations in mussels collected from 11 sites near Monterey Bay, California, were 4.63–8.93 ppm (dry weight) (Martin and Castle 1984). Perwak et al. (1980) reported similar results for mussels (3.9–8.5 ppm) and for clams (8.4–171 ppm). Recent measurements of copper concentrations in zebra and quagga mussels taken from Lakes Erie and Ontario in 1997 ranged from 21 to 41 ppm (dry weight) (Rutzke et al. 2000). In the National Oceanic and Atmospheric Administration (NOAA) Mussel Watch Project, copper concentrations were quantified in mollusks (*M. edulis*, *M. californianus*, *C. virginica*, and *Ostrea equestris*) from 113 sites around the United States in 1993 and compared to copper concentrations measured in mollusks taken from the same site in the EPA2 Mussel Watch Program, 1976–1978 (Lauenstein and Daskalakis 1998). The results of the comparison indicate that the decreasing and increasing trends in copper concentrations in mollusks were approximately equal among the sites, except in California, where increasing trends were noted at five sites.

As a part of the National Contaminant Biomonitoring Program of the U.S. Fish and Wildlife Service, eight species of freshwater fish were collected at 112 stations in the United States in 1978–1979 and 1980–1981 (Lowe et al. 1985). The geometric mean concentrations of copper in ppm (wet weight, whole fish) for these two periods were 0.86 and 0.68, respectively; the 85th percentiles were 1.14 and 0.90, respectively, and the ranges were 0.29–38.75 and 0.25–24.10, respectively. The highest concentration, 38.75 and 24.10 ppm, during both collecting periods was in white perch from the Susquehanna River and the second highest concentration, 19.3 ppm, was found in white perch from the Delaware River near Trenton, New Jersey. However, copper concentrations in common carp and white catfish collected from the same station at the same time were 0.76 and 1.35 ppm, respectively.

6. POTENTIAL FOR HUMAN EXPOSURE

In bluefin tuna caught in the northwest Atlantic off Newfoundland, the mean copper concentration in muscle tissue has been measured at 1.0 ppm (dry weight) (Hellou et al. 1992a). In cod caught off the coast of Newfoundland, mean copper concentrations of <1.2–1.5 µg/g (dry weight) in muscle and 5–10 ppm (dry weight) in liver have been determined (Hellou et al. 1992b).

Copper residues in muscle of 268 fish specimens from 17 species were analyzed over a 5-year period in several surface water systems in eastern Tennessee (Blevins and Pancorbo 1986). The mean residue levels in the muscle of different species of fish from nine stations ranged from 0.12–0.86 ppm (wet weight). Maximum levels ranged from 0.14 to 2.2 ppm.

Concentrations of copper in three species of fish living in storm treatment ponds have been compared to copper concentrations in controls collected from surrounding surface waters near Orlando, Florida (Campbell 1994). In bluegill sunfish collected from storm water ponds, the mean whole body copper concentrations were 6.37 and 2.08 mg/kg wet weight, respectively, and were significantly higher than the mean concentrations of copper, 0.879 and 1.07 mg/kg wet weight, respectively, measured in controls collected in natural lakes or ponds. However, in largemouth bass, the mean copper concentrations in fish collected from storm water ponds and controls did not significantly differ, with values of 3.81 and 4.71 mg/kg wet weight, respectively.

Respective mean and median copper concentrations of 127 samples of finfish from Chesapeake Bay and its tributaries were 1.66 and 0.36 ppm in 1978, and 1.85 and 0.61 ppm in 1979 (Eisenberg and Topping 1986). In striped bass taken from Turkey Point in the bay, copper levels were below the detection limit of the study (<0.1 µg/g) in muscle, but were higher in liver tissue ranging from 0.86 to 23.5 µg/g. In gonad tissue obtained from tissue from a different site on the bay, there was also an increase in the mean copper concentration in this tissue (4.25 µg/g) as compared to muscle (0.76 µg/g). The copper content of muscle tissue of several species of fish collected from metal-contaminated lakes near Sudbury, Ontario, ranged from 0.5 to 1.4 ppm (dry weight). No major pattern in variation was evident among species or among the study lakes (Bradley and Morris 1986). The copper concentration in the livers, however, ranged from 5 to 185 ppm (dry weight) and differed significantly among species and among lakes. Unlike muscle tissue, liver tissue is a good indicator of copper availability, although the data indicate that there are other factor(s) that influence the availability and bioaccumulation of copper in these fish.

6. POTENTIAL FOR HUMAN EXPOSURE

The copper concentrations in the liver of lake trout and grayling taken from four fresh water lakes in Alaska did not correlate well with the concentrations of copper in the sediments of these lakes (Allen-Gil et al. 1997). Lake trout were found to have statistically significant higher burdens of copper in their livers than grayling, and the concentrations of copper in the livers of trout varied considerably depending on the lake from which they were collected. The species and site differences in copper concentrations in fish livers have been attributed to differences in diet, (grayling consume mainly insects, whereas trout consume a mix of snails, insects and small fish) and time spent at various depths of the water column.

Although the concentrations of copper in plants vary widely, they usually range from 1 to 50 ppm (dry weight) (Davies and Bennett 1985) and from 1 to 143 ppm (dry weight) in edible plants (Perwak et al. 1980). Concentration ratios of copper in plants relative to soil (concentration factors or CF) demonstrate that copper uptake differs significantly between plants. For example, CF values have been found to vary from 0.02 (onion), 0.13 (celery), 0.21 (lettuce), and 0.30 (potato) to 2 (grapes), 4.5 (alfalfa), and 6.8 (grass) (Pinochet et al. 1999). Concentration factors in rice were found to vary among soil types (0.59–3.58) with copper concentrations in rice ranging from 1.7 to 5.1 $\mu\text{g/g}$ (Herawati et al. 2000). Copper concentrations in rice grain have been found to increase significantly from 1.4 to 15.5 $\mu\text{g/g}$ when copper concentrations in waste water irrigated soils increased from 17.0 mg/kg (wet weight) to 101.2 mg/kg (wet weight) (Cao and Hu 2000).

Studies of copper in human tissues suggest that copper content in a 70 kg adult ranges from 50–70 mg (Davies and Bennett 1985). Wise and Zeisler (1984) reported an average copper concentration of 10 ppm in the human liver in 36 samples. Despite the wide variation in copper concentrations in the environment, the copper concentration in the liver only varied by a factor of 2–3.5. Copper concentrations in human tissues are given in Table 6-13 (Georgopoulos et al. 2001). The concentration of copper in blood is not expected to be predictive of the total body burden of copper: Saltzman et al. (1990) found that the correlation between copper concentrations measured in blood and total body burden was poor ($r=0.54$).

The mean copper content of tobacco in Finnish cigarettes was 24.7 ppm, with a standard deviation of 10.8 ppm (Mussalo-Rauhamaa et al. 1986). However, only 0.2% of this copper passes into mainstream smoke. This translates to a daily exposure of approximately 1 μg of copper in a pack of 20 cigarettes.

In an EPA-sponsored study conducted to determine the metal concentration in sewage sludge (Feiler et al. 1980), copper concentrations in primary sludge at seven POTWs were reported to be 3.0–77.4 ppm, with

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-13. Copper Content of Human Tissues and Body Fluids

Tissue	Mean content ($\mu\text{g/g}$ dry weight)	
	Normal	Wilson's disease
Adrenal	7.4	17.6
Aorta	6.7	—
Bone	4.2	—
Brain	—	—
Caudate nucleus	—	212
Cerebellum	—	261
Frontal lobe cortex	—	118
Globus pallidus	—	255
Putamen	—	314
Cornea	—	92.9
Erythrocytes (per 100 mL packed red blood cells)	23.1	—
Hair	89.1	—
Heart	16.5	12.7
Kidney	14.9	96.2
Leukocytes (per 10 ⁹ cells)	0.9	—
Liver	25.5	584
Lung	9.5	15.5
Muscle	5.4	25.9
Nail	18.1	—
Ovary	8.1	5.2
Pancreas	7.4	4.2
Placenta	13.5	—
Prostate	6.5	—
Skin	2	5.2
Spleen	6.8	5.6
Stomach and intestines	12.6	22.9
Thymus	6.7	—
Thyroid	6.1	—
Uterus	8.4	—
Aqueous humor	12.4	—
Bile (common duct)	1,050	173
Cerebrospinal fluid	27.8	—
Gastric juice	28.1	—
Pancreatic juice	28.4	—
Plasma, Wilson's disease	—	—
Saliva	50	—

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-13. Copper Content of Human Tissues and Body Fluids

Tissue	Mean content ($\mu\text{g/g}$ dry weight)	
	Normal	Wilson's disease
Serum		
Female	120	—
Male	109	—
Newborn	36	—
Sweat		
Female	148	—
Male	55	—
Tissue		
Synovial fluid	21	—
Urine (24-hour)	18	—

Source: Georgopoulos et al. 2001; Scheinberg 1979; Sternlieb and Scheinberg 1977

6. POTENTIAL FOR HUMAN EXPOSURE

a median concentration of 20.5 ppm. The plant with the highest copper concentrations received wastes from plating industries, foundries, and coking plants. In a comprehensive survey of heavy metals in sewage sludge, 30 sludges from 23 American cities were analyzed (Mumma et al. 1984). The copper concentration in the sludges ranged from 126 to 7,729 ppm (dry weight), with a median value of 991 ppm. Gutenmann et al. (1994) report similar concentrations (217–793 ppm, dry weight) in sewage sludge obtained from 16 major cities in the United States. The proposed limit for copper in sludge spread on agricultural land is 1,000 ppm (Mumma et al. 1984). For comparison, the concentration of copper in cow's manure is ~5 ppm (Mumma et al. 1984).

In municipal solid waste compost obtained from nine sites in the United States, a mean copper concentration of 281 mg/kg (dry weight) was obtained with range of 36.4–424 mg/kg (He et al. 1995). Lisk et al. (1992) reported copper concentrations in composts formed from yard waste ranging from 22.7 to 327 ppm, from sewage sludge ranging from 432 to 1,019 ppm and from municipal solid waste ranging from 191 to 1,143 ppm.

Copper concentrations in waste from the combustion of municipal solid waste and other combustion processes have been reported. Copper in incinerator bottom ash and fly ash has been measured at mean concentrations of 1,700 and 1,000 mg/kg, respectively (Goldin et al. 1992). Buchholz and Landberger (1995) report concentrations of copper of 390–530 µg/g in fly ash, 1,560–2,110 µg/g in bottom ash, and 1,140–1,540 µg/g in combined ash. In sewage sludge incineration process steams, copper concentrations were 4,561 mg/kg in sludge cake, 3,465 mg/kg in bottom ash, 3,707 mg/kg in cyclone ash, 3,684 mg/kg in scrubber particulate matter, and 6,666 mg/kg in stack particulate matter (Balogh 1996). In fossil fuel wastes, copper concentrations of 33–2,200 mg/kg in fly ash, 4–930 mg/kg in bottom ash, 6–340 mg/kg in flue gas desulfurization sludge, 10–130,000 mg/kg oil ash, and 2–190 mg/kg in coal have been obtained (Eary et al. 1990).

Agricultural sources of copper contamination in soils has been summarized by EPA (1995) and are shown in Table 6-14. Concentrations of copper in fertilizers, soil amendments and other agricultural materials have been measured by Raven and Loeppert (1997). The materials and mean concentrations: urea (<0.6 µg/g), ammonium nitrate (<0.6 µg/g), ammonium sulfate (<0.6 µg/g), ammonium phosphate (<2–41.8 µg/g), potassium chloride (<2–3.5 µg/g), potassium-magnesium-sulfate (1.4–5 µg/g), North Carolina rock phosphate (9.6 µg/g), calcite (2.3 µg/g), corn leaves (9.4 µg/g), manure (17.5 µg/g), and austinite (300 µg/g).

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-14. Agricultural Sources of Copper Contamination in Soils

Source	Concentration (ppm dry weight) ^a
Sewage sludges	50–3,300
Phosphate fertilizers	1–300
Limestones	2–125
Nitrogen fertilizers	<1–15
Manure	2–60
Pesticides (percent)	12–50

Source: EPA 1995

^aEquivalent to mg/kg-dry weight

6. POTENTIAL FOR HUMAN EXPOSURE

6.5 GENERAL POPULATION AND OCCUPATIONAL EXPOSURE

Due to the ubiquitousness of copper in the environment and the general occurrence of copper in airborne particulates, exposure to copper through inhalation is commonplace. Estimates of atmospheric copper concentrations from different source categories (e.g., smelters, ore processing, steel production, and combustion) yielded a maximum annual concentration of $30 \mu\text{g}/\text{m}^3$ (EPA 1987a). If a person is assumed to inhale 20 m^3 of air/day, this would amount to an average daily intake of 600 μg of copper. For the reported range of annual atmospheric copper concentrations, 5–200 ng/m^3 (EPA 1987a), the average daily intake by inhalation, would range from 0.1 to 4.0 μg . At the maximum reported ambient air concentration, $100 \mu\text{g}/\text{m}^3$ for a 24-hour period at a location within one-half mile of a major source (EPA 1987a), the average daily intake would rise to 2,000 μg . These estimates assume that all of the copper is attached to particles of inhalable size, which is usually not the case. The average daily dietary intake of copper from food is $\sim 2 \text{ mg}/\text{day}$. The dietary intake of copper is expected to be above this average for those individuals who regularly consume organ meats (e.g., liver and kidney), nuts, seeds (including cocoa powder), legumes, and bran and germ portions of grains; these intakes are not expected to exceed the maximum recommended limits of 10–12 mg/day (WHO 1996). Those individuals who regularly consume oysters or clams may increase their dietary intake of copper by 2–150 mg/day when consuming 250 g of edible tissue per day, based on copper concentrations of 25–600 and 8.4–171 ppm in oysters and clams, respectively (Goldberg 1986; Perwak et al. 1980). Assuming a median copper concentration in drinking water of $75 \mu\text{g}/\text{L}$, the average daily copper exposure from consumption of 2 L of water per day is 0.15 mg . However, many people may have high levels of copper in their tap water that have been acquired during transport through the water distribution system. If the system is not permitted to flush out, average intakes from water may be $>2 \text{ mg}/\text{day}$. It is less likely that high dermal exposures will result from bathing in this tap water because the distribution system will flush itself out as the water is drawn. The total exposure of copper for the average person from all sources (e.g., air, drinking water, and food) is estimated to be 2.75 mg/day .

A National Occupational Exposure Survey (NOES) conducted by NIOSH from 1981 to 1983 estimated that potentially 505,982 workers, including 42,557 women, were occupationally exposed to copper in the United States (NIOSH 1988). The NOES estimate is provisional because all of the data for trade name products that may contain copper have not been analyzed. Of the potential exposures, 1,073 are to pure copper, while in the other cases, the molecular form of copper was unspecified. Additionally, according to the NOES, 125,045 workers, including 38,075 women, were potentially exposed to copper sulfate

6. POTENTIAL FOR HUMAN EXPOSURE

(NIOSH 1988). The NOES was based on field surveys of 4,490 facilities and was designed as a nationwide survey based on a statistically valid sample of virtually all workplace environments in the United States where eight or more persons are employed in all standard industrial codes (SIC) except mining and agriculture. The exclusion of mining and agriculture is significant for estimating exposure to copper since there is a high potential for exposure in these industries. Current occupational exposure limits for copper fume are 0.2 and 1 mg/m³ for dust and mists (Frazier and Hage 1998).

6.6 EXPOSURES OF CHILDREN

This section focuses on exposures from conception to maturity at 18 years in humans. Differences from adults in susceptibility to hazardous substances are discussed in Section 3.7 Children's Susceptibility.

Children are not small adults. A child's exposure may differ from an adult's exposure in many ways. Children drink more fluids, eat more food, breathe more air per kilogram of body weight, and have a larger skin surface in proportion to their body volume. A child's diet often differs from that of adults. The developing human's source of nutrition changes with age: from placental nourishment to breast milk or formula to the diet of older children who eat more of certain types of foods than adults. A child's behavior and lifestyle also influence exposure. Children crawl on the floor, put things in their mouths, sometimes eat inappropriate things (such as dirt or paint chips), and spend more time outdoors. Children also are closer to the ground, and they do not use the judgment of adults to avoid hazards (NRC 1993).

With respect to inhalation, exposures of children to copper are not expected to be very different from those of the rest of the general population. However, exposure of copper through oral routes may differ, due to differences in the consumption of various food groups between children and adults and ingestion of dust and soils. The dietary copper intake for infants who receive the major portion of their nutritional requirements from breast milk is likely to be different from infants whose nutritional needs are either supplemented or entirely received through the consumption of formula. Estimates of copper intake from inhalation and ingestion in children in the United States are limited. From the work of Pennington et al. (1986), the copper intakes for a 6–11-month-old infant and a 2-year-old child were estimated to be 0.47 and 0.58 mg/day, values which are lower than the adult intake of ~1 mg/day. However, one study has provided estimated inhalation and ingestion exposures of copper for children in India (Raghunath et al. 1997). In this work, concentrations of copper in particulates in air were measured at 0.01–0.26 µg/m³. Based on these measurements, estimated inhalation exposures of children to copper were calculated to be

6. POTENTIAL FOR HUMAN EXPOSURE

0.1–3.2 $\mu\text{g}/\text{day}$. In this same work, exposures to copper through ingestion were estimated to be between 684–1,732 $\mu\text{g}/\text{day}$.

Exposures of children to copper are likely to increase in areas where copper concentrations in air are expected to be high, such as mining sites, waste dump sites, smelters, and foundries. For example, copper burdens in children living near a lead smelter, as measured by copper concentration in teeth, increased with decreasing distance from the smelter (Blanuša et al. 1990). Children are also at risk for increased copper intake through consumption of drinking water where leaching of copper from the distribution system has occurred (Murphy 1993; Yannoni and Piorkowski 1995). This route of copper exposure can be minimized through the flushing of drinking water supply lines or increasing the pH of the water in the distribution system.

6.7 POPULATIONS WITH POTENTIALLY HIGH EXPOSURES

In discussing exposure to copper, the important question is whether individuals are exposed to readily available copper, which in general, means free (hydrated) $\text{Cu}(\text{II})$ and perhaps some weakly complexed or adsorbed forms of copper. The data indicate that copper in natural water, sediment, and soil mainly exists in bound form. Even so, the free form of copper can be released readily from ingested materials, for example a child's sampling of soil, following exposure to the low pHs encountered in the stomach (Pizarro et al. 2001). Potential for high uptake copper in the general population may exist where people consume large amounts of tap water that has picked up copper from the distribution system, or already has a high copper background due to natural or anthropogenic activities (e.g., close proximity to mining activities or mine drainage). Leaching of copper from water distribution systems is likely to occur where the water is soft and not allowed to run to flush out the system. In such cases, the concentration of copper frequently exceeds 1 ppm, a large fraction of the copper may be in the form of free cupric ion, and uptake will result by ingestion and, perhaps, dermal contact. Soluble cupric salts are used extensively in agriculture and in water treatment. Workers engaged in the formulation and application of these chemicals and industrial workers, such as those in the plating industry, may come into dermal contact with these chemicals. Exposure to high levels of free $\text{Cu}(\text{II})$ may occur, for example, from swimming in water that has been recently treated with a copper-containing algicide.

Based on the available data, people living close to NPL sites may be at greater risk for exposure to copper than the general population. In this case, exposure can occur through inhalation of airborne particulates from the NPL sites, ingestion of water from private wells which are in close proximity to the sites,

6. POTENTIAL FOR HUMAN EXPOSURE

ingestion of contaminated soil, and/or uptake of copper into fruits and vegetables raised in gardens of residents living near NPL sites.

People living near copper smelters and refineries and workers in these and other industries may be exposed to high levels of dust-borne copper by both inhalation and ingestion. In some industries, workers may be exposed to fumes or very fine dust that may be more hazardous than coarse-grained dust, because it can be inhaled more deeply into the lung, thereby evading the mucocilliary escalator.

6.8 ADEQUACY OF THE DATABASE

Section 104(i)(5) of CERCLA, as amended, directs the Administrator of the ATSDR (in consultation with the Administrator of EPA and agencies and programs of the Public Health Service) to pursue assessment of the adequacy of the available information on the health effects of copper. Where adequate information is not available, the ATSDR, in conjunction with the National Toxicology Program (NTP), is required to assure the initiation of a program of research designed to determine the health effects (and techniques for developing methods to determine such health effects) of copper.

The following categories of possible data needs have been identified by a joint team of scientists from the ATSDR, NTP, and EPA. They are defined as substance-specific informational needs that, if met, would reduce uncertainties of regarding human health assessment. This definition should not be interpreted to mean that all data needs discussed in this section must be filled. In the future, the identified data needs will be evaluated and prioritized and a refined substance-specific research agenda will be proposed.

6.8.1 Identification of Data Needs

Physical and Chemical Properties. In general, the available data on the physical and chemical properties of elemental copper and copper sulfate are sufficient for estimating their environmental fate. That no numerical value is listed for the water solubility of copper in Table 4-3 is of no special significance. For inorganic salts, the solubility product coupled with stability constants for the ionic species in solution are the factors determining how much of a compound goes into solution (i.e., the concentration). The solubility products and stability constants for copper that are required for determining the copper species in natural water and their concentrations are known (Schnoor et al. 1987; Town and Filella 2000). Although no K_{oc} values are listed, copper binds very strongly to organic matter, and values for the binding constants and solubility products to humic acids are available (Schnoor et al.

6. POTENTIAL FOR HUMAN EXPOSURE

1987). Similarly, there are binding constants and solubility products for other species that bind or coprecipitate with copper, such as clay minerals and iron and manganese oxides (Schnoor et al. 1987). Binding constants for copper in specific natural waters are also available (Town and Filella 2000). Other physical and chemical properties in Table 4-3 for which there is no data are not well defined for these copper compounds.

In general, experimental confirmation is required for predicting copper's fate in the environment. The factors which determine the copper species present or the material to which copper may be bound and the strength of the binding can be site specific. If the level of detail requires knowledge of, for example, the percentage of copper associated with iron oxides or that which is easily exchangeable, experimental confirmation is necessary.

Production, Import/Export, Use, Release, and Disposal. Information on the production, use, release, and disposal of copper is used for evaluating the potential for exposure of people to copper who live or work near waste sites and other sources. Copper exposure is widespread; but much of this exposure is to generally benign forms, such as metallic copper. The information available often does not distinguish between these forms and those of greater toxicological significance.

Information on the production, use, release, and disposal of metallic copper and copper sulfate is generally available. These two forms of copper account for most of the copper used. This information is tabulated by the U.S. Geological Survey every year in the Minerals Yearbook and predictions of future trends in production and use are available. Such information is not available for other copper compounds. We also know the major uses of copper and where these uses occur (e.g., the home, workplace, etc.).

According to the Emergency Planning and Community Right-to-Know Act of 1986 (EPCRTKA), (§313), (Pub. L. 99-499, Title III, §313), industries are required to submit release information to the EPA. The TRI contains release information for copper and copper compounds and is updated yearly.

For disposal, industrial waste copper is generally either recycled or landfilled. Data on secondary copper production (i.e., copper produced from scrap) is compiled by the U.S. Geological Survey. Effluent and disposal regulations for copper and its compounds are listed in the Clean Water Act and the Resource Conservation and Recovery Act (RCRA).

6. POTENTIAL FOR HUMAN EXPOSURE

Environmental Fate. Reliable information on how copper and its compounds partition in the environment (i.e., to soil and sediment), and the type of transformations that occur in different media, is available from over 35 published studies included in this chapter. We also have data concerning its transport in the environment from over 50 reliable studies. Although information on the fate of copper in air, water, and soil is available, the fate of copper is both species- and site- specific. Information concerning the forms of copper (i.e., specific compound, to what it is bound or complexed, or, in the case of air, the particle size) or the lability of the copper in particular media is available from only a few, -yet reliable, studies. These are sufficient to identify numerous contributors to the fate of copper and its compounds; but are insufficiently comprehensive for developing accurate fate maps. In addition, studies of how fate data relate to human exposures, especially in regard to projecting copper toxicity in children, is inadequate.

Bioavailability from Environmental Media. Copper is found in food, water, ambient air, and soil. The bioavailability of copper from food and water has been investigated in animals and humans. No information on the availability of copper from air was located. Copper in air originating from smelter sites is predominantly associated with sulfur, and presumably exists as the sulfate. Copper dust from soil in general as well as around mining and smelter sites may occur in ore dust or a silicate. No information was located on the bioavailability of copper in air. Copper in soil often is bound to organic molecules. Therefore, the bioavailability of the copper from soil cannot be assessed based on bioavailability information from drinking water or food studies. Studies on the bioavailability of copper from soil and ambient air would be useful in assessing potential toxicity to people living near a hazardous waste site.

The form and lability of copper in the environment is known in only a few site-specific cases. None of these cases include hazardous waste sites. More information on the forms of copper found at industrial sites and hazardous waste sites would be useful, especially since data from the Hazardous Substances Data Bank (HSDB) indicate that concentrations of copper as high as 182,000 ppm in soil and 14,000 ppm in sediments have been measured offsite of listed NPL sites (HazDat 2002). Monitoring groundwater near industries that use highly acid, copper-containing solutions, such as electroplating, electrowinning, and ore leaching industries, is important for the protection of human populations at risk of exposure to their highly mobile and highly bioavailable copper to human risk populations.

Food Chain Bioaccumulation. Because copper occurs in different forms in the environment, its bioaccumulation is expected to vary according to site and species. Data are available on the bioconcentration of copper in aquatic organisms, plants, and animals, as well as biomagnification in food

6. POTENTIAL FOR HUMAN EXPOSURE

chains. This information is useful in assessing the potential for exposure from ingesting food originating from contaminated areas. However, little information is available on the potential for intoxication from foodstuffs from apparently nonpolluted areas or where they may have accumulated toxic levels of copper through biomagnification resulting from foraging in polluted areas.

Exposure Levels in Environmental Media. Data are available regarding the concentrations of copper in environmental media, including the concentration of copper in soil at some hazardous waste sites. Since copper is naturally present in soil, trace quantitative analytical and statistical techniques can be used to determine whether the copper found at these sites is elevated above normal levels. Monitoring data are reasonably current and human intake of copper from food, water, and air can be estimated.

Exposure Levels in Humans. There are reasonably current data on levels of copper in human tissue and human milk. Although information on copper concentrations in individuals exposed within specific work settings is increasing (for example, Gerhardsson et al. 1993; Saltzman et al. 1990), none of the studies address specific U.S. populations living around hazardous waste sites. There are some quantitative data relating occupation, level and route of exposure to the form of copper to which people are exposed. There is some limited information correlating copper concentration and form to body burden in the general population. However, more information is needed for occupational and other at-risk populations.

Exposures of Children. Reasonably current data report levels of copper intake in infants and children. Information on copper intake by infants from human milk also is available. Exposure of children to copper in drinking water has been assessed and methods to decrease this exposure have been identified and implemented. However, only limited information on inhalation and ingestion is available. Some information on exposure of children to copper near mining, smelting, refining, manufacture facilities, waste sites and other hazardous sites is available; but not for U.S. populations. This information is needed to better estimate exposures of children in U.S. populations living near these facilities and sites. The use of copper concentrations in toenails and hair has been investigated as a surrogate measure of copper exposure in children and adults and more research into establishing the validity of these surrogates is underway.

Child health data needs relating to susceptibility are discussed in Section 3.12.2 Identification of Data Needs: Children's Susceptibility.

6. POTENTIAL FOR HUMAN EXPOSURE

Exposure Registries. No exposure registries for copper and its compounds were located. No subregistry has currently been established for these chemicals. They will be considered in the future when chemical selection is made for subregistries to be established. The information that is amassed in the National Exposure Registry facilitates epidemiological research needed to assess adverse health outcomes that may be related to the exposure to these chemicals.

6.8.2 Ongoing Studies

Ongoing studies of copper in soils, sediments and aquifers have been identified and are listed in Table 6-15. Also included in Table 6-15 are ongoing investigations of human exposures to copper.

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-15. Ongoing Studies on Environmental Fate and the Potential for Human Exposure to Copper

Investigator	Affiliation	Research description	Sponsor ^a
Gardea-Torresdey J	Stanford University	Uptake of copper and lead into creosote bushes in regions with heavy metal contamination	National Center for Research Services
Naqvi SM	Southern University A&M	Bioaccumulation and biomagnification of copper in crayfish	National Institute of General Medical Sciences
Conklin MH	University of Arizona College of Pharmacy	Characterization of abandoned mine sites and mine wastes in Arizona; assess stability of these sites to emission of copper and other metals to surface waters	National Institute of Environmental Health Sciences
Sparks DL	University of Delaware	Influence of aging and competitive sorption on stabilization of metals through surface precipitates in soils	CSREES Delaware
Reeve AS	University of Maine	Use of stable isotope tracing techniques to determine the source of salts in Maine groundwater, including copper	CSREES Maine
Welch RM	Agricultural Research Service, Ithaca, New York	Study the soil chemistry, distribution, and bioavailability to crops of health-related elements (e.g., Fe, Zn, Ca, Se, Cd, B) and their movement into edible plant parts.	ARS
Ahner BA	Cornell University	Determine the nature and source of organic ligands in marine waters to better understand the cycling and bioavailability of Cu and Fe.	SAES New York
Hesterberg DL	North Carolina State University	Determine the significance of heavy-metal sulfides (and other stable chemical species) for reducing the mobility and bioavailability of potentially-toxic metals in complex clay-organic systems.	CSREES North Carolina

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-15. Ongoing Studies on Environmental Fate and the Potential for Human Exposure to Copper

Hunt JR	Agricultural Research Center, Grand Forks, North Dakota	Determine how changes in the U.S. diet may affect nutritional status, with emphasis on intakes and bioavailabilities of iron, zinc, copper, and selenium	ARS
Zelazny LW	Virginia Polytechnic Institute	Determine and compare the quantity, chemical forms, and plant available levels of P, Cu, and Zn from manures treated with phytase after reacting with selected soils for various time periods	CSREES Virginia

Source: CRIS 2003; FEDRIP 2003

^aARS = Agricultural Research Service; CSREES = Cooperative State Research, Education, and Extension Service; SAES = State Agricultural Research Station–Multistate Research Projects

7. ANALYTICAL METHODS

The purpose of this chapter is to describe the analytical methods that are available for detecting, measuring, and/or monitoring copper, its metabolites, and other biomarkers of exposure to and affects of copper. The intent is not to provide an exhaustive list of analytical methods. Rather, the intention is to identify well-established methods that are used as the standard methods of analysis. Many of the analytical methods used for environmental samples are the methods approved by federal agencies and organizations such as EPA and the National Institute for Occupational Safety and Health (NIOSH). Other methods presented in this chapter are those that are approved by groups such as the Association of Official Analytical Chemists (AOAC) and the American Public Health Association (APHA). Additionally, analytical methods are included that modify previously used methods to obtain lower detection limits and/or to improve accuracy and precision.

7.1 BIOLOGICAL MATERIALS

Analytical methods and detection limits for copper in biological materials are given in Table 7-1. Copper in other biological materials such as hair and nails can be determined by using suitable procedures for dissolving the sample matrix and employing the same analytical techniques as with blood and tissue. These methods determine the total amount of copper in the sample. The methodology for analyzing biological material is similar to that used for environmental samples. The most commonly employed methods use atomic adsorption spectroscopy (AAS) or inductively coupled plasma-atomic emission spectroscopy (ICP-AES) (Araki et al. 1990; Lo and Araki 1989; Lopez-Artiguez et al. 1993). Differential-pulse anodic stripping voltammetry techniques have also been used to quantify copper in urine, yielding detection limits of 0.041 $\mu\text{g/L}$ and an accuracy of 97% (Horng 1996).

7.2 ENVIRONMENTAL SAMPLES

Analytical methods and detection limits for copper in environmental media are given in Table 7-2. Analytical methods determine the total copper content of the samples. Determining specific copper compounds and complexes in samples is difficult. The most common methods used for environmental samples are AAS, either flame or graphite furnace, ICP-AES, and inductively coupled plasma-mass spectrometry (ICP-MS). Water and waste water samples can be analyzed for copper by EPA Test Method 200.1 (flame atomic absorption), 200.7 ICP-AES, or EPA Test Method 200.9 (temperature

7. ANALYTICAL METHODS

Table 7-1. Analytical Methods for Determining Copper in Biological Materials

Sample matrix	Preparation method	Analytical method	Sample detection limit	Percent recovery	Reference
Blood or tissue	Acid digestion	Method 8005 ^a ; ICP-AES	1 µg/100 mL blood; 0.2 µg/g tissue	Not available	NIOSH 1987
Urine	Filter and polydithio-carbamate resin collection followed by low temperature plasma ashing or acid digestion	Method 8310 ^a ; ICP-AES	0.1 µg	Not available	NIOSH 1987
Tissue	HNO ₃ digestion	AAS/graphite furnace	0.25 µg/g wet weight	103.1±7.7% mean recovery; 8.2±6.9% mean difference in duplicates ^b ; 0.01% accuracy	Lowe et al. 1985
Toenails	HNO ₃ digestion	AAS/graphite furnace	0.6 µg/g	<5% within run precision; 3.5% day-to-day precision	Wilhelm et al. 1991

^aSimultaneous, multielemental analysis, not compound specific.

^bMean±1 standard deviation

AAS = atomic absorption spectrometry; ICP-AES = inductively coupled plasma-atomic emission spectroscopy

7. ANALYTICAL METHODS

Table 7-2. Analytical Methods for Determining Copper in Environmental Samples

Sample matrix	Preparation method	Analytical method	Sample detection limit	Percent recovery	Reference
Air	Filter collection on 0.8 m μ membrane filter and acid digestion	Method 730, ICP-AES	1 μ g	No bias identified	NIOSH 1987
Air	Filter collection on 0.8 m μ membrane filter and acid digestion	Method 7029, AAS	0.05 μ g	No significant bias	NIOSH 1987
Water, waste water	Acidify with 1:1 HNO ₃ to a pH<2	Method 220.1, AAS/direct aspiration	20 μ g/L	0.9–29.7% bias between 7.5 and 332 μ g/L	EPA 1983
Water, waste water	Sample solutions should contain 0.5% HNO ₃	Method 220.2, AAS/furnace technique	1 μ g/L	Not available	EPA 1983
Water, waste water	Filter and acidify sample	Method 200.7 CLP-M ICP-AES	6 μ g/L	Not available	EMMI 1997
Water, waste water	Digestion with H ₂ SO ₄ and HNO ₃	Neocuproine, spectrometric	120 μ g/L in 1 cm cell	Not available	Greenberg et al. 1985
Waste water	Adjust pH to 1.65–1.85, mix, filter	Method 200.1, flame atomic absorption	4 mg/L	Not available	EMMI 1997
Water, waste water	Filter and acidify	Method 200.7_M, ICP-AES	25 μ g/L	Not available	EMMI 1997
Groundwater, surface water, and drinking water	Filter and acidify	Method 200.8, ICP-MS	20 μ g/L	Not available	EMMI 1997
Marine waters	Digest in HNO ₃ , concentrate on iminodiacetate chelating resin, elute with 1.25 M HNO ₃	Method 200.10, ICP-MS	7 μ g/L	Not available	EMMI 1997
Marine waters, estuarine waters, seawaters, and brines	Digest in HNO ₃ , concentrate on iminodiacetate chelating resin, elute with 1.25 M HNO ₃	Method 200.13, GFAA	5 μ g/L	Not available	EMMI 1997
Soil, sediment, sludge, and solid waste	Digestion with HNO ₃ and H ₂ O ₂ , reflux with dilute HCl	Method 7210, AAS	20 μ g/L	As in Method 220.1	EPA 1986

7. ANALYTICAL METHODS

Table 7-2. Analytical Methods for Determining Copper in Environmental Samples

Sample matrix	Preparation method	Analytical method	Sample detection limit	Percent recovery	Reference
Food	Closed-system digestion	AAS or ASV	0.32 µg/g (ASV), not reported (AAS)	94–100	Holak 1983
Biological tissues	HNO ₃ digestion, reaction with H ₂ O ₂	Method 200.3, ICP-MS	18 µg/L	Not available	EMMI 1997
Fish tissue (fresh edible tissue)	Dissociate tissue in tetraammonium hydroxide, acidify with HNO ₃	Method 200.11, ICP-AES	18 µg/L	Not available	EMMI 1997

AAS = atomic absorption spectrometry; ASV = anodic stripping voltammetry; GFAA = graphite furnace atomic absorption; ICP-AES = inductively coupled plasma-atomic emission spectroscopy; ICP-MS = inductively coupled plasma-mass spectrometry

7. ANALYTICAL METHODS

stabilized graphite furnace atomic absorption spectroscopy) (EMMI 1997). These methods are suitable for groundwater and surface water as well as domestic and industrial effluents. EPA Test Method 200.8 ICP-MS or EPA Test Method 200.15 ICP-AES are suitable for analysis of groundwater, surface water, and drinking water. EPA Test Method 200.8, EPA Test Method 200.10 (on-line chelation and ICP-MS), or EPA Test Method 200.13 (chelation and graphite furnace atomic absorption spectroscopy) are suitable for marine, estuary, and brine waters. If determination of dissolved and suspended copper is required, samples should be filtered using a 0.45 µm membrane filter. Suspended solids, as well as sludge and sediment, may be analyzed by EPA Methods 200.1 and 200.13 after an initial acid digestion with HNO₃. Interference by other elements is not a problem in the analysis. However, background correction may be required when using atomic absorption spectroscopy to correct for nonspecific absorption and scattering, which may be significant at the analytical wavelength, 324.7 nm (EPA 1986). In the determination of trace metals, major concerns are contamination and loss. Contamination can be introduced from impurities in reagents and containers as well as from laboratory dust. Losses may also occur due to adsorption onto containers.

Other analytical methods used for copper analysis include x-ray fluorescence, anodic stripping voltammetry, neutron activation analysis, photon-induced x-ray emission, as well as chemical derivatization, followed by gas chromatographic or liquid chromatographic analysis. Discussion of these methods is beyond the scope of this profile. However, methodology for the determination of copper has been reviewed by Gross et al. (1987) for food, by Fox (1987) for air, by MacCarthy and Klusman (1987) for water, and Lichte et al. (1987) for geological materials.

7.3 ADEQUACY OF THE DATABASE

Section 104(i)(5) of CERCLA, as amended, directs the Administrator of ATSDR (in consultation with the Administrator of EPA and agencies and programs of the Public Health Service) to assess whether adequate information on the health effects of copper is available. Where adequate information is not available, ATSDR, in conjunction with the National Toxicology Program (NTP), is required to assure the initiation of a program of research designed to determine the health effects (and techniques for developing methods to determine such health effects) of copper.

The following categories of possible data needs have been identified by a joint team of scientists from ATSDR, NTP, and EPA. They are defined as substance-specific informational needs that if met would reduce certain uncertainties of human health assessment. This definition should not be interpreted to

7. ANALYTICAL METHODS

mean that all data needs discussed in this section must be filled. In the future, the identified data needs will be evaluated and prioritized and a substance-specific research agenda will be proposed.

7.3.1 Identification of Data Needs

Methods for Determining Biomarkers of Exposure and Effect.

Exposure. Methods for determining background and elevated levels of copper in biological materials are well developed, sensitive, specific, and reliable. Standardized methods are available from NIOSH and other sources. The use of copper concentrations in toenails and hair has been investigated as surrogate markers of copper exposure, with validation studies currently underway.

Effect. No specific biomarkers of copper toxicity have been determined. Until such biomarkers are determined, the methodology needed to identify them cannot be established.

Methods for Determining Parent Compounds and Degradation Products in Environmental

Media. Methods for determining background and elevated levels of copper in environmental media are well-developed, sensitive, and selective. Water is the medium of most concern, since the form of copper generally associated with health effects is soluble copper(II). Standardized methods of analysis for copper in air, water, soil, and food are available from EPA, NIOSH, and other sources. Analytical methods measure total copper. Therefore, the methods can not specifically analyze for a parent compound and a degradation product.

7.3.2 Ongoing Studies

Ongoing studies regarding new analytical methods for measuring copper in biological materials or environmental media were located in the literature. Dr. M. Longnecker at the National Institute of Environmental Health Sciences is working to validate toenail copper concentrations as a surrogate measure of exposure to copper. Development of high-performance liquid chromatography (HPLC) and derivatization techniques for identifying natural copper chelators in marine water is being conducted at Cornell University under the guidance of Drs. B.A. Ahner and J.W. Moffett. Dr. D.L. Sparks, at the University of Delaware, is developing x-ray absorption fine structure (XAFS) and atomic force microscopy (AFS) techniques for the study of metal/metalloid reactions in soil. Dr. J.F. Tyson and

7. ANALYTICAL METHODS

colleagues at the University of Massachusetts at Amherst are developing liquid-liquid extraction pretreatment techniques that can be interfaced with HPLC-ICP-MS instrumentation.

8. REGULATIONS AND ADVISORIES

ATSDR has derived acute- and intermediate-duration oral MRLs for copper. These MRLs are intended to protect against the health effects associated with exposure to copper-contaminated drinking water; it assumes that the affected population will have a normal intake of copper from the diet. The acute-duration oral MRL is 0.01 mg copper/kg/day. It is based on the occurrence of gastrointestinal disturbances in women ingesting 0.0731 mg Cu/kg/day in drinking water for 2 weeks; no adverse effects were observed at a drinking water dose of 0.0272 mg Cu/kg/day (Pizarro et al. 1999). To calculate an MRL, the NOAEL of 0.0272 mg Cu/kg/day was divided by an uncertainty factor of 3 to account for human variability.

An intermediate-duration oral MRL of 0.01 mg copper/kg/day was derived for copper. This MRL is based on the occurrence of gastrointestinal disturbances in men and women ingesting 0.091 mg Cu/kg/day in drinking water for 2 months; no adverse effects were observed at a drinking water dose of 0.042 mg Cu/kg/day (Araya et al. 2003b). To calculate an MRL, the NOAEL of 0.042 mg Cu/kg/day was divided by an uncertainty factor of 3 to account for human variability.

International, national, and state regulations and guidelines regarding human exposure to copper are summarized in Table 8-1.

8. REGULATIONS AND ADVISORIES

Table 8-1. Regulations and Guidelines Applicable to Copper

Agency	Description	Information	Reference
<u>INTERNATIONAL</u>			
Guidelines:			
IARC	Carcinogenicity classification Copper 8-hydroxyquinoline	Group 3 ^a	IARC 2002
<u>NATIONAL</u>			
Regulations and Guidelines:			
a. Air			
ACGIH	TLV (8-hour TWA) Fume (Cu) Dusts and mists (as Cu)	0.2 mg/m ³ 1.0 mg/m ³	ACGIH 2001
EPA	Serious health effects form ambient air exposure (Cu)		EPA 2002b 40CFR1910.1000
NIOSH	REL (10-hour TWA) Fume (as Cu) Dusts and mists (as Cu)	0.1 mg/m ³ 1.0 mg/m ³	NIOSH 2002
OSHA	IDLH Fume, dusts, and mists (as Cu) PEL (8-hour TWA) for general industry Fume (as Cu) Dusts and mists (as Cu) PEL (8-hour TWA) for construction industry Fume (as Cu) Dusts and mists (as Cu) PEL (8-hour TWA) for shipyard industry Fume (as Cu) Dusts and mists (as Cu)	100 mg/m ³ 0.1 mg/m ³ 1.0 mg/m ³ 0.1 mg/m ³ 1.0 mg/m ³ 0.1 mg/m ³ 1.0 mg/m ³ 0.1 mg/m ³ 1.0 mg/m ³	OSHA 2002c 29CFR1910.1000 OSHA 2002b 29CFR1926.55 OSHA 2002a 29CFR1915.1000
b. Water			
DOT	Marine pollutant (Cu metal powder and cupric sulfate)		DOT 2002 49 CFR172.101, Appendix B
EPA	Drinking water standard Action level (Cu) MCLG (Cu) Groundwater monitoring (Cu) Suggested method 6010 7210	1.3 mg/L 1.3 mg/L <u>PQL</u> 60 µg/L 200 µg/L	EPA 2002c EPA 2002d 40CFR141.51(b) EPA 2002g 40CFR264, Appendix IX

8. REGULATIONS AND ADVISORIES

Table 8-1. Regulations and Guidelines Applicable to Copper

Agency	Description	Information	Reference
<u>NATIONAL</u> (cont.)			
EPA	Hazardous substance in accordance with Section 311(b)(2)(A) of the Clean Water Act (cupric sulfate and cupric sulfate ammoniated)		EPA 2002j 40CFR116.4
	Reportable quantity of hazardous substance designated pursuant to Section 311 of the Clean Water Act		EPA 2002k 40CFR117.3
	Cupric sulfate	10 pounds	
	Cupric sulfate, ammoniated	100 pounds	
	Secondary MCL for public water systems (Cu)	1.0 mg/L	EPA 2002e 40CFR143.3
	Toxic pollutant designated pursuant to Section 307(a)(1) of the Federal Water Pollution Control Act and is subject to effluent limitations (Cu and compounds)		EPA 2002a 40CFR401.15
	Water quality criteria (Cu)		EPA 1999
	Fresh water		
	CMC	13.0 µg/L	
	CCC	9.0 µg/L	
	Salt water		
	CMC	4.8 µg/L	
	CCC	3.1 µg/L	
Human health for consumption of water and organism	1,300 µg/L		
Organoleptic effect criteria	1,000 µg/L		
c. Food and Drugs			
EPA	Exemption from requirement of a tolerance in meat, milk, poultry, eggs, fish, shellfish, and irrigated crops when it results from the use as an algacide, herbicide, and fungicide when used in accordance with good agricultural practices (Cu)		EPA 2002f 40CFR180.1021
FDA	Bottled water; allowable level (Cu)	1.0 mg/L	FDA 2001a 21CFR165.110
	Clinical chemistry test system; copper test system measures copper levels in plasma, serum, and urine	Exempt from premarket notification procedures in Subpart E of Part 807	FDA 2001b 21CFR862.1190
	Color additives exempt from certification—copper powder for use in externally applied drugs	Cu not less than 95%	FDA 2001e 21CFR73.1647

8. REGULATIONS AND ADVISORIES

Table 8-1. Regulations and Guidelines Applicable to Copper

Agency	Description	Information	Reference
<u>NATIONAL</u> (cont.)			
FDA	Color additives exempt from certification—copper powder for use in cosmetics		FDA 2001f 21CFR73.2647
	Direct food substance affirmed as generally recognized as safe when used as a nutrient supplement or as a processing aid (cupric sulfate)		FDA 2001c 21CFR184.1261
	Drug products containing certain active ingredients offered over-the-counter; inadequate data to establish general recognition of the safety and effectiveness of these ingredients for the specified uses (Cu)	Weight control drug product	FDA 2001g 21CFR310.545(a)(20)
	Trace minerals added to animal feeds as nutritional dietary supplements are generally recognized as safe when added at levels consistent with good feeding practices (Cu compounds)		FDA 2001i 21CFR582.80
IOM	Recommended dietary allowance (RDA)	0.9 mg/day	IOM 2001
	Tolerable upper intake level	10 mg/day	
d. Other			
EPA	Carcinogenicity classification (Cu)	Group D ^b	IRIS 2004
	RfC	No data	
	RfD	No data	
	Reportable quantity designated as a CERCLA hazardous substance under Section 307(a) of the Clean Water Act (Cu)	5,000 pounds	EPA 2002h 40CFR302.4
	Reportable quantity designated as a CERCLA hazardous substance under Section 311(b) (4) of the Clean Water Act (cupric sulfate)	10 pounds	EPA 2002h 40CFR302.4
	Toxic chemical release reporting; community right-to-know; effective date of reporting (Cu)	01/01/87	EPA 2002i 40CFR372.65(a)
<u>STATE</u>			
Regulations and Guidelines			
a. Air			
Illinois	Toxic air contaminant (Cu)		BNA 2001
Louisiana	Toxic air pollutant ^c		BNA 2001
	Minimum emission rate (Cu and compounds)	25 pounds/year	

8. REGULATIONS AND ADVISORIES

Table 8-1. Regulations and Guidelines Applicable to Copper

Agency	Description	Information	Reference
<u>STATE (cont.)</u>			
New Mexico	Toxic air pollutant		BNA 2001
	Fume (Cu)		
	OEL	0.2 mg/m ³	
	Emissions	0.0133 pounds/ hour	
	Dusts and mists (as Cu)		
	OEL	1.0 mg/m ³	
Vermont	Cu compounds		BNA 2001
	Hazardous ambient air standard	100 µg/m ³	
	Averaging time	8 hours	
	Action level	4 pounds/hour	
b. Water			
Arizona	Drinking water guideline (Cu)	1,300 µg/L	HSDB 2004
North Carolina	Groundwater quality standard (Cu)	1.0 mg/L	BNA 2001
c. Food			
No data			
d. Other			
Arizona	Soil remediation levels (Cu and compounds)		BNA 2001
	Residential	2,800 mg/kg	
	Non-residential	63,000 mg/kg	
Florida	Toxic substance in the workplace (Cu fume, dust, and mist)		BNA 2001

^aGroup 3: unclassifiable as to carcinogenicity to humans

^bGroup D: not classifiable as to human carcinogenicity

^cClass II: suspected human carcinogen and known or suspected human reproductive toxin

ACGIH = American Conference of Governmental Industrial Hygienists; BNA = Bureau of National Affairs; CERCLA = Comprehensive Environmental Response Compensation and Liability Act; CFR = Code of Federal Regulations; CCC = criterion continuous concentration; CMC = criteria maximum concentration; Cu = copper; DOT = Department of Transportation; EPA = Environmental Protection Agency; FDA = Food and Drug Administration; HSDB = Hazardous Substances Data Bank; IARC = International Agency for Research on Cancer; IDLH = immediately dangerous to life and health; IOM = Institute of Occupational Medicine; IRIS = Integrated Risk Information System; MCL = maximum contaminant level; MCLG = maximum contaminant level goal; NIOSH = National Institute for Occupational Safety and Health; OEL = occupational exposure limit; OSHA = Occupational Safety and Health Administration; PEL = permissible exposure limits; PQL = practical quantitation limits; RDA = recommended dietary allowance; REL = recommended exposure limit; RfC = inhalation reference concentration; RfD = oral reference dose; TLV = threshold limit value; TWA = time-weighted average

9. REFERENCES

- Aaseth J, Norseth T. 1986. Copper. In: Friberg L, Nordberg GF, Vouk V, eds. Handbook on the toxicology of metals 2. New York, NY: Elsevier Science Publishers, 233-254.
- *Aburto EM, Cribb AE, Fuentealba IC. 2001a. Effect of chronic exposure to excess dietary copper and dietary selenium supplementation on liver specimens from rats. *Am J Vet Res* 62(9):1423-1427.
- *Aburto EM, Cribb AE, Fuentealba IC, et al. 2001b. Morphological and biochemical assessment of the liver response to excess dietary copper in Fischer 344 rats. *Can J Vet Res* 65(2):97-103.
- Aburto EM, Cribb A, Fuentealba IC, et al. 2001c. The failure of selenium supplementation to prevent copper-induced liver damage in Fischer 344 rats. *Can J Vet Res* 65(2):104-110.
- *ACGIH. 1988. Threshold limit values and biological exposure indices for 1988-1989. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.
- *ACGIH. 2001. Threshold limit values for chemical substances and physical agents and biological exposure indices. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.
- Ackerman DJ, Reinecke AJ, Els HJ, et al. 1999. Sperm abnormalities associated with high copper levels in impala (*Aepyceros melampus*) in the Kruger National Park, South Africa. *Ecotoxicol Environ Saf* 43(3):261-266.
- Adachi A, Okiyasu M, Nishikawa A, et al. 1998. Metal levels in rain water from Kobe City in Japan. *Bull Environ Contam Toxicol* 60:892-897.
- Adamo P, Dudka S, Wilson MJ. 1996. Chemical and mineralogical forms of Cu and Ni in contaminated soils from the Sudbury mining and smelting region, Canada. *Environ Pollut* 91(1):11-19.
- *Adinolfi M. 1985. The development of the human blood-CSF-brain barrier. *Dev Med Child Neurol* 27:532-537.
- *Adlercreutz H. 1995. Phytoestrogens: Epidemiology and a possible role in cancer protection. *Environ Health Perspect Suppl* 103(7):103-112.
- Agarwal K, Sharma A, Talukder G. 1989. Effects of copper on mammalian cell components. *Chem Biol Interact* 69(1):1-16.
- *Agarwal K, Sharma A, Talukder G. 1990. Clastogenic effects of copper sulfate on the bone marrow chromosomes of mice in vivo. *Mutat Res* 243(1):1-6.
- *Agency for Toxic Substances and Disease Registry. 1989. Decision guide for identifying substance-specific data needs related to toxicological profiles; Notice. *Federal Register* 54(174):37618-37634.

* Cited in text

9. REFERENCES

*Agency for Toxic Substances and Disease Registry. 1990. Biomarkers of organ damage or dysfunction for the renal, hepatobiliary, and immune systems. Subcommittee on Biomarkers of Organ Damage and Dysfunction, Agency for Toxic Substances and Disease Registry, Atlanta, GA.

Aggett PJ, Fairweather-Tait S. 1998. Adaptation to high and low copper intakes: Its relevance to estimated safe and adequate daily dietary intakes. *Am J Clin Nutr* 67:1061S-1063S.

*Ahasan HAMN, Chowdhury MAJ, Azhar MA, et al. 1994. Copper sulphate poisoning. *Trop Doct* 24(2):52-53.

Ahmad MS, Fazal F, Rahman A, et al. 1992. Activities of flavonoids for the cleavage of DNA in the presence of Cu(II) correlation with generation of active oxygen species. *Carcinogenesis* 13(4):605-608.

Ahmed HM, Shoka AA. 1994. Toxic interactions between copper sulphate and some organic agrochemicals. *Toxicol Lett* 70(1):109-119.

Ahmed KO, Al-Swaidan HM, Davies BE. 1993. Simultaneous elemental analysis in dust of the city of Riyadh, Saudi Arabia by inductively coupled plasma-mass spectrometry (ICP/MS). *Sci Total Environ* 138:207-212.

Aisen P, Morell AG, Alpert S, et al. 1964. Biliary excretion of caeruloplasmin copper. *Nature* 203:873-874.

*Akintonwa A, Mabadeje AFB, Odutola TA. 1989. Fatal poisonings by copper sulfate ingested from "spiritual water". *Vet Hum Toxicol* 31(5):453-454.

Algerwie MH, Khatri PC. 1998. Serum copper in newborns and their mothers. *Indian J Pediatr* 65(6):899-903.

Alkhatib E, Castor K. 2000. Parameters influencing sediments resuspension and the link to sorption of inorganic compounds. *Environ Monit Assess* 65(3):531-546.

*Allen SK, Allen JM, Lucas S. 1996. Dissolved metal concentrations in surface waters from west-central Indiana contaminated with acidic mine drainage. *Bull Environ Contam Toxicol* 56:240-243.

*Allen-Gil SM, Gubala CP, Landers DH, et al. 1997. Heavy metal accumulation in sediment and freshwater fish in U.S. arctic lakes. *Environ Toxicol Chem* 16(4):733-741.

*Altman PL, Dittmer DS. 1974. Biological handbooks: Biology data book. Vol. III. 2nd ed. Bethesda, MD: Federation of American Societies for Experimental Biology, 1987-2008, 2041.

*American Institute of Nutrition. 1977. Report of the AID ad hoc committee on standards for nutritional studies. *J Nutr* 107:1340-1348.

*American Society for Testing and Materials. 1990. Standard test methods for copper in water. ASTM D 1688, Verlag.

*American Society for Testing and Materials. 2000. Standard test method for elements in water by inductively coupled plasma - mass spectrometry. ASTM D 5673-96, Verlag.

9. REFERENCES

- Amrhein C, Mosher PA, Strong JE, et al. 1994. Trace metal solubility in soils and waters receiving deicing salts. *J Environ Qual* 23(2):219-227.
- *Amrhein C, Strong JE, Mosher PA. 1992. Effect of deicing salts on metal and organic matter mobilization in roadside soils. *Environ Sci Technol* 26(4):703-709.
- *Andersen ME, Krishnan K. 1994. Relating in vitro to in vivo exposures with physiologically based tissue dosimetry and tissue response models. In: Salem H, ed. *Animal test alternatives: Refinement, reduction, replacement*. New York, NY: Marcel Dekker, Inc., 9-25.
- *Andersen ME, Clewell HJ III, Gargas ML, et al. 1987. Physiologically based pharmacokinetics and the risk assessment process for methylene chloride. *Toxicol Appl Pharmacol* 87:185-205.
- *Anderson JR, Aggett FJ, Buseck PR, et al. 1988. Chemistry of individual aerosol particles from Chandler, Arizona, an arid urban environment. *Environ Sci Technol* 22(7):811-818.
- Andrzejak R, Antonowicz J, Tomczyk J, et al. 1993. Lead and cadmium concentrations in blood of people living near a copper smelter in Legnica, Poland. *Sci Total Environ Suppl*:233-236.
- Apgar GA, Kornegay ET. 1996. Mineral balance of finishing pigs fed copper sulfate or a copper-lysine complex at growth-stimulating levels. *J Anim Sci* 74:1594-1600.
- Apte SC, Gardner MJ, Ravenscroft JE. 1990. An investigation of copper complexation in the Severn Estuary using differential pulse cathodic stripping voltammetry. *Mar Chem* 29:63-75.
- Araki S, Murata K, Uchida E, et al. 1993. Radial and median nerve conduction velocities in workers exposed to lead, copper, and zinc: A follow-up study for 2 years. *Environ Res* 61(2):308-316.
- *Araki S, Sata F, Murata K. 1990. Adjustment for urinary flow rate: an improved approach to biological monitoring. *Int Arch Occup Environ Health* 62(6):471-477.
- *Araya M, Chen B, Klevay LM, et al. 2003a. Confirmation of an acute no-observed-adverse-effect and low-observed-adverse-effect level for copper in bottled drinking water in a multi-site international study. *Reg Tox Pharmacol* 38:389-399.
- *Araya M, McGoldrick MC, Klevay LM, et al. 2001. Determination of an acute no-observed-adverse-effect level (NOAEL) for copper in water. *Regul Toxicol Pharmacol* 34(2):137-148.
- *Araya M, Olivares M, Pizarro F, et al. 2003b. Gastrointestinal symptoms and blood indicators of copper load in apparently healthy adults undergoing controlled copper exposure. *Am J Clin Nutr* 77(3):646-650.
- *Araya M, Pena C, Pizarro F, et al. 2003c. Gastric response to acute copper exposure. *Sci Total Environ* 303(3):253-257.
- *Armstrong CW, Moore LW, Hackler RL, et al. 1983. An outbreak of metal fume fever. Diagnostic use of urinary copper and zinc determinations. *J Occup Med* 25:886-888.
- *Arnaud J, Favier A. 1995. Copper, iron, manganese and zinc contents in human colostrum and transitory milk of French women. *Sci Total Environ* 159:9-15.

9. REFERENCES

- *Arredondo M, Uauy R, Gonzalez M. 2000. Regulation of copper uptake and transport in intestinal cell monolayers by acute and chronic copper exposure. *Biochim Biophys Acta* 1474(2):169-176.
- *Askergren A, Mellgren M. 1975. Changes in the nasal mucosa after exposure to copper salt dust. A preliminary report. *Scand J Work Environ Health* 1:45-49.
- Aston N, Morris P, Tanner S. 1996. Retrosine in breast milk influences copper handling in suckling rat pups. *J Hepatol* 25(5):748-755.
- Aston NS, Morris PA, Tanner MS, et al. 1998. An animal model for copper-associated cirrhosis in infancy. *J Pathol* 186(2):215-221.
- Aston NS, Watt N, Tanner MS, et al. 2000. Copper toxicity affects proliferation and viability of human hepatoma cells (HepG2 line). *Hum Exp Toxicol* 19:367-376.
- August D, Janghorbani M, Young VR. 1989. Determination of zinc and copper absorption at three dietary Zn-Cu ratios by using stable isotope methods in young adult and elderly subjects. *Am J Clin Nutr* 50:1457-1463.
- *Aulenbach DB, Meyer MA, Beckwith E, et al. 1987. Removal of heavy metals in POTW. *Environ Progress* 6:91-98.
- *Aulerich RJ, Ringer RK, Bleavins MR, et al. 1982. Effects of supplemental dietary copper on growth, reproductive performance and kit survival of standard dark mink and the acute toxicity of copper to mink. *J Anim Sci* 55(2):337-343.
- *Badri MA, Aston SR. 1983. Observations on heavy metal geochemical associations in polluted and non-polluted estuarine sediments. *Environ Pollut Ser B* 6:181-193.
- Baker DH, Odle J, Funk MA, et al. 1991. Research note: Bioavailability of copper in cupric oxide, cuprous oxide and in copper-lysine complex. *Poult Sci* 70:177-179.
- *Balogh S. 1996. The fate of metals in sewage sludge incinerators. *Water Air Soil Pollut* 91:249-254.
- Baranowska I, Czernicki K, Aleksandrowicz R. 1995. The analysis of lead, cadmium, zinc, copper and nickel content in human bones from the Upper Silesian industrial district. *Sci Total Environ* 159:155-162.
- Baranowska-Dutkiewicz B, Dutkiewicz T. 1991. Evaluation of simultaneous industrial and environmental exposure to metals. *Sci Total Environ* 101:149-151.
- Barash A, Shoham Z, Borenstein R, et al. 1990. Development of human embryos in the presence of a copper intrauterine device. *Gynecol Obstet Invest* 29(3):203-206.
- Barceloux DG. 1999. Copper. *J Toxicol Clin Toxicol* 37(2):217-230.
- Bargagli R, Barghigiani C, Siegel BZ, et al. 1991. Trace metal anomalies in surface soils and vegetation on two active island volcanos: Stromboli and Vulcano (Italy). *Sci Total Environ* 102:209-222.
- Bargagli R, Cateni D, Nelli L, et al. 1997. Environmental impact of trace element emissions from geothermal power plants. *Arch Environ Contam Toxicol* 33:172-181.

9. REFERENCES

Barnea A, Cho G. 1984. Evidence that copper-amino acid complexes are potent stimulators of the release of luteinizing hormone-releasing hormone from isolated hypothalamic granules. *Endocrinology* 115:936-943.

*Barnes DG, Dourson M. 1988. Reference dose (RfD): Description and use in health risk assessments. *Regul Toxicol Pharmacol* 8:471-486.

*Barranco VP. 1972. Eczematous dermatitis caused by internal exposure to copper. *Arch Dermatol* 106:386-387.

*Barrie LA, Lindberg SE, Chan WH, et al. 1987. On the concentration of trace metals in precipitation. *Atmos Environ* 21:1133-1135.

*Batsura YD. 1969. Electron microscope investigation of penetration of copper oxide aerosol from the lungs into the blood and internal organs. *Byull Eksp Biol Med* 68:1175-1178.

Batzevich VA. 1995. Hair trace element analysis in human ecology studies. *Sci Total Environ* 164:89-98.

Bauer G, Schachermayer E. 1996. Statistical analysis of heavy metal data from municipal waste incineration residues. *Environ Sci Pollut Res* 3(1):10-16.

*Bearn AG, Kunkel HG. 1955. Metabolic studies in Wilson's disease using Cu^{64} . *J Lab Clin Med* 45:623-631.

Beck JN, Sneddon J. 2000. Metal concentrations in soils and sediments in Southwest Louisiana. *Anal Lett* 33(10):1913-1959.

Beliaeff B, O'Connor TP, Daskalakis DK, et al. 1997. U.S. Mussel watch data from 1986 to 1994: Temporal trend detection at large spatial scales. *Environ Sci Technol* 31:1411-1415.

Beltran-Garcia MJ, Espinosa A, Herrera N, et al. 2000. Formation of copper oxychloride and reactive oxygen species as causes of uterine injury during copper oxidation of Cu-IUD. *Contraception* 61(2):99-103.

Benders AA, Li J, Lock RA, et al. 1994. Copper toxicity in cultured human skeletal muscle cells: the involvement of $\text{Na}^+/\text{Ca}^{2+}$ -ATPase and the $\text{Na}^+/\text{Ca}^{2+}$ -exchanger. *Pflugers Arch* 428(5-6):461-467.

Benedetti MF, Milne CJ, Kinniburgh DG, et al. 1995. Metal ion binding to humic substances: application of the non-ideal competitive adsorption model. *Environ Sci Technol* 29:446-457.

*Bentur Y, Koren G, McGuigan M, et al. 1988. An unusual skin exposure to copper; clinical and pharmacokinetic evaluation. *J Toxicol Clin Toxicol* 26(5-6):371-380.

Berg G, Kohlmeier L, Brenner H. 1998. Effect of oral contraceptive progestins on serum copper concentration. *Eur J Clin Nutr* 52:711-715.

*Berger GS. 1994. Epidemiology of endometriosis. In: Berger GS, ed. *Endometriosis: Advanced management and surgical techniques*. New York, NY: Springer-Verlag.

9. REFERENCES

- *Beveridge SJ, Boettcher B, Walker WR, et al. 1984. Biodistribution of ^{64}Cu in rats after topical application of two lipophilic anti-inflammatory Cu(II) formulations. *Agents Actions* 14(2):291-295.
- *Beyer WN, Cromartie EJ. 1987. A survey of Pb, Cu, Zn, Cd, As, and Se in earthworms and soil from diverse sites. *Environ Monit Assess* 8:27-36.
- Bhandari P, Andrews PLR. 1991a. Erratum: Preliminary evidence for the involvement of the putative 5-HT₄ receptor in zacopride-and copper sulphate-induced vomiting in the ferret. *Eur J Pharmacol* 211(3):430.
- Bhandari P, Andrews PLR. 1991b. Preliminary evidence for the involvement of the putative-5-HT₄ receptor in zacopride-and copper sulphate-induced vomiting in the ferret. *Eur J Pharmacol* 211(3):273-280.
- *Bhave SA, Pandi AN, Pradhan AM, et al. 1982. Liver disease in India. *Arch Dis Child* 57:922-928.
- *Bhave SA, Panditschein AN, Tanner MS. 1987. Comparison of feeding history of children with Indian childhood cirrhosis and paired controls. *J Pediatr Gastroenterol Nutr* 6:562-567.
- *Bhunya SP, Jena GB. 1996. Clastogenic effect of copper sulphate in chick in vivo test system. *Mutat Res* 367(2):57-63.
- *Bhunya SP, Pati PC. 1987. Genotoxicity of an inorganic pesticide, copper sulphate in mouse in vivo test system. *Cytologia* 52:801-808.
- Bingham MJ, McArdle HJ. 1994. A comparison of copper uptake by liver plasma membrane vesicles and uptake by isolated cultured rat hepatocytes. *Hepatology* 20(4):1024-1031.
- Bires J, Kovac G, Vrzgula L. 1991a. Interactions between copper and selenium in sheep in the course of experimentally-produced copper intoxication. *Vet Hum Toxicol* 33(5):489-491.
- Bires J, Kovac G, Vrzgula L. 1991b. Mineral profile of serum in experimental copper intoxication of sheep from industrial emissions. *Vet Hum Toxicol* 33(5):431-435.
- *Blanuša M, Ivicic N, Simeon V. 1990. Lead, iron, copper, zinc and ash in deciduous teeth in relation to age and distance from a lead smelter. *Bull Environ Contam Toxicol* 45(4):478-485.
- Blanuša M, Prester L, Matek M, et al. 1999. Trace elements in soil and coniferous needles. *Bull Environ Contam Toxicol* 62:700-707.
- *Blevins RD, Pancorbo OC. 1986. Metal concentrations in muscle of fish from aquatic systems in east Tennessee. *Water Air Soil Pollut* 29:361-371.
- Blincoe C. 1992. Simulation of copper metabolism by mammals. *Comput Biol Med* 22(1-2):113-122.
- *BNA. 2001. Environmental and Safety Library on the Web. States and Territories. Washington, DC. Bureau of National Affairs Inc. <http://www.esweb.bna.com/>.
- *Bopp RF, Simpson HJ, Chillrud SN, et al. 1993. Sediment-derived chronologies of persistent contaminants in Jamaica Bay, New York. *Estuaries* 16(3B):608-616.

9. REFERENCES

- *Borak J, Cohen H, Hethmon TA. 2000. Copper exposure and metal fume fever: Lack of evidence for a causal relationship. *Am Ind Hyg Assoc J* 61(6):832-836.
- Borga P, Elowson T, Liukko K. 1996. Environmental loads from water-sprinkled softwood timber. 1. Characteristics of an open and a recycling water system. *Environ Toxicol Chem* 15(6):856-867.
- *Boyden R, Potter VR, Elvehjem CA. 1938. Effect of feeding high levels of copper to albino rats. *J Nutr* 15:397-402.
- *Bradley RW, Morris JR. 1986. Heavy metals in fish from a series of metal-contaminated lakes near Sudbury, Ontario. *Water Air Soil Pollut* 27:341-354.
- Braune B, Muir D, DeMarch B, et al. 1999. Spatial and temporal trends of contaminants in Canadian Arctic freshwater and terrestrial ecosystems: A review. *Sci Total Environ* 230:145-207.
- *Breault RF, Colman JA, Aiken GR, et al. 1996. Copper speciation and binding by organic matter in copper-contaminated streamwater. *Environ Sci Technol* 30:3477-3486.
- Bremner I. 1998. Manifestations of copper excess. *Am J Clin Nutr* 67:1069S-1073S.
- *Breslin VT. 1999. Retention of metals in agricultural soils after amending with MSW and MSW-biosolids compost. *Water Air Soil Pollut* 109:163-178.
- *Brewer GJ. 1995. Interactions of zinc and molybdenum with copper in therapy of Wilson's disease. *Nutrition* 11:114-116.
- Brewer GJ. 1998. Wilson disease and canine copper toxicosis. *Am J Clin Nutr* 67:1087S-1090S.
- *Brewer GJ, Yuzbasiyan-Gurkan V. 1992. Wilson disease. *Medicine* 71(3):139-164.
- *Brewer GJ, Dick RD, Schall W, et al. 1992. Use of zinc acetate to treat copper toxicosis in dogs. *J Am Vet Med Assoc* 201(4):564-568.
- *Brewer GJ, Yuzbasiyan-Gurkan V, Johnson V, et al. 1993. Treatment of Wilson's disease with zinc: XI: Interaction with other anticopper agents. *J Am Coll Nutr* 12(1):26-30.
- *Brewer GJ, Yuzbasiyan-Gurkan V, Lee D-Y, et al. 1989. Treatment of Wilson's disease with zinc. VI. Initial treatment studies. *J Lab Clin Med* 114(6):633-638.
- Brown KR, McPherson RG. 1992. Concentrations of copper, zinc and lead in the Sydney rock oyster, *Saccostrea commercialis* (Iredale and Roughley) from the Georges River, New South Wales. *Sci Total Environ* 12:27-33.
- *Brown KW, Thomas JC, Slowey JF. 1983. The movement of metals applied to soils in sewage effluent. *Water Air Soil Pollut* 19:43-54.
- *Bruce BW, McMahon PB. 1996. Shallow ground-water quality beneath a major urban center: Denver, Colorado, USA. *J Hydrol* 186:129-151.
- *Buchanan SD, Diseker RA III, Sinks T, et al. 1999. Copper in drinking water, Nebraska, 1994. *Int J Occup Environ Health* 5(4):256-261.

9. REFERENCES

- *Buchholz BA, Landsberger S. 1995. Leaching dynamics studies of municipal solid waste incinerator ash. *J Air Waste Manage Assoc* 45:579-590.
- Buckley DE, Smith JN, Winters GV. 1995. Accumulation of contaminant metals in marine sediments of Halifax Harbour, Nova Scotia: Environmental factors and historical trends. *Appl Geochem* 10:175-195.
- *Budavari S, O'Neil MJ, Smith A, et al., eds. 2001. *The Merck index: an encyclopedia of chemicals, drugs and biologicals*. Whitehouse Station, NJ: Merck & Co. Inc., 440, 462.
- Bu-Olayan AH, Subrahmanyam MNV. 1996. Trace metals in fish from the Kuwait coast using the microwave acid digestion technique. *Env Int* 22(6):753-758.
- Burba P, Rocha J, Klockow D. 1994. Labile complexes of trace metals in aquatic humic substances: Investigations by means of an ion exchange-based flow procedure. *Fresenius J Anal Chem* 349:800-807.
- Burkitt MJ. 1994. Copper-DNA adducts. *Methods Enzymol* 234:66-79.
- *Bush JA, Mahoney JP, Markowitz H, et al. 1955. Studies on copper metabolism. XVI. Radioactive copper studies in normal subjects and in patients with hepatolenticular degeneration. *J Clin Invest* 34:1766-1778.
- *Butterman WC. 1982. Copper. In: *Bureau of Mines Minerals Yearbook*. Pittsburgh, PA: U.S. Department of the Interior, 279-285.
- Byczkowski JZ, Gearhart JM, Fisher JW. 1994. "Occupational" exposure of infants to toxic chemicals via breast milk. *Nutrition* 10(1):43-48.
- *Cadle SH, Mulawa PA, Hunsanger EC, et al. 1999. Composition of light-duty motor vehicle exhaust particulate matter in the Denver, Colorado area. *Environ Sci Technol* 33:2328-2339.
- *Calabrese EJ, Moore GS. 1979. Can elevated levels of copper in drinking water precipitate acute hemolysis in G-6-PD deficient individuals. *Med Hypotheses* 5:493-498.
- *Camakaris J, Voskoboinik I, Mercer JF. 1999. Molecular mechanisms of copper homeostasis. *Biochem Biophys Res Commun* 261(2):225-232.
- Camner P, Johansson A. 1992. Reaction of alveolar macrophages to inhaled metal aerosols. *Environ Health Perspect* 97:185-188.
- *Campbell KR. 1994. Concentrations of heavy metals associated with urban runoff in fish living in stormwater treatment ponds. *Arch Environ Contam Toxicol* 27:352-356.
- Camusso M, Vigano L, Balestrini R. 1995. Bioconcentration of trace metals in rainbow trout: A field study. *Ecotoxicol Environ Saf* 31:133-141.
- *Cao ZH, Hu ZY. 2000. Copper contamination in paddy soils irrigated with wastewater. *Chemosphere* 41:3-6.
- *Cao Y, Conklin M, Betterton E. 1995. Competitive complexation of trace metals with dissolved humic acid. *Environ Health Perspect Suppl* 103(Suppl 1):29-32.

9. REFERENCES

- *Capar SG, Cunningham WC. 2000. Element and radionuclide concentrations in food: FDA total diet study 1991-1996. *J AOAC Int* 83(1):157-177.
- Carmalt JL, Baptiste KE, Blakley B. 2001. Suspect copper toxicity in an alpaca. *Can Vet J* 42:554-556.
- Carpenter TO, Pendrak ML, Anast CS. 1988. Metabolism of 25-hydroxyvitamin D in copper-laden rat: a model of Wilson's disease. *Am J Physiol* 254(2 Pt 1):E150-E154.
- *Cartwright GE, Wintrobe MM. 1964. Copper metabolism in normal subjects. *Am J Clin Nutr* 14:224-232.
- Castillo RO, Thaler MM, O'Toole C, et al. 1990. Hepatic copper metabolism in a mouse model for Menkes' Kinky Hair Syndrome. *Pediatr Res* 27(5):492-496.
- Catsiki VA, Bei F. 1992. Determination of trace metals in benthic organisms from an unpolluted area: Cyclades Islands (Aegean Sea). *Fresenius Environ Bull* 1(Suppl):S60-S65.
- Catsiki VA, Papathanassiou E, Bei F. 1991. Heavy metal levels in characteristic benthic flora and fauna in the Central Aegean Sea. *Mar Pollut Bull* 13:566-569.
- *CEIDARS. 2000. Chemical speciation. California Emission Inventory and Reporting System. <http://www.arb.ca.gov/emisinv/speciate/speciate.htm>.
- Chaim W, Mazor M. 1992. Pregnancy with an intrauterine device in situ and preterm delivery. *Arch Gynecol Obstet* 252(1):21-24.
- *Chan WH, Tang AJS, Chung DHS, et al. 1986. Concentration and deposition of trace metals in Ontario - 1982. *Water Air Soil Pollut* 29:373-389.
- Chang CC, Tatum HJ. 1970. A study of the antifertility effect of intrauterine copper. *Contraception* 1:265-270.
- Chang CC, Tatum HJ, Kincl FA. 1970. The effect of intrauterine copper and other metals on implantation in rats and hamsters. *Fertil Steril* 21:274-278.
- Chang SI, Reinfelder JR. 2000. Bioaccumulation, subcellular distribution, and trophic transfer of copper in a coastal marine diatom. *Environ Sci Technol* 34:4931-4935.
- Charlesworth SM, Lees JA. 1999. Particulate-associated heavy metals in the urban environment: their transport from source to deposit, Coventry, UK. *Chemosphere* 39(5):833-848.
- *Chattopadhyay A, Sarkar M, Sengupta R, et al. 1999. Antitesticular effect of copper chloride in albino rats. *J Toxicol Sci* 24(5):393-397.
- *Chen M, Ma LQ, Harris WG. 1999. Baseline concentrations of 15 trace elements in Florida surface soils. *J Environ Qual* 28(4):1173-1181.
- *Chen R, Wei L, Chen R-L. 1995. Lung cancer mortality update and prevalence of smoking among copper miners and smelters. *Scand J Work Environ Health* 21:513-516.

9. REFERENCES

- *Chen R, Wei L, Huang H. 1993. Mortality from lung cancer among copper miners. *Br J Ind Med* 50(6):505-509.
- Chowrimootoo GF, Ahmed HA, Seymour CA. 1996. New insights into the pathogenesis of copper toxicosis in Wilson's disease: evidence for copper incorporation and defective canalicular transport of caeruloplasmin. *Biochem J* 315(Pt 3):851-855.
- *Christensen TH, Kjeldsen P, Albrechtsen HJ, et al. 1994. Attenuation of landfill leachate pollutants in aquifers. *Crit Rev Environ Sci* 24:119-202.
- *Chugh KS, Sakhuja V. 1979. Acute copper intoxication. *Int J Artif Organs* 2(4):181-182.
- Chukwuma C. 1994. Contamination of soils and rice by heavy metals in the Enyigba-Abakaliki lead and zinc mine, Nigeria. *Toxicol Environ Chem* 41:125-130.
- *Chuttani HK, Gupta PS, Gulati S, et al. 1965. Acute copper sulfate poisoning. *Am J Med* 39:849-854.
- Clark DR, Bickham JW, Baker DL, et al. 2000. Environmental contaminants in Texas, USA, wetland reptiles: evaluation using blood samples. *Environ Toxicol Chem* 19(9):2259-2265.
- *Clemens S. 2001. Review: Molecular mechanisms of plant metal tolerance and homeostasis. 212:475-486.
- *Clewell HJ III, Andersen ME. 1985. Risk assessment extrapolations and physiological modeling. *Toxicol Ind Health* 1(4):111-131.
- Coakley JP, Serodes JB. 1994. Spatial and vertical trends in sediment-phase contaminants in the upper estuary of the St. Lawrence River. *Estuaries* 16(3B):653-669.
- *Coale KH, Bruland KW. 1988. Copper complexation in the Northeast Pacific. *Limnol Oceanogr* 33:1084-1101.
- Cohen HJ, Powers BJ. 1994. A study of respirable versus nonrespirable copper and zinc oxide exposures at a nonferrous foundry. *Am Ind Hyg Assoc J* 55(11):1047-1059.
- Cohen HJ, Powers BJ. 2000. Particle size characterizations of copper and zinc oxide exposures of employees working in a nonferrous foundry using cascade impactors. *Am Ind Hyg Assoc J* 61(3):422-430.
- Cohen JM, Kamphake LJ, Harris EK, et al. 1960. Taste threshold concentrations of metals in drinking water. *J Am Water Works Assoc* 52:660-661.
- *Colborn T, Clement C, eds. 1992. Chemically-induced alterations in sexual and functional development: The wildlife human connection. In: *Advances in modern environmental toxicology* Vol. XXI: Princeton, NJ: Princeton Scientific Publishing.
- *Cole KL, Engstrom DR, Futyma RP, et al. 1990. Past atmospheric deposition of metals in Northern Indiana measured in a peat core from Cowles Bog. *Environ Sci Technol* 24:543-549.
- Cole RH, Frederick RE, Healy RP, et al. 1984. Preliminary findings of the priority pollutant monitoring project of the National Urban Runoff Program. *J Water Pollut Control Fed* 56:898-908.

9. REFERENCES

- *Coleman ME, Elder RS, Basu P. 1992. Trace metals in edible tissues of livestock and poultry. *J AOAC Int* 75(4):615-625.
- Comber SDW, Gardner MJ, Gunn AM, et al. 1996. Kinetics of trace metal sorption to estuarine. *Chemosphere* 33(6):1027-1040.
- *Cordano A. 1998. Clinical manifestations of nutritional copper deficiency in infants and children. *Am J Clin Nutr* 67:1012S-1016S.
- *Cotton FA, Wilkinson G. 1980. Copper. *Advanced inorganic chemistry*. New York, NY: John Wiley and Sons, 798-821.
- Couillard D, Chartier M, Mercier G. 1994. Major factors influencing bacterial leaching of heavy metals (Cu and Zn) from anaerobic sludge. *Environ Pollut* 85:175-184.
- Cox DW. 1999. Disorders of copper transport. *Br Med Bull* 55(3):544-555.
- *Crampton RF, Matthews DM, Poisner R. 1965. Observation on the mechanism of absorption of copper by the small intestine. *J Physiol* 178:111-126.
- *Crawford DW, Bonnevie NL, Wenning RJ. 1995. Sources of pollution and sediment contamination in Newark Bay, New Jersey. *Ecotoxicol Environ Saf* 30:85-100.
- *CRIS. 2003. CRIS Database. Current Research Information System.
- *Cristofori P, Terron A, Marella M, et al. 1992. Copper supplementation in the rat: Preliminary observations on the clinical, hematological and histopathological profile. *Agents Actions* 108:C118-C120.
- Cromwell GL, Lindemann MD, Monegue HJ, et al. 1998. Tribasic copper chloride and copper sulfate as copper sources for weanling pigs. *J Anim Sci* 76:118-123.
- Cunningham WC, Stroube WB Jr. 1987. Application of an instrumental neutron activation analysis procedure to analysis of food. *Sci Total Environ* 63:29-43.
- Cuzzocrea S, Persichini T, Dugo L, et al. 2003. Copper induces type II nitric oxide synthase in vivo. *Free Radic Biol Med* 34(10):1253-1262.
- *Cyr F, Mehra MC, Mallet VN. 1987. Leaching of chemical contaminants from a municipal landfill site. *Bull Environ Contam Toxicol* 38:775-782.
- *Dameron CT, Harrison MD. 1998. Mechanisms for protection against copper toxicity. *Am J Clin Nutr* 67(5):1091S-1097S.
- *Danks DM. 1988. Copper deficiency in humans. *Annu Rev Nutr* 8:235-257.
- Darwish HM, Cheney JC, Schmitt RC, et al. 1984. Mobilization of copper(II) from plasma components and mechanism of hepatic copper transport. *Am J Physiol* 246:G72-79.

9. REFERENCES

- *Daskalakis KD, O'Connor TP. 1995. Distribution of chemical concentrations in US coastal and estuarine sediment. *Mar Environ Res* 40:381-398.
- Dassel de Vergara J, Zietz B, Schneider HB, et al. 1999. Determination of the extent of excessive copper concentrations in the tap-water of households with copper pipes and an assessment of possible health hazards for infants. *Eur J Med Res* 4(11):475-482.
- *Davidson CI, Goold WD, Mathison TP, et al. 1985. Airborne trace elements in Great Smoky Mountains, Olympic, and Glacier National Parks. *Environ Sci Technol* 19:27-34.
- Davidson LA, McOrmond SL, Harris ED. 1994. Characterization of a particulate pathway for copper in K562 cells. *Biochim Biophys Acta* 1221:1-6.
- *Davies DJA, Bennett BG. 1985. Exposure of man to environmental copper - an exposure commitment. *Sci Total Environ* 46:215-227.
- *Davies NT, Campbell JK. 1977. The effect of cadmium on intestinal copper absorption and binding in the rat. *Life Sci* 20:955-960.
- Davies NT, Nightingale R. 1975. The effects of phytate on intestinal absorption and secretion of zinc, and whole-body retention of Zn, copper, iron and manganese in rats. *Br J Nutr* 34:243-258.
- *Davies-Colley RJ, Nelson PO, Williamson KJ. 1984. Copper and cadmium uptake by estuarine sedimentary phases. *Environ Sci Technol* 18:491-499.
- *Davies-Colley RJ, Nelson PO, Williamson KJ. 1985. Sulfide control of cadmium and copper concentrations in anaerobic estuarine sediments. *Mar Chem* 16:173-186.
- *Davis AP, Shokouhian M, Ni S. 2001. Loading estimates of lead, copper, cadmium, and zinc in urban runoff from specific sources. *Chemosphere* 44:997-1009.
- *Davis CD, Johnson WT. 2002. Dietary copper affects azoxymethane-induced intestinal tumor formation and protein kinase C isozyme protein and mRNA expression. *J Nutr* 132:1018-1025.
- Davis GK, Mertz W. 1987. Copper. Mertz W eds. *Trace elements in human and animal nutrition*: San Diego: Academic Press, Inc., 301-364.
- *Dean JA, ed. 1985. *Lange's handbook of chemistry*. New York, NY: McGraw-Hill Book Co., 10-11.
- De Gregori I, Pinochet H, Delgado D, et al. 1994. Heavy metals in bivalve mussels and their habitats from different sites along the Chilean coast. *Bull Environ Contam Toxicol* 52:261-268.
- *Demerec M, Bertani G, Flint J. 1951. A survey of chemicals for mutagenic action on E coli. *Am Nat* 85:119-136.
- *De Vries DJ, Sewell RB, Beart PM. 1986. Effects of copper on dopaminergic function in the rat corpus striatum. *Exp Neurol* 91:546-558.
- Diaz-Barriga F, Santos MA, Mejia JDJ, et al. 1993. Arsenic and cadmium exposure in children living near a smelter complex in San Luis Potosi, Mexico. *Environ Res* 62(2):242-250.

9. REFERENCES

- Dicarlo FJ. 1979. Copper-induced heart malformations in hamsters. *Experientia* 35:827-828.
- Dicarlo FJ. 1980. Syndromes of cardiovascular malformations induced by copper citrate in hamsters. *Teratology* 21:89-101.
- *Diks DM, Allen HE. 1983. Correlation of copper distribution in a freshwater-sediment system to bioavailability. *Bull Environ Contam Toxicol* 30:37-43.
- Di Toro DM, Allens HE, Bergman HL, et al. 2001. Hazard/risk assessment biotic ligand model of the acute toxicity of metals. 1. Technical basis. *Environ Toxicol Chem* 20(10):2383-2396.
- DOI. 1982. Copper. Bureau of mines minerals yearbook: Pittsburgh, PA: Department of the Interior. 279-285.
- *Domergue FL, Védry JC. 1992. Mobility of heavy metals in soil profiles. *Int J Environ Anal Chem* 46:13-23.
- *Domingo JL, Gomez M, Jones MM. 2000. Comparative efficacy of several potential treatments for copper mobilization in copper-overloaded rats. *Biol Trace Elem Res* 74(2):127-139.
- Donley, SA, Ilagan BJ, Rim H, et al. 2002. Copper transport to mammary gland and milk during lactation in rats. *Am J Physiol Endocrinol Metab* 283(4):E667-E675.
- *Dörner K, Dziadzka S, Höhn A, et al. 1989. Longitudinal manganese and copper balances in young infants and preterm infants fed on breast-milk and adapted cow's milk formulas. *Br J Nutr* 61:559-572.
- *DOT. 2002. List of marine pollutants. Hazardous materials regulations and procedures. U.S. Department of Transportation. Code of Federal Regulations. <http://63.141.231.97/cgi-bin/om>. April 10, 2002.
- *Dressler RL, Storm GL, Tzilkowski WM, et al. 1986. Heavy metals in cottontail rabbits on mined lands treated with sewage sludge. *J Environ Qual* 15(3):278-281.
- *Drummond JG, Aranyi C, Schiff LJ, et al. 1986. Comparative study of various methods used for determining health effects of inhaled sulfates. *Environ Res* 41:514-528.
- *Duby P. 1980. Extractive metallurgy. Kirk-Othmer encyclopedia of chemical technology. New York, NY: John Wiley and Sons, 739-767.
- Dudka S, Ponce-Hernandez R, Tate G, et al. 1996. Forms of Cu, Ni, and Zn in soils of Sudbury, Ontario and the metal concentrations in plants. *Water Air Soil Pollut* 90:531-542.
- Dunn MA, Green MH, Leach RM. 1991. Kinetics of copper metabolism in rats: a compartmental model. *Am J Physiol* 261:E115-125.
- Dusing DC, Bishop PL, Keener TC. 1992. Effect of redox potential on leaching from stabilized/solidified waste materials. *J Air Waste Manage Assoc* 42:56-62.
- *Eary LE, Rai D, Mattigod SV, et al. 1990. Geochemical factors controlling the mobilization of inorganic constituents from fossil fuel combustion residues: II. Review of the minor elements. *J Environ Qual* 19:202-214.

9. REFERENCES

- *Eckel WP, Jacob TA. 1988. Ambient levels of 24 dissolved metals in U.S. surface and ground waters. *Prepr Pap Natl Meet Am Chem Soc Div Environ Chem* 28:371-372.
- *Eckel WP, Langley WD. 1988. A background-based ranking technique for assessment of elemental enrichment in soils at hazardous waste sites. *Superfund '88. Proceedings of the 9th National Conference* Nov. 28-30, 1988. Washington, DC 282-286.
- Ecker FJ, Hirai E, Chohji T. 1990. Airborne trace metals in snow on the Japan sea side of Japan. *Atmos Environ* 24A(10):2593-2600.
- Eckert GE, Greene LW, Carstens GE, et al. 1999. Copper status of ewes fed increasing amounts of copper from copper sulfate or copper proteinate. *J Anim Sci* 77:244-249.
- *Effler SW, Litten S, Field SD, et al. 1980. Whole lake response to low level copper sulfate treatment. *Water Res* 14:1489-1499.
- Eife R, Weiss M, Barros V, et al. 1999. Chronic poisoning by copper in tap water: I. Copper intoxications with predominantly gastrointestinal symptoms. *Eur J Med Res* 4(6):219-223.
- *Eisenberg M, Topping JJ. 1986. Trace metal residues in fin fish from Maryland waters, 1978-1979. *J Environ Sci Health B* 21:87-102.
- Eitzer BD, Iannucci-Berger WA, Mark G, et al. 1997. Fate of toxic compounds during composting. *Bull Environ Contam Toxicol* 58:953-960.
- Eklind Y, Beck-Friis B, Bengtsson S, et al. 1997. Chemical characterization of source-separated organic household wastes. *Swed J Agric Res* 27:167-178.
- *Ellenhorn MJ, Schonwald S, Ordog G, eds. 1997. *Ellenhorn's medical toxicology*, 2nd ed. Baltimore, MD: Williams and Wilkins, 1554-1556.
- *Elliott HA, Liberati MR, Huang CP. 1986. Competitive adsorption of heavy metals by soils. *J Environ Qual* 15(3):214-219.
- *Ellis R, Morris ER, Hill AD, et al. 1997. Selected mineral intakes of adult African-Americans in the Washington, DC area. *J Food Comp Anal* 10:334-342.
- *EMMI. 1997. Environmental monitoring methods index. Version 1.1. PC#4082. Rockville, MD: U.S. Environmental Protection Agency, Government Institutes.
- EPA. 1979a. National secondary drinking water regulations. U.S. Environmental Protection Agency. *Fed Regist* 40(143):373-374.
- EPA. 1979b. Copper. Water-related environmental fate of 129 priority pollutants. U.S. Environmental Protection Agency. EPA440479029a.
- EPA. 1980a. Ambient water quality criteria for copper. Washington, DC: U.S. Environmental Protection Agency. EPA440580036.

9. REFERENCES

- EPA. 1980b. Primary copper industry. Industrial process profiles for environmental use. Cincinnati, OH: U.S. Environmental Protection Agency. EPA600280170.
- *EPA. 1981. Treatability manual. Volume 1. U.S. Environmental Protection Agency. EPA600282001a, I.4.7-1 to I.4.7-5.
- *EPA. 1983. Methods for chemical analysis of water and wastes. Washington, DC: U.S. Environmental Protection Agency. EPA600479020.
- *EPA. 1984. Air quality data for metals 1977 through 1979 from The National Air Surveillance Networks. U.S. Environmental Protection Agency. EPA600S483053.
- EPA. 1985. Notification requirements; reportable quantity adjustments, final rule and proposed rule. U.S. Environmental Protection Agency. Fed Regist 50(65):13456-13490.
- EPA. 1986. Test methods for evaluating solid waste1A: Laboratory manual physical/chemical methods. Washington, DC: U.S. Environmental Protection Agency.
- *EPA. 1987a. Assessment of copper as a potentially toxic air pollutant. U.S. Environmental Protection Agency. Fed Regist 52(35):5496-5499.
- EPA. 1987b. Drinking water criteria document for copper. Cincinnati, OH: U.S. Environmental Protection Agency. Research and Development.
- EPA. 1987c. Processes, coefficients, and models for simulating toxic organics and heavy metals in surface waters. Athens, GA: U.S. Environmental Protection Agency. EPA600387015.
- *EPA. 1988a. Mining waste exclusion. Fed Regist 53(203):41288.
- EPA. 1988b. Analysis of clean water act effluent guidelines pollutants: Summary of the chemicals regulated by industrial point source category. Code of Federal Regulations. 40 CFR Parts 400-475. Washington, DC: U.S. Environmental Protection Agency.
- EPA. 1988c. Copper. Chronic health assessment for noncarcinogenic effects. Washington, DC: U.S. Environmental Protection Agency.
- EPA. 1988d. Drinking water regulations; maximum contaminant level goals and national primary drinking water regulations for lead and copper. U.S. Environmental Protection Agency. Fed Regist 53(160):31516-31578.
- *EPA. 1990. Interim methods for development of inhalation reference concentrations. Washington, DC: U.S. Environmental Protection Agency, Office of Health and Environmental Assessment, Office of Research and Development, Environmental Criteria and Assessment Office. EPA600890066A.
- *EPA. 1991. Maximum contaminated level, goals and national primary drinking-water regulation for lead and copper. Final rule. Fed Regist 56:438-470.
- EPA. 1992. Pesticides in ground water database-A compilation of monitoring studies: 1971-1991. United States Environmental Protection Agency. EPA7341292001.

9. REFERENCES

- *EPA 1994. Methods for derivation of inhalation reference concentrations and application of inhalation dosimetry. Washington, DC. U.S. Environmental Protection Agency, Office of Research and Development.
- *EPA. 1995. Engineering forum issue: Determination of background concentrations of inorganics in soils and sediments at hazardous waste sites: U.S. Environmental Protection Agency, Office of Research and Development, Office of Solid Waste and Emergency Response.
- *EPA. 1997a. Automated Form R for Windows: User's guide (RY97). Washington, DC: U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics.
- *EPA. 1997b. Special report on environmental endocrine disruption: An effects assessment and analysis. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. EPA630R96012.
- *EPA. 1999. National recommended water quality criteria-correction. U.S. Environmental Protection Agency, Office of Water. EPA822Z99001.
- *EPA. 2002a. Effluent guidelines and standards. Toxic pollutants. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 401.15. <http://ecfrback.access.gpo.gov>. April 16, 2002.
- *EPA. 2002b. National emission standards for hazardous air pollutants. Lists of pollutants and applicability of part 61. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 61.01(b). <http://ecfrback.access.gpo.gov>. April 09, 2002.
- *EPA. 2002c. National primary drinking water regulations. Consumer factsheet on: Copper. [Wysiwyg://196/http://epa.gov/safewater/dwh/c-ioc/copper.html](http://196/http://epa.gov/safewater/dwh/c-ioc/copper.html). April 09, 2002.
- *EPA. 2002d. National primary drinking water regulations. Maximum contaminant level goals for inorganic contaminants. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 141.51(b). <http://ecfrback.access.gpo.gov>. April 09, 2002.
- *EPA. 2002e. National secondary drinking water regulations. Secondary maximum contaminant levels. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 143.3. <http://ecfrback.access.gpo.gov/otcg>. April 09, 2002.
- *EPA. 2002f. Pesticide programs. Copper; exemption from the requirement of a tolerance. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 180.1021. <http://ecfr.access.gpo.gov/otcgi/cf>. April 10, 2002.
- *EPA. 2002g. Standards for owner and operators of hazardous waste treatment, storage, and disposal facilities. Groundwater monitoring list. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 264, Appendix IX. <http://ecfr.access.gpo.gov>. April 10, 2001.
- *EPA. 2002h. Superfund, emergency planning, and community right-to-know programs. Designation, reportable quantities, and notifications. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 302.4. <http://ecfr.access.gpo.gov/otcgi>. April 10, 2002.
- *EPA. 2002i. Toxic chemical release reporting: Community right-to-know. Chemicals and chemical categories to which this part applies. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 377.65(a). <http://ecfr.access.gpo.gov/otcgi/cf>. April 09, 2002.

9. REFERENCES

- *EPA. 2002j. Water programs. Designation of hazardous substances. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 116.4. <http://ecfrback.access.gpo.gov/otcg>. April 10, 2002.
- *EPA. 2002k. Water programs. Determination of reportable quantities for hazardous substances. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 117.3. <http://ecfrback.access.gpo.gov/otcg>. April 10, 2002.
- *Epstein O, Spisni R, Parbhoo S, et al. 1982. The effect of oral copper loading and portasystemic shunting on the distribution of copper in the liver, brain, kidney, and cornea of the rat. *Am J Clin Nutr* 35:551-555.
- Ettinger MJ, Darwish HM, Schmitt RC. 1986. Mechanism of copper transport from plasma to hepatocytes. *Fed Proc* 45:2800-2804.
- Evans DW, Dodoo DK, Hanson PJ. 1993. Trace elements concentrations in fish livers: Implications of variations with fish size in pollution monitoring. *Mar Pollut Bull* 26(6):329-334.
- *Evans GW, Leblanc FN. 1976. Copper-binding protein in rat intestine: Amino acid composition and function. *Nutr Rep Int* 14(3):281-288.
- *Evans GW, Majors PF, Cornatzer WE. 1970a. Mechanism for cadmium and zinc antagonism of copper metabolism. *Biochem Biophys Res Commun* 40:1142-1148.
- *Evans GW, Majors PF, Cornatzer WE. 1970b. Induction of ceruloplasmin synthesis by copper. *Biochem Biophys Res Commun* 41(5):1120-1125.
- *Evering WE, Haywood S, Bremner I, et al. 1991a. The protective role of metallothionein in copper overload: I. Differential distribution of immunoreactive metallothionein in copper-loaded rat liver and kidney. *Chem Biol Interact* 78(3):283-295.
- *Evering WE, Haywood S, Bremner I, et al. 1991b. The protective role of metallothionein in copper-overload. II. Transport and excretion of immunoreactive MT-1 in blood, bile and urine of copper-loaded rats. *Chem Biol Interact* 78(3):297-305.
- Fairfax R. 1996. Exposure to copper, lead, and confined spaces in a foundry. *Appl Occup Environ Hyg* 11(12):1379-1381.
- Fairweather-Tait SJ. 1997. Bioavailability of copper. *Eur J Clin Nutr* 51:S24-S26.
- Frag AM, Woodward DF, Goldstein JN, et al. 1998. Concentrations of metals associated with mining waste in sediments, biofilm, benthic macroinvertebrates, and fish from the Coeur d'Alene River Basin, Idaho. *Arch Environ Contam Toxicol* 34:119-127.
- Farrah H, Pickering WF. 1993. Factors influencing the potential mobility and bioavailability of metals in dried lake sediments. *Chem Speciat Bioavail* 5(3):81-96.
- *Farrer P, Mistilis SP. 1967. Absorption of exogenous and endogenous biliary copper in the rat. *Nature* 213:291-292.

9. REFERENCES

- *FDA. 2000. Total diet study statistics on element results. Washington, DC: U.S. Food and Drug Administration.
- *FDA. 2001a. Beverages. Bottled water. U.S. Food and Drug Administration. Code of Federal Regulations. 21 CFR 165.110. <http://www.frwebgate.access.gpo.gov/cgi>. April 09, 2002.
- *FDA. 2001b. Clinical chemistry and clinical toxicology devices. Copper test system. U.S. Food and Drug Administration. Code of Federal Regulations. 21 CFR 862.1190. <http://www.frwebgate.accessgpo.gov/cgi>. April 09, 2002.
- *FDA. 2001c. Direct food substances affirmed as generally recognized as safe. Copper sulfate. U.S. Food and Drug Administration. Code of Federal Regulations. 21 CFR 184.1261. <http://www.frwebgate.accessgpo.gov/cgi>. April 10, 2002.
- *FDA. 2001d. Food labeling. Nutritional labeling of food. U.S. Food and Drug Administration. Code of Federal Regulations. 21 CFR 101.9(c)(8). <http://www.frwebgate.accessgpo.gov/cgi>. April 09, 2002.
- *FDA. 2001e. Listing of color additives exempt from certification. Copper powder. U.S. Food and Drug Administration. Code of Federal Regulations 21 CFR 73.1647. <http://www.frwebgate.accessgpo.gov/cgi>. April 09, 2002.
- *FDA. 2001f. Listing of color additives exempt from certification. Copper powder. U.S. Food and Drug Administration. Code of Federal Regulations 21 CFR 73.2647. <http://www.frwebgate.accessgpo.gov/cgi>. April 09, 2002.
- *FDA. 2001g. New drugs. Drug products containing certain active ingredients offered over-the-counter. U.S. Food and Drug Administration. Code of Federal Regulations. 21 CFR 310.545(a)(20). <http://www.frwebgate.accessgpo.gov/cgi>. April 09, 2002.
- *FDA. 2001h. Nutritional quality guidelines for foods. Statement of purpose. U.S. Food and Drug Administration. Code of Federal Regulations. 21 CFR 104.20(d)(3). <http://www.frwebgate.accessgpo.gov/cgi>. April 09, 2002.
- *FDA. 2001i. Substances generally recognized as safe. Trace minerals added to animal feeds. U.S. Food and Drug Administration. Code of Federal Regulations. 21 CFR 582.80. <http://www.frwebgate.accessgpo.gov/cgi>. April 10, 2002.
- FEDRIP. 1988. Federal Research In Progress Database. National Technical Information Service, Springfield, VA.
- *FEDRIP. 2003. Federal Research In Progress. Dialog Information Services, Inc.
- *Feiler HD, Storch PJ, Southworth R. 1980. Organics in municipal sludges survey of forty cities. Silver Spring, MD; National Conference: Municipal Industrial Sludge Utility Disposal.
- *Feng X, Melander AP, Klaue B. 2000. Contribution of municipal waste incineration to trace metal deposition on the vicinity. *Water Air Soil Pollut* 119:295-316.
- *Fergusson JE, Stewart C. 1992. The transport of airborne trace elements copper, lead, cadmium, zinc and manganese from a city into rural areas. *Sci Total Environ* 121:247-269.

9. REFERENCES

- Ferm VH, Hanlon DP. 1974. Toxicity of copper salts in hamster embryonic development. *Biol Reprod* 11:97-101.
- *Fernandes AC, Filipe PM, Manso CF. 1992. Protective effects of a 21-aminosteroid against copper-induced erythrocyte and plasma lipid peroxidation. *Eur J Pharmacol* 220(2-3):211-216.
- Fernandes A, Mira ML, Azevedo MS, et al. 1988. Mechanisms of hemolysis induced by copper. *Free Radic Res Commun* 4(5):291-298.
- Fewtrell L, Kay D, MacGill S. 2001. A review of the science behind drinking water standards for copper. *Int J Environ Health Res* 11(2):161-167.
- Fields M, Holbrook J, Scholfield D, et al. 1986. Effect of fructose or starch on copper-67 absorption and excretion by the rat. *J Nutr* 116:625-632.
- *Finelli VN, Boscolo P, Salimei E, et al. 1981. Anemia in men occupationally exposed to low levels of copper. *Heavy Met Environ Int Conf* 4th, 475-478.
- *Finley EB, Cerklewski FL. 1983. Influence of ascorbic acid supplementation on copper status in young adult men. *Am J Clin Nutr* 37(4):553-556.
- *Fischer, PWF, Giroux A, L'Abbe AR. 1984. Effect of zinc supplementation on copper status in adult man. *Am J. Clin Nutr* 40:743-746.
- Fischer PWF, L'Abbe MR, Giroux A. 1990. Effects of age, smoking, drinking, exercise and estrogen use on indices of copper status in healthy adults. *Nutr Res* 10(10):1081-1090.
- Fishman JH, Fishman J. 1988. Copper and endogenous mediators of estradiol action. *Biochem Biophys Res Commun* 152(2):783-788.
- Fitzgerald DJ. 1995. Copper guideline values for drinking water: Reviews in need of review? *Regul Toxicol Pharmacol* 21:177-179.
- Fitzgerald DJ. 1996. Copper regulatory level in drinking-water as proposed by Sidhu et al. *Regul Toxicol Pharmacol* 23:173-175.
- Fitzgerald DJ. 1998. Safety guidelines for copper in water. *Am J Clin Nutr* 67(Suppl):1098S-1102S.
- *Fleckman P. 1985. Anatomy and physiology of the nail. *Dermatol Clin* 3(3):373-381.
- *Fomon SJ. 1966. Body composition of the infant: Part I: The male "reference infant". In: Falkner F, ed. *Human development*. Philadelphia, PA: WB Saunders, 239-246.
- *Fomon SJ, Haschke F, Ziegler EE, et al. 1982. Body composition of reference children from birth to age 10 years. *Am J Clin Nutr* 35:1169-1175.
- *Ford ES. 2000. Serum copper concentration and coronary heart disease among U.S. adults. *Am J Epidemiol* 151(12):1182-1188.
- Forman SJ, Kumar KS, Redeker AG, et al. 1980. Hemolytic anemia in Wilson disease: Clinical findings and biochemical mechanisms. *Am J Hematol* 9:269-275.

9. REFERENCES

- *Fox DL. 1987. Air pollution. *Anal Chem* 59:280R-294R.
- Fox JG, Zeman DH, Mortimer JD. 1994. Copper toxicosis in sibling ferrets. *J Am Vet Med Assoc* 205(8):1154-1156.
- Fraser JK, Butler CA, Timperley MH, et al. 2000. Formation of copper complexes in landfill leachate and their toxicity to zebrafish embryos. *Environ Toxicol Chem* 19(5):1394-1402.
- *Frazier LM, Hage ML. 1998. Appendix 1 Occupational exposure limits for chemicals. In: *Reproductive hazards of the workplace*. New York: Van Nostrand Reinhold, 537-543.
- Freedman JM, Peisach J. 1989a. Intracellular copper transport in cultured hepatoma cells. *Biochem Biophys Res Commun* 164(1):134-140.
- *Freedman JM, Peisach J. 1989b. Resistance of cultured hepatoma cells to copper toxicity. Purification and characterization of the hepatoma metallothionein. *Biochim Biophys Acta* 992:145-154.
- Freedman JH, Ciriolo MR, Peisach J. 1989. The role of glutathione in copper metabolism and toxicity. *J Biol Chem* 264(10):5598-5605.
- Freedman JH, Pickart L, Weinstein B, et al. 1982. Structure of the Glycyl-L-lysine-Copper(II) complex in solution. *Biochemistry* 21:4540-4544.
- Frias-Espericueta MG, Osuna-Lopez JI, Sandoval-Salazar G, et al. 1999. Distribution of trace metals in different tissues in the rock oyster *Crassostrea iridescens*: Seasonal variation. *Bull Environ Contam Toxicol* 63:73-79.
- *Frost DV, Olson OE. 1972. The two faces of selenium-can selenophobia be cured? *CRC Crit Rev Toxicol* 1:467-514.
- FSTRAC. 1988. Summary of state and federal drinking water standards and guidelines: 8 & 16.
- Fuentealba IC, Bratton GR. 1994. The role of the liver, kidney and duodenum in tolerance in the copper-loaded rat. *Toxicol* 6(4):345-358.
- *Fuentealba IC, Haywood S. 1988. Cellular mechanisms of toxicity and tolerance in the copper-loaded rat. I. Ultrastructural changes in the liver. *Liver* 8:372-380.
- Fuentealba IC, Davis RW, Elmes ME, et al. 1993. Mechanisms of tolerance in the copper-loaded rat liver. *Exp Mol Pathol* 59(1):71-84.
- Fuentealba IC, Haywood S, Foster J. 1989a. Cellular mechanisms of toxicity and tolerance in the copper-loaded rat. II. Pathogenesis of copper toxicity in the liver. *Exp Mol Pathol* 50:26-37.
- Fuentealba IC, Haywood S, Foster J. 1989b. Cellular mechanisms of toxicity and tolerance in the copper-loaded rat. III. Ultrastructural changes and copper localization in the kidney. *Br J Exp Pathol* 70(5):543-556.

9. REFERENCES

- Fuentealba IC, Haywood S, Trafford J. 1989c. Variations in the intralobular distribution of copper in the livers of copper-loaded rats in relation to the pathogenesis of copper storage diseases. *J Comp Pathol* 100(1):1-11.
- *Fuentealba IC, Mullins JE, Aburto EM, et al. 2000. Effect of age and sex on liver damage due to excess dietary copper in Fischer 344 rats. *Clin Toxicol* 38(7):709-717.
- *Fuhrer GJ. 1986. Extractable cadmium, mercury, copper, lead, and zinc in the lower Columbia River Estuary, Oregon and Washington. U.S. Geological Survey Water Resources Investigations Report 86(4088). Portland, Oregon: U.S. Department of Interior.
- *Fukui H, Yamamoto M, Sasaki S, et al. 1993. Involvement of 5-HT₃ receptors and vagal afferents in copper sulfate- and cisplatin-induced emesis in monkeys. *Eur J Pharmacol* 249(1):13-18.
- *Fukui H, Yamamoto M, Sasaki S, et al. 1994. Possible involvement of peripheral 5-HT₄ receptors in copper sulfate-induced vomiting in dogs. *Eur J Pharmacol* 257(1-2):47-52.
- *Furst A. 1971. Trace elements related to specific chronic diseases: Cancer. *Environmental geochemistry in health and disease*. The Geological Society of America, Inc., 123:109-130.
- Gaetke LM, Chow CK. 2003. Copper toxicity, oxidative stress, and antioxidant nutrients. *Toxicology* 189(1-2):147-163.
- Galasinska-Pomykol I, Moniuszko-Jakoniuk J, Pietrewicz T. 1993. Changes in neurosecretory material content in neurohypophysis of rats submitted to chronic treatment with copper. *Ann Med Univ Bialyst Pol* 38(1):72-78.
- *Gallagher CH. 1979. Biochemical and pathological effects of copper deficiency. In: Nriagu JO, ed. *Copper in the environment*. New York, NY: John Wiley & Sons, 58-82.
- Gallagher MD, Nuckols JR, Stallones L, et al. 1998. Exposure to trihalomethanes and adverse pregnancy outcomes. *Epidemiology* 9(5):484-489.
- *Galloway JN, Norton SA, Volchok HL, et al. 1982. Trace metals in atmospheric deposition: A review and assessment. *Atmos Environ* 16(7):1677-1700.
- Gambling L, Danzeisen R, Fosset C, et al. 2003. Iron and copper interactions in development and the effect on pregnancy outcome. *J Nutr* 133(5 Suppl 1):1554S-1556S.
- *Gao L, Li R, Wang K. 1989. Kinetic studies of mobilization of copper (II) from human serum albumin with chelating agents. *J Inorg Biochem* 36(2):83-92.
- *Gao S, Walker WJ, Dahlgren RA. 1997. Simultaneous sorption of Cd, Cu, Ni, Zn, Pb, and Cr on soils treated with sewage sludge supernatant. *Water Air Soil Pollut* 93:331-345.
- Gardner M, Ravenscroft J. 1991. The behavior of copper complexation in rivers and estuaries: Two studies in North East England. *Chemosphere* 23(6):695-713.
- Garnier J-M, Pham MK, Ciffroy P, et al. 1997. Kinetics of trace element complexation with suspended matter and with filterable ligands in freshwater. *Environ Sci Technol* 31:1597-1606.

9. REFERENCES

- *Garrett NE, Lewtas J. 1983. Cellular toxicity in Chinese hamster ovary cells culture I. Environ Res 32:455-465.
- Garty J, Kauppi M, Kauppi A. 1996. Accumulation of airborne elements from vehicles in transplanted lichens in urban sites. J Environ Qual 25:265-272.
- *Georgopoulos AR, Yonone-Lioy MJ, Opiekun RE, et al. 2001. Environmental copper: Its dynamics and human exposure issues. J Toxicol Environ Health Part B Crit Rev 4(4):341-394.
- *Gerhardsson L, Brune D, Lundstrom N-G, et al. 1993. Biological specimen bank for smelter workers. Sci Total Environ 139/140:157-173.
- Gerhardsson L, Brune D, Nordberg GF, et al. 1988. Occupation-related cancer in a Nordic copper smeltery. Arctic Med Res 47:628-631.
- *Gerritse RG, Driel WV. 1984. The relationship between adsorption of trace metals, organic matter, and pH in temperate soils. J Environ Qual 13(2):197-204.
- Gibbs PJ, Miskiewicz AG. 1995. Heavy metals in fish near a major primary treatment sewage plant outfall. Mar Pollut Bull 30(10):667-674.
- *Giesy JP, Briese LA, Leversee GJ. 1978. Metal binding capacity of selected Maine surface waters. Environ Geol 2(5):257-268.
- *Giesy JP, Newell A, Leversee GJ. 1983. Copper speciation in soft, acid, humic waters: Effects on copper bioaccumulation by and toxicity to *Simocephalus serrulatus*. Sci Total Environ 28:23-36.
- *Gill JS, Bhagat CI. 1999. Acute copper poisoning from drinking lime cordial prepared and left overnight in an old urn. Med J Aust 170(10):510.
- *Gilman JPW. 1962. Metal carcinogenesis: II. A study on the carcinogenic activity of cobalt, copper, iron, and nickel compounds. Cancer Res 22:158-162.
- *Gitlin D, Hughes WL, Janeway CA. 1960. Absorption and excretion of copper in mice. Nature 188(4745):150-151.
- *Giusquiani PL, Gigliotti G, Businelli D. 1992. Mobility of heavy metals in urban waste-amended soils. J Environ Qual 21:330-335.
- *Giusti L, Yang Y-L, Hewitt CN, et al. 1993. The solubility and partitioning of atmospherically derived trace metals in artificial and natural waters: A review. Atmos Environ 27A(10):1567-1578.
- *Giwerzman A, Carlsen E, Keiding N, et al. 1993. Evidence for increasing incidence of abnormalities of the human testis: A review. Environ Health Perspect Suppl 101(2):65-71.
- *Gleason RP. 1968. Exposure to copper dust. Am Ind Hyg Assoc J 29:461-462.
- *Goldberg ED. 1986. The mussel watch concept. Environ Monit Assess 7:91-103.
- *Goldin A, Bigelow C, Veneman PLM. 1992. Concentrations of metals in ash from municipal solid waste combustors. Chemosphere 24(3):271-280.

9. REFERENCES

- *Gollan JL, Deller DJ. 1973. Studies on the nature and excretion of biliary copper in man. *Clin Sci* 44:9-15.
- *Golomb D, Ryan D, Eby N, et al. 1997. Atmospheric deposition of toxics onto Massachusetts Bay-I. Metals. *Atmos Environ* 31(9):1349-1359.
- Gonzalez-Vila FJ, Bautista JM, Del Rio JC, et al. 1993. Evolution of chemicals within the dump profile in a controlled landfill. *Chemosphere* 31(3):2817-2825.
- Gooneratne SR, Gawthorne JM, Howell JM. 1989a. Distribution of Cu, Zn, and Fe in the soluble fraction of the kidney in normal, copper-poisoned, and thiomolybdate-treated sheep. *J Inorg Biochem* 35:37-53.
- Gooneratne SR, Howell JM, Gawthorne JM, et al. 1989b. Subcellular distribution of copper in the kidneys of normal, copper-poisoned, and thiomolybdate-treated sheep. *J Inorg Biochem* 35:23-36.
- Gorlach U, Boutron CF. 1992. Variations in heavy metals concentrations. *J Atmos Chem* 14:205-222.
- *Gotteland M, Araya M, Pizarro F, et al. 2001. Effect of acute copper exposure on gastrointestinal permeability in healthy volunteers. *Dig Dis Sci* 46(9):1909-1914.
- *Gralak MA, Leontowicz M, Morawiec M, et al. 1996. Comparison of the influence of dietary fibre sources with different proportions of soluble and insoluble fibre on Ca, Mg, Fe, Zn, Mn and Cu apparent absorption in rats. *Arch Tierernaehr* 49(4):293-299.
- *Greenberg AE, Trussell RR, Clesceri LS, eds. 1985. Copper: Neocuprione method. In: *Standard methods: For the examination of water and wastewater*. Washington, DC: American Public Health Association, 205-207.
- *Greene FL, Lamb LS, Barwick M, et al. 1987. Effect of dietary copper on colonic tumor production and aortic integrity in the rat. *J Surg Res* 42:503-512.
- Greger JL, Zaikis SC, Abernathy RP, et al. 1978. Zinc, nitrogen, copper, iron and manganese balance in adolescent females fed two levels of zinc. *J Nutr* 108:1449-1456.
- Gregori ID, Pinochet H, Arancibia M, et al. 1996a. Grain size effect on trace metals distribution in sediments from two coastal areas of Chile. *Bull Environ Contam Toxicol* 57:163-170.
- Gregori ID, Pinochet H, Gras N, et al. 1996b. Variability of cadmium, copper and zinc levels in molluscs and associated sediments from Chile. *Environ Pollut* 92(3):359-368.
- *Gregus Z, Klaassen CD. 1986. Disposition of metals in rats: A comparative study of fecal, urinary, and biliary excretion and tissue distribution of eighteen metals. *Toxicol Appl Pharmacol* 85:24-38.
- Gromping AHJ, Ostapczuk P, Emons H. 1997. Wet deposition in Germany: Long-term trends and the contribution of heavy metals. *Chemosphere* 34(9/10):2227-2236.
- *Gross AF, Given PS, Athnasios AK. 1987. Food. *Anal Chem* 59:212R-252R.
- Gulliver JM. 1991. A fatal copper sulfate poisoning. *J Anal Toxicol* 15(6):341-342.

9. REFERENCES

- Gulson BL, Mizon KJ, Korsch, MJ, et al. 2001. Dietary intakes of selected elements from longitudinal 6-day duplicate diets for pregnant and nonpregnant subjects and elemental concentrations of breast milk and infant formula. *Environ Res Sect A* 87:160-174.
- *Gupta UC. 1979. Copper in agricultural crops. Nriagu JO, ed. In: *Copper in the environment. Part I: Ecological Cycling*. New York: John Wiley & Sons Inc.
- *Gutenmann WH, Rutzke M, Kuntz HT, et al. 1994. Elements and polychlorinated biphenyls in sewage sludges of large cities in the United States. *Chemosphere* 28(4):725-728.
- *Guzelian PS, Henry CJ, Olin SS, eds. 1992. *Similarities and differences between children and adults: Implications for risk assessment*. Washington, DC: International Life Sciences Institute Press.
- *Haddad DS, al-Alousi A, Kantarjian AH. 1991. The effect of copper loading on pregnant rats and their offspring. *Funct Dev Morphol* 1(3):17-22.
- *Haines RC. 1984. Environmental contamination-surveys of heavy metals in urban soils and hazard assessment. *Trace Subst Environ Health* 18:450-460.
- *Hall AC, Young BW, Bremner I. 1979. Intestinal metallothionein and the mutual antagonism between copper and zinc in the rat. *J Inorg Biochem* 11:57-66.
- Hall WS, Pulliam GW. 1995. An assessment of metals in an estuarine wetlands ecosystem. *Arch Environ Contam Toxicol* 29:164-173.
- Hanna LA, Peters JM, Wiley LM, et al. 1997. Comparative effects of essential and nonessential metals on preimplantation mouse embryo development in vitro. *Toxicology* 116(1-3):123-131.
- Harada M, Sakisaka S, Yoshitake S, et al. 1993. Biliary copper excretion in acutely and chronically copper-loaded rats. *Hepatology* 17(1):111-117.
- Harris ED. 1991. Copper transport: An overview. *Proc Soc Exp Biol Med* 196(2):130-140.
- *Harris ED. 1993. The transport of copper: Essential and toxic trace elements in human health and disease. *Prog Clin Biol Res* 380:163-179.
- Harris ED, Percival SS. 1989. Copper transport: Insights into a ceruloplasmin-based delivery system. *Adv Exp Med Biol* 258:95-102.
- Harris ED, Percival SS. 1991. A role for ascorbic acid in copper transport. *Am J Clin Nutr* 54:1193S-1197S.
- *Harris ED, Qian Y, Tiffany-Castiglioni E, et al. 1998. Functional analysis of copper homeostasis in cell culture models: a new perspective on internal copper transport. *Am J Clin Nutr* 67:988S-995S.
- Harris ZL, Gitlin JD. 1996. Genetic and molecular basis for copper toxicity. *Am J Clin Nutr* 63(5):836S-841S.
- *Harrison BJ. 1998. Table 1. Copper concentrations in the environment. In: *Copper information sourcebook- 1998- the world's scientific literature on copper in the environment and health*, 21-62.

9. REFERENCES

- Harrison FL. 1982. A review of the impact of copper released into marine and estuarine environments. U.S. Nuclear Regulatory Commission. Washington, DC: Lawrence Livermore Laboratory. NUREG/CR-2823.
- *Harrison FL, Bishop DJ. 1984. A review of the impact of copper released into freshwater environments. U.S. Nuclear Regulatory Commission. Livermore, CA: Lawrence Livermore National Laboratory. NUREG/CR-3478.
- *Harrison FL, Bishop DJ, Emerson RR, et al. 1980. In: U.S. Nuclear Regulatory Commission eds. Concentration and speciation of copper in waters collected near the San Onofre and Diablo Canyon nuclear power stations: Washington, DC: Lawrence Livermore Laboratory. NUREG/CR-0750.
- Harrison RM, Jones M. 1995. The chemical composition of airborne particles in the UK atmosphere. *Sci Total Environ* 168:195-214.
- Hasan N, Emery D, Baithun D. 1995. Chronic copper intoxication due to ingestion of coins: a report of an unusual case. *Hum Exp Toxicol* 14(6):500-502.
- *Haschke F, Ziegler EE, Edwards BB, et al. 1986. Effect of iron fortification of infant formula on trace mineral absorption. *J Pediatr Gastroenterol Nutr* 5(5):768-773.
- Hassan SM, Garrison AW, Allen HE, et al. 1996. Estimation of partition coefficients for five trace metals in sandy sediments and application to sediment quality criteria. *Environ Toxicol Chem* 15(12):2198-2208.
- *Hawley GG. 1981. *The condensed chemical dictionary*. New York, NY: Van Nostrand Reinhold Company. 273-274.
- Hayashi M, Kuge T, Endoh D, et al. 2000. Hepatic copper accumulation induces DNA strand breaks in the liver cells of Long-Evans Cinnamon strain rats. *Biochem Biophys Res Commun* 276(1):174-178.
- Hayward DG, Petreas MX, Winkler JJ, et al. 1996. Investigation of a wood treatment facility: Impact on an aquatic ecosystem in the San Joaquin River, Stockton, California. *Arch Environ Contam Toxicol* 30:30-39.
- *Haywood S. 1980. The effect of excess dietary copper on the liver and kidney of the male rat. *J Comp Pathol* 90:217-232.
- *Haywood S. 1985. Copper toxicosis and tolerance in the rat: I-changes in copper content of the liver and kidney. *J Pathol* 145:149-158.
- *Haywood S, Comerford B. 1980. The effect of excess dietary copper on plasma enzyme activity and on the copper content of the blood of the male rat. *J Comp Pathol* 90:233-238.
- *Haywood S, Loughran M. 1985. Copper toxicosis and tolerance in the rat. II. Tolerance-a liver protective adaptation. *Liver* 5:267-275.
- *Haywood S, Loughran M, Batt RM. 1985a. Copper toxicosis and tolerance in the rat. III. Intracellular localization of copper in the liver and kidney. *Exp Mol Pathol* 43:209-219.

9. REFERENCES

- *Haywood S, Trafford J, Loughran M. 1985b. Copper toxicosis and tolerance in the rat. IV. Renal tubular excretion of copper. *Br J Exp Pathol* 66:699-707.
- *HazDat. 2002. Agency for Toxic Substances and Disease Registry (ATSDR). Atlanta, GA. <http://www.atsdr.cdc.gov/hazdat.html>. May 8, 2002.
- *HazDat. 2004. Agency for Toxic Substances and Disease Registry (ATSDR). Atlanta, GA. <http://www.atsdr.cdc.gov/hazdat.html>. July 29, 2004.
- *He X-T, Logan TJ, Traina SJ. 1995. Physical and chemical characteristics of selected US municipal solid waste compost. *J Environ Qual* 24:543-552.
- Headley AD. 1996. Heavy metal concentrations in peat profiles from the high Arctic. *Sci Total Environ* 177:105-111.
- Hebert CD, Elwell MR, Travlos GS, et al. 1993. Subchronic toxicity of cupric sulfate administered in drinking water and feed to rats and mice. *Fundam Appl Toxicol* 21(4):461-475.
- Heiny JS, Tate CM. 1997. Concentration, distribution, and comparison of selected trace elements in bed sediment and fish tissue in the South Platte River Basin, U.S.A., 1992-1993. *Arch Environ Contam Toxicol* 32:246-259.
- *Heit M, Klusek CS. 1985. Trace element concentrations in the dorsal muscle of white suckers and brown bullheads from two acidic Adirondack lakes. *Water Air Soil Pollut* 25:87-96.
- Helgen SO, Moore JN. 1996. Natural background determination and impact quantification in trace metal-contaminated river sediments. *Environ Sci Technol* 30:129-135.
- *Hellou J, Fancey LL, Payne JF. 1992a. Concentrations of twenty-four elements in bluefin tuna, *Thunnus thynnus* from the Northwest Atlantic. *Chemosphere* 24(2):211-218.
- *Hellou J, Warren WG, Payne JF, et al. 1992b. Heavy metals and other elements in three tissues of cod, *Gadus morhua* from the Northwest Atlantic. *Mar Pollut Bull* 24(9):452-458.
- Helmers E, Schrems O. 1995. Wet deposition of metals to the tropical north and the south Atlantic Ocean. *Atmos Environ* 29(18):2475-2484.
- *Helz GR, Huggett RJ, Hill JM. 1975. Behavior of Mn, Fe, Cu, Zn, Cd, and Pb discharged from a wastewater treatment plant into an estuarine environment. *Water Res* 9:631-636.
- Heramanson MH. 1993. Historical accumulation of atmospherically derived pollutant trace metals in the arctic as measured in dated sediment cores. *Water Sci Technol* 28(8-9):33-41.
- *Herawati N, Suzuki S, Hayashi K, et al. 2000. Cadmium, copper, and zinc levels in rice and soil of Japan, Indonesia, and China by soil type. *Bull Environ Contam Toxicol* 64:33-39.
- Hering JG, Morel FMM. 1988. Kinetics of trace metal complexation: Role of alkaline-earth metals. *Environ Sci Technol* 22:1469-1478.
- *Hermann R, Newmann-Mahlkau P. 1985. The mobility of zinc, cadmium, copper, lead, iron, and arsenic in ground water as a function of redox potential and pH. *Sci Total Environ* 43:1-12.

9. REFERENCES

- *Hernandez LM, Gonzalez MJ, Rico MC, et al. 1985. Presence and biomagnification of organochlorine pollutants and heavy metals in mammals of Donana National Park (Spain) 1982-1983. *J Environ Sci Health B* 20:633-650.
- Hildemann LM, Markowski GR, Cass GR. 1991. Chemical composition of emissions from urban sources of fine organic aerosol. *Environ Sci Technol* 25:744-759.
- Hirano S, Ebihara H, Sakai S, et al. 1993. Pulmonary clearance and toxicity of intratracheally instilled cupric oxide in rats. *Arch Toxicol* 67(5):312-317.
- *Hirano S, Sakai S, Ebihara H, et al. 1990. Metabolism and pulmonary toxicity of intratracheally instilled cupric sulfate in rats. *Toxicology* 64(3):223-233.
- Hochstein P, Kumar KS, Forman SJ. 1978. Mechanisms of copper toxicity in red cells. *Prog Clin Biol Res* 21:669-686.
- *Hoel DG, Davis DL, Miller AB, et al. 1992. Trends in cancer mortality in 15 industrialized countries, 1969-1986. *J Natl Cancer Inst* 84(5):313-320.
- Hoffmann P, Dedik AN, Deutsch F, et al. 1997. Solubility of single chemical compounds from an atmosphere aerosol in pure water. *Atmos Environ* 31(17):2777-2785.
- *Holak W. 1983. Determination of copper, nickel, and chromium in foods. *J Assoc Off Anal Chem* 66(3):620-624.
- *Holland MK, White IG. 1988. Heavy metals and human spermatozoa. III. The toxicity of copper ions for spermatozoa. *Contraception* 38(6):685-695.
- *Holleran RS. 1981. Copper sulfate overdose. *J Emerg Nurs* 7:136-137.
- *Holmgren GGS, Meyer MW, Chaney RL, et al. 1993. Cadmium, lead, zinc, copper, and nickel in agricultural soils of the United States of America. *J Environ Qual* 22:335-348.
- *Holtzman NA, Elliott DA, Heller RH. 1966. Copper intoxication. *N Engl J Med* 275:347-352.
- *Hoogenraad TU, Koevuet R, deRuyter Korver EG. 1979. Oral zinc sulphate as long-term treatment in Wilson's disease (hepatolenticular degeneration). *Eur Neurol* 18:205-211.
- *Hopps HC. 1977. The biologic bases for using hair and nail for analyses of trace elements. *Sci Total Environ* 7:71-89.
- *Horng C-J. 1996. Simultaneous determination of urinary zinc, cadmium, lead and copper concentrations in steel production workers by differential-pulse anodic stripping voltammetry. *Analyst* 121(10):1511-1514.
- Horowitz AJ, Lum KR, Garbarino JR, et al. 1996. Problems associated with using filtration to define dissolved trace element concentrations in natural water samples. *Environ Sci Technol* 30:954-963.
- Hosovski E, Sunderic D, Sindji M. 1990. [Functional and histologic changes in the kidney in copper poisoning in rats.] *Srp Arh Celok Lek* 118(11-12):445-449. (Serbo-Croatian, Cyrillic)

9. REFERENCES

- *HSDB. 2004. Hazardous Substances Data Bank. <http://toxnet.nlm.nih.gov/cgi-bin/sis/html>.
- *Huh CA. 1996. Fluxes and budgets of anthropogenic metals in the Santa Monica and San Pedro Basins off Los Angeles: Review and reassessment. *Sci Total Environ* 179:47-60.
- *Humphries WR, Morrice PC, Bremner I. 1988. A convenient method for the treatment of chronic copper poisoning in sheep using subcutaneous ammonium tetrathiomolybdate. *Vet Rec* 123(2):51-53.
- Hurley JP, Shafer MM, Cowell SE, et al. 1996. Trace metal assessment of Lake Michigan tributaries using low-level techniques. *Environ Sci Technol* 30:2093-2098.
- *Hutchinson TC. 1979. Copper contamination of ecosystems caused by smelter activities. In: Nriagu JO, ed. *Copper in the environment. Part I: Ecological cycling*. New York: John Wiley and Sons Inc.
- Hwang DF, Wang LC, Cheng HM. 1998. Effect of taurine on toxicity of copper in rats. *Food Chem Toxicol* 36(3):239-244.
- *Iannuzzi TJ, Huntley SL, Schmidt CW, et al. 1997. Combined sewer overflows (CSOs) as sources of sediment contamination in the lower Passaic River, New Jersey. I. Priority pollutants and inorganic chemicals. *Chemosphere* 34(2):213-231.
- *IARC. 1987. IARC monographs on the evaluation of carcinogenic risks to humans. 31-32, 61.
- *IARC. 2002. Overall evaluation of carcinogenicity to humans. Group 3: Unclassifiable as to carcinogenicity to humans. IARC monographs programme on the evaluation of carcinogenic risks to humans. International Agency for Research on Cancer. <http://193.51.164.11/monoeval/crthgr03htm>. April 09, 2002.
- *IOM. 2001. Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium and zinc. Institute of Medicine, Food and Nutrition Board. Washington DC. National Academy Press.
- *IRIS. 2004. Copper. Integrated Risk Information System. <http://www.epa.gov/iris/subst/0368.htm>. February 10, 2004.
- *Isaac RA, Gil L, Cooperman AN, et al. 1997. Corrosion in drinking water distribution systems: A major contributor of copper and lead to wastewaters and effluents. *Environ Sci Technol* 31:3198-3203.
- *Iyengar V, Woittiez J. 1988. Trace elements in human clinical specimens: Evaluation of literature data to identify reference values. *Clin Chem* 34(3):474-481.
- Jackson DR, Garrett BC, Bishop TA. 1984. Comparison of batch and column methods for assessing leachability of hazardous waste. *Environ Sci Technol* 18(9):668-673.
- *Jacob RA, Skala JH, Omaye ST, et al. 1987. Effect of varying ascorbic acid intakes on copper absorption and ceruloplasmin levels of young men. *J Nutr* 117:2109-2115.
- *Janssen RPT, Peijnenburg WJGM, Posthuma L, et al. 1997. Equilibrium partitioning of heavy metals in Dutch field soils: I. Relationship between metal partition coefficients and soil characteristics. *Environ Toxicol Chem* 16(12):2470-2478.

9. REFERENCES

- *Jantsch W, Kulig K, Rumack BH. 1985. Massive copper sulfate ingestion resulting in hepatotoxicity. *Clin Toxicol* 22(6):585-588.
- *Jenkins D, Russell LL. 1994. Heavy metals contribution of household washing products to municipal wastewater. *Water Environ Res* 66:805-813.
- Jiménez I, Aracena P, Letelier ME, et al. 2002. Chronic exposure of HepG2 cells to excess copper results in depletion of glutathione and induction of metallothionein. *Toxicol in Vitro* 16:167-175.
- *Johanson CE. 1980. Permeability and vascularity of the developing brain: Cerebellum vs cerebral cortex. *Brain Res* 190:3-16.
- *Johansson A, Camner P, Jastrand C, et al. 1983. Rabbit alveolar macrophages after inhalation of soluble cadmium, cobalt, and copper: A comparison with the effects of soluble nickel. *Environ Res* 31:340-354.
- *Johansson A, Curstedt T, Robertson B, et al. 1984. Lung morphology and phospholipids after experimental inhalation of soluble cadmium, copper, and cobalt. *Environ Res* 34:295-309.
- Johns C, Timmerman. 1998. Total cadmium, copper, and zinc in two dreissenid mussels, *Dreissena polymorpha* and *Dreissena bugensis*, at the outflow of Lake Ontario. *J Great Lakes Res* 24(1):55-64.
- *Johnson CA, Sigg L, Zobrist J. 1987. Case studies on the chemical composition of fogwater: The influence of local gaseous emissions. *Atmos Environ* 21(11):2365-2374.
- Johnson PE. 1989. Factors affecting copper absorption in humans and animals. *Adv Exp Med Biol* 258:71-79.
- *Johnson PE, Milne DB, Lykken GI. 1992. Effects of age and sex on copper absorption, biological half-life, and status in humans. *Am J Clin Nutr* 56(5):917-925.
- Johnson PE, Stuart MA, Bowman TD. 1988a. Bioavailability of copper in rats from various foodstuffs and in the presence of different carbohydrates. *Proc Soc Exp Biol Med* 187(1):44-50.
- *Johnson PE, Stuart MA, Hunt JR, et al. 1988b. Copper absorption by women fed intrinsically and extrinsically labeled goose meat, goose liver, peanut butter and sunflower butter. *J Nutr* 118(12):1522-1528.
- *Jolly JL, Edelstein D. 1987. Bureau of mines minerals yearbook-Copper, 1-57.
- Jones MM, Singh PK, Zimmerman LJ, et al. 1995. Effects of some chelating agents on urinary copper excretion by the rat. *Chem Res Toxicol* 8:942-948.
- Junge RE, Thornburg L. 1989. Copper poisoning in four llamas. *J Am Vet Med Assoc* 195(7):987-989.
- Kabala C, Singh BR. 2001. Fractionation and mobility of copper, lead, and zinc in soil profiles in the vicinity of a copper smelter. *J Environ Qual* 30(2):485-492.

9. REFERENCES

- Kahkonen MA, Pantsar-Kallio M, Manninen PKG. 1997. Analysing heavy metal concentrations in the different parts of *Elodea canadensis* and surface sediment with PCA in two boreal lakes in Southern Finland. *Chemosphere* 35(11):2645-2656.
- Kahkonen MA, Suominen KP, Manninen PKG, et al. 1998. 100 Years of sediment accumulation history of organic halogens and heavy metals in recipient and nonrecipient lakes of pulping industry in Finland. *Environ Sci Technol* 32(12):1741-1746.
- Kalac P, Niznanska M, Bevilaqua D, et al. 1996. Concentrations of mercury, copper, cadmium, and lead in fruiting bodies of edible mushrooms in the vicinity of a mercury smelter and a copper smelter. *Sci Total Environ* 177:251-258.
- *Kamamoto Y, Makiura S, Sugihara S, et al. 1973. The inhibitory effects of copper on DL-ethionine carcinogenesis in rats. *Cancer Res* 33:1129-1135.
- Kapur MM, Mokkapatil S, Farooq A, et al. 1984. Copper intravas device (IVD) and male contraception. *Contraception* 29(1):45-54.
- *Karlsson B, Noren L. 1965. Ipecacuanha and copper sulphate as emetics in intoxications in children. *Acta Paediatr Scand* 54:331-335.
- Karthikeyan KG, Elliott HA, Cannon FS. 1997. Adsorption and coprecipitation of copper with the hydrous oxides of iron and aluminum. *Environ Sci Technol* 31:2721-2725.
- Kasama T, Tanaka H. 1989. Effects of oral copper administration to pregnant heterozygous brindled mice on fetal viability and copper levels. *J Nutr Sci Vitaminol* 35(6):627-638.
- Kashulin NA, Ratkin NE, Dauvalter VA, et al. 2001. Impact of airborne pollution on the drainage area of subarctic lakes and fish. *Chemosphere* 42:51-59.
- *Keller C, Vedy J-C. 1994. Heavy metals in the environment-Distribution of copper and cadmium fractions in two forest soils. *J Environ Qual* 23:987-999.
- *Keller JC, Kaminski EJ. 1984. Toxic effects of Cu implants on liver. *Fundam Appl Toxicol* 4:778-783.
- *Kennish MJ. 1998. Trace metal-sediment dynamics in estuaries: Pollution assessment. *Rev Environ Contam Toxicol* 155:69-110.
- Kidwell JM, Phillips LJ, Birchard GF. 1995. Comparative analyses of contaminant levels in bottom feeding and predatory fish using the national contaminant biomonitoring program data. *Bull Environ Contam Toxicol* 54:919-923.
- *Kilbride KM, Paveglio FL, Altstatt AL, et al. 1998. Contaminant loading in drainage and fresh water used for wetland management at Stillwater National Wildlife Refuge. *Arch Environ Contam Toxicol* 35:236-248.
- Kim K-H, Kim D-Y. 1996. Heavy metal pollution in agricultural soils: Measurements in the proximity of abandoned mine land sites (AMLs). *J Environ Sci Health Part A* A31(4):783-795.
- *Kim N, Fergusson J. 1993. Concentrations and sources of cadmium, copper, lead and zinc in house dust in Christchurch, New Zealand. *Sci Total Environ* 138:1-21.

9. REFERENCES

- Kim ND, Fergusson JE. 1994. The concentrations, distribution and sources of cadmium, copper, lead and zinc in the atmosphere of an urban environment. *Sci Total Environ* 144:179-189.
- *Kimball KD. 1973. Seasonal fluctuations of ionic copper in Knights Pond, Massachusetts. *Limnol Oceanogr* 18:169-172.
- *King LD. 1988. Retention of metals by several soils of the southeastern United States. *J Environ Qual* 17(2):239-246.
- King SO, Mach CE, Brezonik PL. 1992. Changes in trace metal concentrations in lake water and biota during experimental acidification of Little Rock Lake, Wisconsin, USA. *Environ Pollut* 78:9-18.
- *Klein D, Scholz P, Drasch GA, et al. 1991. Metallothionein, copper and zinc in fetal and neonatal human liver: changes during development. *Toxicol Lett* 56:61-67.
- Klevay LM. 1998. Lack of recommended dietary allowance for copper may be hazardous to your health. *J Am Coll Nutr* 17(4):322-326.
- *Kline RD, Hays VW, Cromwell GL. 1971. Effects of copper, molybdenum and sulfate on performance, hematology and copper stores of pigs and lambs. *J Anim Sci* 33(4):771-779.
- Klomp AEM, Tops BBJ, van den Berg IET, et al. 2002. Biochemical characterization and subcellular localization of human copper transporter 1 (hCTR1) *Biochem J* 364(2):497-505.
- *Knobeloch L, Schubert C, Hayes J, et al. 1998. Gastrointestinal upsets and new copper plumbing- is there a connection? *Wis Med J* 97(1):49-53.
- Knobeloch L, Ziarnik M, Howard J, et al. 1992. Gastrointestinal upsets associated with ingestion of copper-contaminated water. *Dig Dis Sci* 37(11):1785-1790.
- *Knobeloch L, Ziarnik M, Howard J, et al. 1994. Gastrointestinal upsets associated with ingestion of copper-contaminated water. *Environ Health Perspect* 102:958-961.
- Knudsen E, Sandstrom B, Solgaard P. 1996. Zinc, copper and magnesium absorption from a fibre-rich diet. *J Trace Elem Med Biol* 10:68-76.
- Knulst J, Sodergren A. 1994. Occurrence and toxicity of persistent pollutants in surface microlayers near an incinerator plant. *Chemosphere* 29(6):1339-1347.
- Koch M, Rotard W. 2001. On the contribution of background sources to the heavy metal content of municipal sewage sludge. *Water Sci Technol* 43(2):67-74.
- Komatsu Y, Sadakata I, Ogra Y, et al. 2000. Excretion of copper complexed with thiomolybdate into the bile and blood in LEC rats. *Chem Biol Interact* 124(3):217-231.
- *Komori M, Nishio K, Kitada M, et al. 1990. Fetus-specific expression of a form of cytochrome P-450 in human livers. *Biochemistry* 29:4430-4433.
- Kostova V. 1995. [The neurological screening of workers in the manufacture of copper and aluminum rolled wire]. *Probl Khig* 20:198-209. (Bulgarian)

9. REFERENCES

- *Koutrakis P, Briggs SLK, Leaderer BP. 1992. Source appointment of indoor aerosols in Suffolk and Onondaga Counties, New York. *Environ Sci Technol* 26:521-527.
- Kozanoglou C, Catsiki VA. 1997. Impact of products of a ferronickel smelting plant to the marine benthic life. *Chemosphere* 34(12):2673-2682.
- *Krishnan K, Andersen ME. 1994. Physiologically based pharmacokinetic modeling in toxicology. In: Hayes AW, ed. *Principles and methods of toxicology*. 3rd ed. New York, NY: Raven Press, Ltd., 149-188.
- *Krishnan K, Andersen ME, Clewell HJ III, et al. 1994. Physiologically based pharmacokinetic modeling of chemical mixtures. In: Yang RSH, ed. *Toxicology of chemical mixtures: Case studies, mechanisms, and novel approaches*. San Diego, CA: Academic Press, 399-437.
- Krumgalz BS. 1993. "Fingerprints" approach to the identification of anthropogenic trace metal sources in the nearshore and estuarine environments. *Estuaries* 16(3A):488-495.
- *Kumar A, Sharma CB. 1987. Hematological indices in copper-poisoned rats. *Toxicol Lett* 38:275-278.
- Kumaratilake JS, Howell JM. 1989a. Intracellular distribution of copper in the liver of copper-loaded-sheep - a subcellular fractionation study. *J Comp Pathol* 101(2):161-176.
- Kumaratilake JS, Howell JM. 1989b. Intravenously administered tetra-thiomolybdate and the removal of copper from the liver of copper-loaded sheep. *J Comp Pathol* 101:177-199.
- Kumaratilake JS, Howell JM. 1989c. Lysosomes in the pathogenesis of liver injury in chronic copper poisoned sheep: An ultrastructural and morphometric study. *J Comp Pathol* 100:381-390.
- Kurisaki E, Kuroda Y, Sato M. 1988. Copper-binding protein in acute copper poisoning. *Forensic Sci Int* 38(1-2):3-11.
- *Kust RN. 1978. Copper compounds. In: Kirk-Othmer encyclopedia of chemical technology, Vol 7. 3rd ed. New York, NY: John Wiley & Sons, 97-109.
- Lagos GE, Maggi LC, Peters D, et al. 1999. Model for estimation of human exposure to copper in drinking water. *Sci Total Environ* 239:49-70.
- Lal S, Papeschi R, Duncan RJS, et al. 1974. Effect of copper loading on various tissue enzymes and brain monoamines in the rat. *Toxicol Appl Pharmacol* 28:395-405.
- Lal S, Papeschi R, Duncan RJS. 1983. Trace metals distribution in the surficial sediments of Penobscot Bay, Maine. *Bull Environ Contam Toxicol* 31:566-573.
- Lamb DJ, Avades TY, Allen MD, et al. 2002. Effect of dietary copper supplementation on cell composition and apoptosis in atherosclerotic lesions in cholesterol-fed rabbits. *Atherosclerosis* 164(2):229-236.
- *Lamont DL, Duflou JALC. 1988. Copper sulfate. Not a harmless chemical. *Am J Forensic Med Pathol* 9(3):226-227.

9. REFERENCES

- *Landing WM, Perry JJ Jr, Guentzel JL. 1995. Relationships between the atmospheric deposition of trace elements, major ions, and mercury in Florida: The FAMS Project (1992-1993). *Water Air Soil Pollut* 80:343-352.
- Larsen PF, Gadbois DF, Johnson AC, et al. 1983a. Distribution of polycyclic aromatic hydrocarbons in the surficial sediments of Casco Bay, Maine. *Bull Environ Contam Toxicol* 30:530-535.
- *Larsen PF, Zdanowicz V, Johnson AC. 1983b. Trace metal distribution in the surficial sediments of Penobscot Bay, Maine. *Bull Environ Contam Toxicol* 31:566-573.
- *Lauenstein GG, Daskalakis KD. 1998. U.S. Long-term coastal contaminant temporal trends determined from mollusk monitoring programs, 1965-1993. *Mar Pollut Bull* 37(1-2):6-13.
- *Law LW. 1938. The effects of chemicals on the lethal mutation rate in *Drosophila melanogaster*. *Proc Natl Acad Sci USA* 24:546-550.
- *Lecyk M. 1980. Toxicity of CuSO_4 in mice embryonic development. *Zool Pol* 28:101-105.
- Lee D-Y, Brewer GJ, Wang Y. 1989. Treatment of Wilson's disease with zinc. VII. Protection of the liver from copper toxicity by zinc-induced metallothionein in a rat model. *J Lab Clin Med* 114(6):639-645.
- Lee DY, Schroeder K, Gordon DT. 1988. Enhancement of Cu bioavailability in the rat by phytic acid. *J Nutr* 118(6):712-717.
- *Leeder JS, Kearns GL. 1997. Pharmacogenetics in pediatrics: Implications for practice. *Pediatr Clin North Am* 44(1):55-77.
- *Leung H-W. 1993. Physiologically-based pharmacokinetic modelling. In: Ballentine B, Marro T, Turner P, eds. *General and applied toxicology*. Vol. 1. New York, NY: Stockton Press, 153-164.
- Levenson CW. 1998. Mechanisms of copper conservation in organs. *Am J Clin Nutr* 67:978S-981S.
- *Levenson CW, Janghorbani M. 1994. Long-term measurement of organ copper turnover in rats by continuous feeding of a stable isotope. *Anal Biochem* 221(2):243-249.
- *Levy DB, Barbarick KA, Siemer EG. 1992. Distribution and partitioning of trace metals in contaminated soils near Leadville, Colorado. *J Environ Qual* 21:185-195.
- *Lewis RJ, ed. 1997. *Hawley's condensed chemical dictionary*. New York, NY: John Wiley & Sons, Inc., 297, 302.
- *Lewis RJ. 2000. *Sax's dangerous properties of industrial materials*. 10th edition. New York: John Wiley & Sons, Inc., 990.
- Lichte FE, Seeley JL, Jackson LL, et al. 1987. Geological and inorganic materials. *Anal Chem* 59:197R-212R.
- *Lide DR, ed. 2000. *CRC handbook of chemistry and physics*. New York, NY: CRC Press, 4-56, 4-58.

9. REFERENCES

- *Lind Y, Glynn AW. 1999. Intestinal absorption of copper from drinking water containing fulvic acids and an infant formula mixture studied in a suckling rat model. *BioMetals* 12:181-187.
- Linde AR, Sanchez-Calan S, Izquierdo JI, et al. 1998. Brown trout as biomonitor of heavy metal pollution: Effect of age on the reliability of the assessment. *Ecotoxicol Environ Saf* 40:120-125.
- Lindeman JH, Lentjes EG, Berger HM. 1995. Diminished protection against copper-induced lipid peroxidation by cord blood plasma of preterm and term infants. *J Parenter Enteral Nutr* 19(5):373-375.
- Linder MC, Hazegh-Azam M. 1996. Copper biochemistry and molecular biology. *Am J Clin Nutr* 63:797S-811S.
- Linder MC, Wooten L, Cerveza P, et al. 1998. Copper transport. *Am J Clin Nutr* 67:965S-971S.
- Linder MC, Zerounian NR, Moriya M, et al. 2003. Iron and copper homeostasis and intestinal absorption using the Caco2 cell model. *Biometals* 16:145-160.
- *Lioy PJ, Daisey JM, Morandi MT, et al. 1987. The airborne toxic element and organic substances (ATEOS) study design. In: Lioy PJ, Daisey JM, eds. *Toxic air pollution: A comprehensive study of non-criteria air pollutants*. Chelsea, MI: Lewis Publishing, Inc., 3-42.
- *Lisk DJ, Gutenmann WH, Rutzke M, et al. 1992. Survey of toxicants and nutrients in composted waste materials. *Arch Environ Contam Toxicol* 22:190-194.
- Liu C, Jiao K. 1990. Ultraviolet irradiation and polarographic adsorptive complex wave techniques for the simple and rapid simultaneous determination of trace amounts of zinc, lead and copper in human hair. *Anal Chim Acta* 238(2):367-374.
- Liu CF, Medeiros DM. 1986. Excess diet copper increases systolic blood pressure in rats. *Biol Trace Elem Res* 9:15-24.
- Liu J, Kashimura S, Hara K, et al. 2001. Death following cupric sulfate emesis. *J Toxicol Clin Toxicol* 39(2):161-163.
- *Livingston, AL. 1978. Forage plant estrogens. *J Toxicol Environ Health* 4:301-324.
- *Llanos RM, Mercer JFB. 2002. The molecular basis of copper homeostasis and copper-related disorders. *DNA Cell Biol* 21(4):259-279.
- *Llewellyn GC, Floyd EA, Hoke GD, et al. 1985. Influence of dietary aflatoxin, zinc, and copper on bone size, organ weight, and body weight in hamsters and rats. *Bull Environ Contam Toxicol* 35:149-156.
- *Lo FB, Araki DK. 1989. Biological monitoring of toxic metals in urine by simultaneous inductively coupled plasma-atomic emission spectrometry. *Am Ind Hyg Assoc J* 50(5):245-251.
- *Lodenius M, Braunschweiler H. 1986. Volatilisation of heavy metals from a refuse dump. *Sci Total Environ* 57:253-255.
- Loehr RC, Rogers LA, Erickson DC. 1992. Mobility of residues at petroleum industry hazardous waste land treatment sites. *Water Sci Technol* 25(3):191-196.

9. REFERENCES

- *Long DT, Angino EE. 1977. Chemical speciation of Cd, Cu, Pb, and Zn in mixed freshwater, seawater, and brine solutions. *Geochim Cosmochim Acta* 41:1183-1191.
- Longerich HP, Friel JK, Fraser C, et al. 1991. Analysis of drinking water of mothers of neural tube defect infants and of normal for 14 selected trace elements by inductively coupled plasma-mass spectrometry (ICP-MS). *Can J Appl Spectrosc* 36(1):15-21.
- Lonnerdal B. 1996. Bioavailability of copper. *Am J Clin Nutr* 63(5):8215-8295.
- Lonnerdal B. 1998. Copper nutrition during infancy and childhood. *Am J Clin Nutr* 67(5):10465-10535.
- *Lopez-Artiguez M, Camean A, Repetto M. 1993. Preconcentration of heavy metals in urine and quantification by inductively coupled plasma atomic emission spectrometry. *J Anal Toxicol* 17(1):18-22.
- Lopez-Artiguez M, Camean AM, Repetto M. 1996. Determination of nine elements in sherry wine by inductively coupled plasma-atomic emission spectrometry. *J AOAC Int* 79(5):1191-1197.
- Lopez-Artiguez M, Grilo A, Soria M-L, et al. 1990. Levels of zinc, copper, and lead in wines from the area south of Seville. *Bull Environ Contam Toxicol* 45:711-717.
- *Loranger S, Tetrault M, Kennedy G, et al. 1996. Manganese and other trace elements in urban snow near an expressway. *Environ Pollut* 92(2):203-211.
- *Lores EM, Pennock JR. 1998. The effect of salinity on binding of Cd, Cr, Cu and Zn to dissolved organic matter. *Chemosphere* 37(5):861-874.
- *Lowe TP, May TW, Brumbaugh WG. 1985. National contaminant biomonitoring program: Concentration of seven elements in freshwater fish, 1978-1981. *Arch Environ Contam Toxicol* 14:363-388.
- Lubach VD, Wurzinger R. 1986. [Trace elements in brittle nails.] *Derm Beruf Umwelt* 34(2):37-39. (German)
- *Luncan-Bouché ML, Couderchet M, Vernet G, et al. 1997. The simultaneous influence of pH and temperature on binding and mobilization of metals in sand: 1-Copper. *Fresenius Environ Bull* 6:711-718.
- Lundborg M, Camner P. 1984. Lysozyme levels in rabbit lung after inhalation of nickel, cadmium, cobalt, and copper chlorides. *Environ Res* 34:335-342.
- *Ma H, Kim SD, Allen HE, et al. 2002. Effect of copper binding by suspended particulate matter on toxicity. *Environ Toxicol Chem* 21(4):710-714.
- *Ma J, Betts NM5. 2000. Zinc and copper intakes and their major food sources for older adults in the 1994-1996 continuing survey of food intakes by individuals (CSFII). *J Nutr* 130(11):2838-2843.
- *Ma LQ, Rao GN. 1997. Heavy metals in the environment-chemical fractionation of cadmium, copper, nickel, and zinc in contaminated soils. *J Environ Qual* 26:259-264.

9. REFERENCES

- *Ma LQ, Tan F, Harris WG. 1997. Concentrations and distributions of eleven metals in Florida soils. *J Environ Qual* 26:769-773.
- *MacCarthy P, Klusman RW. 1987. Water analysis. *Anal Chem* 59:308R-337R.
- *Maessen O, Freedman B, McCurdy R. 1985. Metal mobilization in home well water systems in Nova Scotia. *J Am Water Works Assoc* 77:73-80.
- *Magee AC, Matrone G. 1960. Studies on growth, copper metabolism and iron metabolism of rats fed high levels of zinc. *J Nutr* 72:233-242.
- Maggiore G, De Giacomo C, Sessa F, et al. 1987. Idiopathic hepatic copper toxicosis in a child. *J Pediatr Gastroenterol Nutr* 6:980-983.
- *Makale MT, King GL. 1992. Surgical and pharmacological dissociation of cardiovascular and emetic responses to intragastric CuSO₄. *Am J Physiol* 263(2 Pt 2):R284-R291.
- *Mannsville Chemical Products. 1984. Copper sulfate. *Chemical products synopsis*: 115-117.
- *Marceau N, Aspin N, Sass-Kortsak A. 1970. Absorption of copper 64 from gastrointestinal tract of the rat. *Am J Physiol* 218(2):377-383.
- Marcisz C, Jonderko G, Wieczorek-Latka U, et al. 1998. Respiratory system of workers dealing with casting and processing of copper. *Pneumol Alergol Pol* 66(9-10):433-439.
- *Marinussen PJC, van der Zee SEATM, de Haan F, et al. 1997. Heavy metal (copper, lead, and zinc) accumulation and excretion by the earthworm, *Dendrobaena veneta*. *J Environ Qual* 26:278-284.
- *Mart L, Nurnberg HW. 1984. Trace metal levels in the Eastern Arctic Ocean. *Sci Total Environ* 39:1-14.
- Martens D, Balta-Brouma K, Brotsack R, et al. 1998. Chemical impact of uncontrolled solid waste combustion to the vicinity of the Kourpoupitos Ravine, Crete, Greece. *Chemosphere* 36(14):2855-2866.
- *Martin M, Castle W. 1984. Petrowatch: Petroleum hydrocarbons, synthetic organic compounds, and heavy metals in mussels from the Monterey Bay area of central California. *Mar Pollut Bull* 15:259-266.
- Martinez CE, McBride MB. 1999. Dissolved and labile concentrations of Cd, Cu, Pb, and Zn in aged ferrihydrite-organic matter systems. *Environ Sci Technol* 33:745-750.
- Marx J. 2003. Possible role of environmental copper in Alzheimer's disease. *Science* 301(5635):905.
- *Marzin DR, Phi HV. 1985. Study of the mutagenicity of metal derivatives with *Salmonella typhimurium*. *Mutat Res* 155:49-51.
- *Mason KE. 1979. A conceptus of research on copper metabolism and requirements of man. *J Nutr* 109:1979-2066.
- Mason RW, Edwards IR, Fisher LC. 1989. Teratogenicity of combinations of sodium dichromate, sodium arsenate and copper sulphate in the rat. *Comp Biochem Physiol C* 93(2):407-411.

9. REFERENCES

- *Massie HR, Aiello VR. 1984. Excessive intake of copper: Influence on longevity and cadmium accumulation in mice. *Mech Ageing Dev* 26:195-203.
- Mattie MD, Freedman JH. 2001. Protective effects of aspirin and vitamin E (α -Tocopherol) against copper- and cadmium-induced toxicity. *Biochem Biophys Res Commun* 285(4):921-925.
- *May TW, Wiedmeyer RH, Gober J, et al. 2001. Influence of mining-related activities on concentrations of metals in water and sediment from streams of the Black Hills, South Dakota. *Arch Environ Contam Toxicol* 40:1-9.
- *Mayr U, Butsch A, Schneider S. 1992. Validation of two in vitro test systems for estrogenic activities with zearalenone, phytoestrogens and cereal extracts. *Toxicology* 74:135-149.
- McArdle HJ, Erlich R. 1991. Copper uptake and transfer of the mouse fetus during pregnancy. *J Nutr* 121(2):208-214.
- McArdle HJ, Gross SM, Danks DM. 1988. Uptake of copper by mouse hepatocytes. *J Cell Physiol* 136(2):373-378.
- McArdle HJ, Gross SM, Danks DM, et al. 1990. Role of albumin's copper binding site in copper uptake by mouse hepatocytes. *Am J Physiol* 258:G988-G991.
- McArdle HJ, Guthrie JR, Ackland ML, et al. 1987. Albumin has no role in the uptake of copper by human fibroblasts. *J Inorg Biochem* 31:123-131.
- *McCrary JK, Chapman GA. 1979. Determination of copper complexing capacity of natural river water, well water and artificially reconstituted water. *Water Res* 13:143-150.
- *McIlroy LM, DePinto JV, Young TC, et al. 1986. Partitioning of heavy metals to suspended solids of the Flint River, Michigan. *Environ Toxicol Chem* 5:609-623.
- *Mehra RK, Bremner I. 1984. Species differences in the occurrence of copper-metallothionein in the particulate fractions of the liver of copper-loaded animals. 219:539-546.
- *Meister RT, Sharp DT, Ponkivar E, et al., eds. 2001. *Farm chemicals handbook*, Volume 87. Willoughby, OH: Meister Publishing Co., C106-107.
- *Meranger JC, Subramanian KS, Chalifoux C. 1979. A national survey for cadmium, chromium, copper, lead, zinc, calcium, and magnesium in Canadian drinking water supplies. *Environ Sci Technol* 13:707-711.
- *Mercer JFB, Lazdins I, Stevenson T, et al. 1981. Copper induction of translatable metallothionein messenger RNA. *Biosci Rep* 1:793-800.
- Michael GE, Miday RK, Bercz JP, et al. 1981. Chlorine dioxide water disinfection: A prospective epidemiology study. *Arch Environ Health* 36:20-27.
- *Miller DR, Byrd JE, Perona MJ. 1987. The source of Pb, Cu, and Zn in fogwater. *Water Air Soil Pollut* 32:329-340.

9. REFERENCES

- Miller GD, Keen CL, Stern JS, et al. 1996. Copper absorption, endogenous excretion, and distribution in Sprague-Dawley and lean (Fa/Fa) Zucker rats. *Biol Trace Elem Res* 53:261-279.
- *Mills GL, Quinn JG. 1984. Dissolved copper and copper-organic complexes in the Narragansett Bay estuary. *Mar Chem* 15:151-172.
- *Minear RA, Ball RO, Church RL. 1981. Data base for influent heavy metals in publicly owned treatment works. 1-4. EPA600S281220.
- Miszta H. 1989. *In vitro* effect of copper on the stromal cells of bone marrow in rats. *Toxicol Ind Health* 5(6):1117-1123.
- *Moffett JW, Zika RG. 1987. Photochemistry of copper complexes in sea water. *Photochemistry of environmental aquatic systems*. ACS Symp Ser 327:116-130.
- Moreno MA, Marin C, Vinagre F, et al. 1999. Trace element levels in whole blood samples from residents of the city Badajoz, Spain. *Sci Total Environ* 229:209-215.
- Moriya M, Ohta T, Watanabe K, et al. 1983. Further mutagenicity studies on pesticides in bacterial reversion assay systems. *Mutat Res* 116:185-216.
- Morselli L, Zappoli S, Tirabassi T. 1992. Characterization of the effluents from a municipal solid waste incinerator plant and of their environmental impact. *Chemosphere* 24(12):1775-1784.
- *Morselli PL, Franco-Morselli R, Bossi L. 1980. Clinical pharmacokinetics in newborns and infants: Age-related differences and therapeutic implications. *Clin Pharmacokin* 5:485-527.
- Mudroch A. 1993. Lake Ontario sediments in monitoring pollution. *Environ Monit Assess* 28:117-129.
- *Müller T, Feichtinger H, Berger H, et al. 1996. Endemic tyrolean infantile cirrhosis: an ecogenetic disorder. *Lancet* 347:887-880.
- *Müller T, Müller W, Feichtinger H. 1998. Idiopathic copper toxicosis. *Am J Clin Nutr* 67:1082S-1086S.
- Müller T, Schaefer H, Rodeck B, et al. 1999. Familial clustering of infantile cirrhosis in Northern Germany: A clue to the etiology of idiopathic copper toxicosis. *J Pediatr* 135(2 Pt 1):189-196.
- *Müller-Höcker J, Meyer U, Wiebecke B, et al. 1988. Copper storage disease of the liver and chronic dietary copper intoxication in two further German infants mimicking Indian Childhood Cirrhosis. *Pathol Res Pract* 183(1):39-45.
- Müller-Höcker J, Summer KH, Schramel P, et al. 1998. Different pathomorphologic patterns in exogenic infantile copper intoxication of the liver. *Pathol Res Pract* 194(6):377-384.
- Mullins JE, Fuentealba IC. 1998. Immunohistochemical detection of metallothionein in liver, duodenum and kidney after dietary copper-overload in rats. *Histol Histopathol* 13:627-633.
- *Mullins MJP, Norman JB. 1994. Solubility of metals in windblown dust from mine waste dump sites. *Appl Occup Environ Hyg* 9(3):218-223.

9. REFERENCES

- *Mumma RO, Raupach DC, Waldman JP, et al. 1984. National survey of elements and other constituents in municipal sewage sludges. *Arch Environ Contam Toxicol* 13:75-83.
- *Murphy EA. 1993. Effectiveness of flushing on reducing lead and copper levels in school drinking water. *Environ Health Perspect* 101(3):240-241.
- *Murthy RC, Lal S, Saxena DK, et al. 1981. Effect of manganese and copper interaction on behavior and biogenic amines in rats fed a 10% casein diet. *Chem Biol Interact* 37:299-308.
- *Musci G, Bonaccorsi di Patti MC, Calabrese L. 1993. The state of copper sites in human ceruloplasmin. *Arch Biochem Biophys* 306:111-118.
- *Mussalo-Rauhamaa H, Salmela SS, Leppanen A, et al. 1986. Cigarettes as a source of some trace and heavy metals and pesticides in man. *Arch Environ Health* 41(1):49-55.
- *Myers BM, Prendergast FG, Holman R, et al. 1993. Alterations in hepatocyte lysosomes in experimental hepatic copper overload in rats. *Gastroenterology* 105(6):1814-1823.
- Nair J, Sone H, Nagao M, et al. 1996. Copper-dependent formation of miscoding etheno-DNA adducts in the liver of long evans cinnamon (LEC) rats developing hereditary hepatitis and hepatocellular carcinoma. *Cancer Res* 56:1267-1271.
- *NAS. 1980. Copper. Recommended dietary allowances. Washington, DC: National Academy of Sciences, 151-154.
- Nash TH, Gries C. 1995. The use of lichens in atmospheric deposition studies with an emphasis on the Arctic. *Sci Total Environ* 160/161:729-736.
- *NAS. 2000. Copper in drinking water. Prepared by the Board of Environmental Studies and Toxicology, Commission on Life Sciences, National Research Council. Washington, DC: National Academy Press.
- *NAS/NRC. 1989. Report of the oversight committee. In: *Biologic markers in reproductive toxicology*. Washington, DC: National Academy of Sciences, National Research Council, National Academy Press.
- NATICH. 1988. NATICH data base report on state, local and EPA air toxics activities. Research Triangle Park, NC. National Air Toxics Information Clearinghouse, U.S. Environmental Protection Agency.
- *NCI. 1968. Evaluation of carcinogenic, teratogenic, and mutagenic activities of selected pesticides and industrial chemicals1. Vol 1. Bethesda, MD. Bionetics Research Laboratories, National Cancer Institute. 1-150. NCI-DCCP-CG-1973-1-1. PB223159.
- Nellessen JE, Fletcher JS. 1993. Assessment of published literature on the uptake, accumulation, and translocation of heavy metals by vascular plants. *Chemosphere* 27(9):1669-1680.
- *Nerin C, Domeno C, Garcia JI, et al. 1999. Distribution of Pb, V, Cr, Ni, Cd, Cu and Fe in particles formed from the combustion of waste oils. *Chemosphere* 38(7):1533-1540.
- *Neuhauser EF, Cukic ZV, Malecki MR, et al. 1995. Bioconcentration and biokinetics of heavy metals in the earthworm. *Environ Pollut* 89(3):293-301.

9. REFERENCES

- *Nicholas PO, Brist MB. 1968. Food-poisoning due to copper in the morning tea. *Lancet* 2:40-42.
- NIOSH. 1985. NIOSH pocket guide to chemical hazards. Washington, DC; National Institute for Occupational Safety and Health.
- *NIOSH. 1987. In: Eller PM, ed. NIOSH manual of analytical methods. Cincinnati, OH; National Institute for Occupational Safety and Health.
- *NIOSH. 1988. National occupational exposure survey as of 05/10/88. National Institute for Occupational Safety and Health.
- *NIOSH. 2002. Copper. NIOSH pocket guide to chemical hazards. National Institute for Occupational Safety and Health. <http://www.cdc.gov/niosh/npg/npgd0150>.
- *Nishioka H. 1975. Mutagenic activities of metal compounds in bacteria. *Mutat Res* 31:185-189.
- *Nolte J. 1988. Pollution source analysis of river water and sewage sludge. *Environ Technol Lett* 9:857-868.
- Nomiyama K, Nomiyama H, Kameda N, et al. 1999. Mechanism of hepatorenal syndrome in rats of Long-Evans Cinnamon strain, an animal model of fulminant Wilson's disease. *Toxicology* 132(2-3):201-214.
- Norrstrom AC, Jacks G. 1998. Concentration and fractionation of heavy metals in roadside soils receiving de-icing salts. *Sci Total Environ* 218:161-174.
- Nowack B, Xue H, Sigg L. 1997. Influence of natural and anthropogenic ligands on metal transport during infiltration of river water to groundwater. *Environ Sci Technol* 31:866-872.
- *NRC. 1977. Medical and biologic effects of environmental pollutants: Copper. Washington, DC: National Research Council. National Academy of Sciences.
- *NRC. 1993. Pesticides in the diets of infants and children. Washington, DC; National Research Council. National Academy Press.
- *NRC. 1995. Nutrient requirements of laboratory animals, fourth revised edition. Subcommittee on Laboratory Animal Nutrition, Committee on Animal Nutrition, Board of Agriculture, National Research Council. Washington, DC: National Academy Press.
- NRC. 2000. Copper in drinking water. Washington, DC: National Research Council, National Academy Press.
- *Nriagu JO. 1989. A global assessment of natural sources of atmospheric trace metals. *Nature* 338:47-49.
- *Nriagu JO, Coker RD. 1980. Trace metals in humic and fulvic acids from Lake Ontario sediments. *Environ Sci Technol* 14(:):443-446.
- *Nriagu JO, Pacyna JM. 1988. Quantitative assessment of worldwide contamination of air, water and soil by trace metals. *Nature* 333:134-139.

9. REFERENCES

- *Nriagu JO, Lawson G, Wong HKT, et al. 1996. Dissolved trace metals in Lakes Superior, Erie, and Ontario. *Environ Sci Technol* 30:178-187.
- *NTP. 1993. NTP Technical Report on toxicity studies of cupric sulfate administered in drinking water and feed to F344/N rats and B6C3F₁ mice. National Toxicology Program. United States Department of Health and Human Services. NIH Publication 93-3352.
- Nyberg P, Gottfries C-G, Holmgren G, et al. 1982. Advanced catecholaminergic disturbances in the brain in a case of Wilson's disease. *Acta Neurol Scand* 65:71-75.
- *Nygren O, Nilsson C-A, Lindahl R. 1992. Occupational exposure to chromium, copper and arsenic during work with impregnated wood in joinery shops. *Ann Occup Hyg* 36(5):509-517.
- Obata H, Sawada N, Isomura H, et al. 1996. Abnormal accumulation of copper in LEC rat liver induces expression of p53 and nuclear matrix-bound p21^{waf1/cip1}. *Carcinogenesis* 17(10):2157-2161.
- O'Day PA, Carroll SA, Randall S, et al. 2000. Metal speciation and bioavailability in contaminated estuary sediments, Alameda Naval Air Station, California. *Environ Sci Technol* 34:3665-3673.
- *O'Dell BL. 1984. Copper. Present knowledge in nutrition. The Nutrition Foundation, Inc. Washington, DC.
- *O'Donohue JW, Reid MA, Varghese A, et al. 1993. Micronodular cirrhosis and acute liver failure due to chronic copper self-intoxication. *Eur J Gastroenterol Hepatol* 5(7):561-562.
- O'Donohue JW, Reid MA, Varghese A, et al. 1999. A case of adult chronic copper self-intoxication resulting in cirrhosis. *Eur J Med Res* 4:252.
- *Ogra Y, Ohmichi M, Suzuki KT. 1996. Mechanisms of selective copper removal by tetrathiomolybdate from metallothionein in LEC rats. *Toxicology* 106(1-3):75-83.
- Okayasu T, Tochimara H, Hyuga T, et al. 1992. Inherited copper toxicity in Long-Evans Cinnamon rats exhibiting spontaneous hepatitis: A model of Wilson's disease. *Pediatr Res* 31(3):253-257.
- *Okonkwo AC, Ku PK, Miller ER, et al. 1979. Copper requirement of baby pigs fed purified diets. *J Nutr* 109:939-948.
- Oldenquist G, Salem M. 1999. Parenteral copper sulfate poisoning causing acute renal failure. *Nephrol Dial Transplant* 14(2):441-443.
- *Olivares M, Araya M, Pizarro F, et al. 2001. Nausea threshold in apparently healthy individuals who drink fluids containing graded concentrations of copper. *Regul Toxicol Pharmacol* 33(3):271-275.
- *Olivares M, Araya M, Uauy R. 2000. Copper homeostasis in infant nutrition: Deficit and excess. *J Pediatr Gastroenterol Nutr* 31(2):102-111.
- *Olivares M, Lonnerdal B, Abrams SA, et al. 2002. Age and copper intake do not affect copper absorption, measured with the use of ⁶⁵Cu as a tracer, in young infants. *Am J Clin Nutr* 76(3):641-645.

9. REFERENCES

- *Olivares M, Pizarro F, Speisky H, et al. 1998. Copper in infant nutrition: safety of World Health Organization provisional guideline value for copper content of drinking water. *J Pediatr Gastroenterol Nutr* 26:251-257.
- Olmez I, Kotra JP, Lowery S, et al. 1985. Airborne lead and trace elements in an indoor shooting range: A study of the DC National Guard armory pistol range. *Environ Toxicol Chem* 4:447-452.
- Olmez I, Sheffield AE, Gordon GE, et al. 1988. Compositions of particles from selected sources in Philadelphia for receptor modeling applications. *J Air Pollut Control Assoc* 38(11):1392-1402.
- Ong CN, Chia SE, Foo SC, et al. 1993. Concentrations of heavy metals in maternal and umbilical cord blood. *BioMetals* 6:61-66.
- OSHA. 1985. Permissible exposure limits (Table Z-1). Occupational Safety and Health Administration. CFR 29(1910.1000):655-659.
- *OSHA. 2002a. Air contaminants. Occupational safety and health standards for shipyard employment. Occupational Safety and Health Administration. Code of Federal Regulations. 29 CFR 1910.119. <http://www.osha.gov/pls/oshaweb/owa>. April 11, 2002.
- *OSHA. 2002b. Gases, vapors, fumes, dusts, and mists. Safety and health regulations for construction. Occupational Safety and Health Administration. Code of Federal Regulations. 29 CFR 1926.55. <http://www.osha.gov/pls/oshaweb/owa>. April 09, 2002.
- *OSHA. 2002c. Limits for air contaminants. Occupational safety and health standards. Occupational Safety and Health Administration. Code of Federal Regulations. 29 CFR 1910.1000. <http://www.osha.gov/pls/oshaweb/owa>. April 09, 2002.
- O'Shea KS, Kaufman MH. 1979. Influence of copper on the early post-implantation mouse embryo: An in vivo and in vitro study. *Wilhelm Roux's Arch Dev Biol* 186:297-308.
- O'Shea KS, Kaufman MH. 1980. Copper-induced microtubule degeneration and filamentous inclusions in the neuroepithelium of the mouse embryo. *Acta Neuropathol* 49:237-240.
- Ostapczuk P, Burow M, May K, et al. 1997. Mussels and algae as bioindicators for long-term tendencies of element pollution in marine ecosystems. *Chemosphere* 34(9/10):2049-2058.
- Otero XL, Sanchez JM, Macias F. 2000. Bioaccumulation of heavy metals in thionic fluvisols by a marine polycheate: The role of metal studies. *J Environ Qual* 29:1133-1141.
- Othman I, Spyrou NM. 1980. The abundance of some elements in hair and nail from the Machakos District of Kenya. *Sci Total Environ* 16:267-278.
- Overvad K, Wang DY, Olsen J, et al. 1993. Copper in human mammary carcinogenesis: A case-cohort study. *Am J Epidemiol* 137(4):409-414.
- *Owen CA. 1965. Metabolism of radiocopper (Cu^{64}) in the rat. *Am J Physiol* 209:900-904.
- *Owen GM, Brozek J. 1966. Influence of age, sex and nutrition on body composition during childhood and adolescence. In: Falkner F, ed. *Human development*. Philadelphia, PA: WB Saunders, 222-238.

9. REFERENCES

- Pacakova V, Pockeviciute D, Armalis S, et al. 2000. A study of the distribution of lead, cadmium and copper between water and kaolin, bentonite and a river sediment. *J Environ Monitor* 2:187-191.
- *Page WG. 1981. Comparison of groundwater and surface water for patterns and levels of contamination by toxic substances. *Environ Sci Technol* 15(12):1475-1480.
- *Pandit A, Bhawe S. 1996. Present interpretation of the role of copper in Indian childhood cirrhosis. *Am J Clin Nutr* 63(5):830S-835S.
- Pang Y, MacIntosh DL, Ryan B. 2001. A longitudinal investigation of aggregate oral intake of copper. *J Nutr* 131(8):2171-2176.
- *Paode RD, Sofuoglu SC, Sivadechathep J, et al. 1998. Dry deposition fluxes and mass size distributions of Pb, Cu, and Zn measured in southern Lake Michigan during AEOLOS. *Environ Sci Technol* 32:1629-1635.
- Paris I, Dagnino-Subiabre A, Marcelain K, et al. 2001. Copper neurotoxicity is dependent on dopamine-mediated copper uptake and one-electron reduction of aminochrome in a rat substantia nigra neuronal cell line. *J Neurochem* 77(2):519-529.
- Parmer P, Daya S. 2001. The effect of copper on (³H)-tryptophan metabolism in organ cultures of rat pineal glands. *Metab Brain Dis* 16(3/4):199-205.
- Parmer P, Limson J, Nyokong T, et al. 2002. Melatonin protects against copper-mediated free radical damage. *J Pineal Res* 32(4):237-242.
- Parrish CS, Urchrin CG. 1990. Runoff-induced metals in Lakes Bay, New Jersey. *Environ Toxicol Chem* 9:559-567.
- Pascoe GA, Blanchet RJ, Linder G. 1994. Bioavailability of metals and arsenic to small mammals at a mining waste-contaminated wetland. *Arch Environ Contam Toxicol* 27(1):44-50.
- Pascoe GA, Blanchet RJ, Linder G. 1996. Food chain analysis of exposures and risks to wildlife at metals-contaminated wetland. *Arch Environ Contam Toxicol* 30:306-318.
- Pashalidis I, Kontoghiorghes GJ. 2002. Molecular factors affecting the complex formation between deferiprone (L1) and Cu (II) *Arzneim Forsch* 51(12):998-1003.
- Paulson AJ, Gendron JF. 2001. Partitioning of copper at concentrations below the marine water quality criteria. *Environ Toxicol Chem* 20(5):952-959.
- *Pekelharing HLM, Lemmens Ag, Beynen AC. 1994. Iron, copper and zinc status in rats fed on diets containing various concentrations of tin. *Br J Nutr* 71:103-109.
- *Pena MMO, Lee J, Thiele DJ. 1999. A delicate balance: Homeostatic control of copper uptake and distribution. *J Nutr* 129(7):1251-1260.
- *Pennington JAT. 1983. Revision of the Total Diet Study food list and diets. *J Am Diet Assoc* 82:166-173.

9. REFERENCES

- *Pennington JAT, Schoen SA. 1996. Contributions of food groups to estimated intakes of nutritional elements: Results from the FDA total diet studies, 1982-1991. *Int J Vitam Nutr Res* 66(4):342-349.
- Pennington JAT, Schoen SA. 1997. Corrections-Contributions of food groups to estimated intakes of nutritional elements: Results from the FDA total diet studies, 1982-1991. *Int J Vitam Nutr Res* 67(3):350-362.
- Pennington JAT, Capar S, Parfitt C, et al. 1996. History of the Food and Drug Administration's total diet study (part II), 1987-1993. *J Assoc Off Anal Chem* 79(1):163-170.
- *Pennington JAT, Young BE, Wilson DB, et al. 1986. Mineral content of foods and total diets: The selected minerals in foods survey, 1982-1984. *J Am Diet Assoc* 86:876-891.
- *Percival SS, Harris ED. 1989. Ascorbate enhances copper transport from ceruloplasmin into human K562 cells. *J Nutr* 119:779-784.
- *Percival SS, Harris ED. 1990. Copper transport from ceruloplasmin: Characterization of the cellular uptake mechanism. *Am J Physiol* 258:C140-C146.
- *Perwak J, Bysshe S, Goyer M, et al. 1980. An exposure and risk assessment for copper. Washington, DC: EPA. EPA-440/4-81-015.
- *Petruzzelli G. 1997. Soil sorption of heavy metals. Chapter 5. In: Ecological issues and environmental impact assessment, 145-175.
- *Petruzzelli G, Lubrano L, Petronio BM, et al. 1994. Soil sorption of heavy metals as influenced by sewage sludge addition. *J Environ Sci Health Part A* A29(1):31-50.
- Petta CA, Faundes D, Pimentel E, et al. 1996. The use of vaginal ultrasound to identify copper T IUDs at high risk of expulsion. *Contraception* 54:287-289.
- Pettersson R, Kjellman B. 1989. Vomiting and diarrhea are the most common symptoms in children who drink water with high levels of copper. *Lakartidningen* 86(25):2361-2362. (Swedish)
- Pettersson R, Rasmussen F, Oskarsson A. 2003. Copper in drinking water: not a strong risk factor for diarrhoea among young children. A population-based study from Sweden. *Acta Paediatr* 92:473-480.
- Phaneuf D, Cote I, Dumas P, et al. 1999. Evaluation of the contamination of marine algae (seaweed) from the St. Lawrence River and likely to be consumed by humans. *Environ Res* 80:S175-S182.
- Pickart L, Freedman JH, Loker WJ, et al. 1980. Growth modulating plasma tripeptide may function by facilitating copper uptake into cells. *Nature* 288:715-717.
- *Pimentel JC, Marquez F. 1969. 'Vineyard sprayer's lung': a new occupational disease. *Thorax* 24:678-688.
- *Pimentel JC, Menezes AP. 1975. Liver granulomas containing copper in Vineyard Sprayer's lung: A new etiology of hepatic granulomatosis. *Am Rev Respir Dis* 3:189-195.
- *Pinochet H, De Gregori I, Lobos MG, et al. 1999. Selenium and copper in vegetables and fruits grown on long-term impacted soils from Valparaiso region, Chile. *Bull Environ Contam Toxicol* 63:327-334.

9. REFERENCES

- Pip E. 1991. Cadmium, copper and lead in soils and garden produce near a metal smelter at Flin Flon, Manitoba. *Bull Environ Contam Toxicol* 46:790-796.
- Pip E. 1993. Cadmium, copper and lead in wild rice from central Canada. *Arch Environ Contam Toxicol* 24:179-181.
- *Pirof F, Millet J, Kalia YN, et al. 1996b. In vitro study of percutaneous absorption, cutaneous bioavailability and bioequivalence of zinc and copper from five topical formulations. *Skin Pharmacol* 9:259-269.
- *Pirof F, Panisset F, Agache P, et al. 1996a. Simultaneous absorption of copper and zinc through human skin in vitro. *Skin Pharmacol* 9:43-52.
- *Pirrone N, Keeler GJ. 1993. Deposition of trace metals in urban and rural areas in the Lake Michigan basin. *Water Sci Technol* 28(3-5):261-270.
- *Pitt R, Field R, Lalor M, et al. 1995. Urban stormwater toxic pollutants: Assessment, sources, and treatability. *Water Environ Res* 67(3):260-275.
- *Pizarro F, Olivares M, Araya M, et al. 2001. Gastrointestinal effects associated with soluble and insoluble copper in drinking water. *Environ Health Perspect* 109(9):949-952.
- *Pizarro F, Olivares M, Uauy R, et al. 1999. Acute gastrointestinal effects of graded levels of copper in drinking water. *Environ Health Perspect* 107(2):117-121.
- *Plamenac P, Santic Z, Nikulin A, et al. 1985. Cytologic changes of the respiratory tract in vineyard spraying workers. *Eur J Respir Dis* 67:50-55.
- Plette ACC, Nederlof MM, Temminghoff EJM, et al. 1999. Bioavailability of heavy metals in terrestrial and aquatic systems: A quantitative approach. *Environ Toxicol Chem* 18(9):1882-1890.
- *Pocino M, Baute L, Malave I. 1991. Influence of the oral administration of excess copper on the immune response. *Fundam Appl Toxicol* 16(2):249-256.
- *Pocino M, Malave I, Baute L. 1990. Zinc administration restores the impaired immune response observed in mice receiving excess copper by oral route. *Immunopharmacol Immunotoxicol* 12(4):697-713.
- Poulsen OM, Christensen JM, Sabbioni E, et al. 1994. Trace element reference values in tissues from inhabitants of the European community. 5. Review of trace elements in blood, serum and urine and critical evaluation of reference values for the Danish population. *Sci Total Environ* 141:197-215.
- *Poulton DJ, Simpson KJ, Barton DR, et al. 1988. Trace metals and benthic invertebrates in sediments of nearshore Lake Ontario at Hamilton Harbour. *J Great Lakes Res* 14(1):52-65.
- *Prasad AS, Brewer GJ, Schoomaker EB, et al. 1978. Hypocupremia induced by zinc therapy in adults. *JAMA* 240:2166-2168.
- Prasad R, Kaur G, Nath R, et al. 1996. Molecular basis of pathophysiology of Indian childhood cirrhosis: Role of nuclear copper accumulation in liver. *Mol Cell Biochem* 156(1):25-30.

9. REFERENCES

- *Pratt WB, Omdahl JL, Sorenson JRJ. 1985. Lack of effects of copper gluconate supplementation. *Am J Clin Nutr* 42:681-682.
- Price LA, Walker NI, Clague AE, et al. 1996. Chronic copper toxicosis presenting as liver failure in an Australian child. *Pathology* 28:316-320.
- Prieditis H, Adamson IYR. 2002. Comparative pulmonary toxicity of various soluble metals found in urban particulate dusts. *Exp Lung Res* 28(7):563-576.
- Puig S, Thiele DJ. 2002. Molecular mechanisms of copper uptake and distribution. *Curr Opin Chem Biol* 6(2):171-180.
- *Que Hee SS, Finelli VN, Fricke FL, et al. 1982. Metal content of stack emissions, coal and fly ash from some eastern and western power plants in the U.S.A. as obtained by ICP-AES. *Int J Environ Anal Chem* 13:1-18.
- *Raghunath R, Tripathi RM, Khandekar RN, et al. 1997. Retention times of Pb, Cd, Cu and Zn in children's blood. *Sci Total Environ* 207(2-3):133-139.
- Raie RM. 1996. Regional variation in As, Cu, Hg, and Se and interaction between them. *Ecotoxicol Environ Saf* 35:248-252.
- *Rana SVS, Kumar A. 1980. Biological, haematological and histological observations in copper poisoned rats. *Ind Health* 18:9-17.
- *Raspor B, Nurnberg HW, Valenta P, et al. 1984. Studies in seawater and lake water on interactions of trace metals with humic substances isolated from marine and estuarine sediments. *Mar Chem* 15:217-230.
- *Raven KP, Loeppert RH. 1997. Heavy metals in the environment: Trace element composition of fertilizers and soil amendments. *J Environ Qual* 26:551-557.
- *Reed JS, Henningson JC. 1984. Acid precipitation and drinking water supplies. *J Am Water Works Assoc* 76:60-65.
- Reimann C, De Caritat P, Halleraker JH, et al. 1997. Rainwater composition in eight arctic catchments in Northern Europe (Finland, Norway and Russia). *Atmos Environ* 31:159-170.
- *Renwick WH, Edenborn HM. 1983. Metal and bacterial contamination in New Jersey estuarine sediments. *Environ Pollut Ser B* 5:175-185.
- *Rhee HM, Dunlap M. 1990. Acute cardiovascular toxic effects of copper in anesthetized rabbits. *Neurotoxicology* 11:355-360.
- *Rice KC. 1999. Trace-element concentrations in steambed sediment across the conterminous United States. *Environ Sci Technol* 33:2499-2504.
- Rice TM, Clarke RW, Godleski JL, et al. 2001. Differential ability of transition metals to induce pulmonary inflammation. *Toxicol Appl Pharmacol* 177:46-53.

9. REFERENCES

- *Richards BK, Steenhuis TS, Peverly JH. 1998. Metal mobility at an old, heavily loaded sludge application site. *Environ Pollut* 99(3):365-377.
- *Richards MP. 1999. Zinc, copper, and iron metabolism during porcine fetal development. *Biol Trace Elem Res* 69:27-44.
- *Rieuwerts JS, Thornton I, Farago ME, et al. 1998. Factors influencing metal bioavailability in soils: Preliminary investigations for the development of a critical loads approach for metals. *Chem Speciat Bioavail* 10(2):61-75.
- Riley RG, Zachara JM, Wobber FJ. 1992. Chemical contaminants on DOE lands and selection of contaminant mixtures for subsurface science research. Department of Energy. Washington, DC. DE 92 014 826.
- Riondato J, Vanhaecke F, Moens L, et al. 2000. Fast and reliable determination of (ultra-) trace and/or spectrally interfered elements in water by sector field ICP-MS. *J Anal Atom Spectrom* 15(4):341-345.
- *Ritter WF, Eastburn RP. 1978. Leaching of heavy metals from sewage sludge through coastal plain soils. *Commun Soil Sci Plant Anal* 9(9):785-798.
- *Rodeck B, Kardorff R, Melter M. 1999. Treatment of copper associated liver disease in childhood. *Eur J Med Res* 4(6):253-256.
- *Romo-Kroger CM, Morales JR, Dinator MI, et al. 1994. Heavy metals in the atmosphere coming from a copper smelter in Chile. *Atmos Environ* 28(4):705-711.
- *Rope SK, Arthur WJ, Craig TH, et al. 1988. Nutrient and trace elements in soil and desert vegetation of Southern Idaho. *Environ Monit Assess* 10:1-24.
- *Rösner U. 1998. Effects of historical mining activities on surface water and groundwater-an example from northwest Arizona. *Environ Geol* 33(4):224-230.
- *Roy WR. 1994. Groundwater contamination from municipal landfills in the USA. In: Adriano DC, ed. *Contamination of groundwaters: Case studies*. Northwood, UK. Scientific Review, 411-446.
- *Rubin ES. 1999. Toxic releases from power plants. *Environ Sci Technol* 33:3062-3067.
- Rule JH, Alden RW. 1996a. Interactions of Cd and Cu in aerobic estuarine sediments. I. Partitioning in geochemical fractions of sediments. *Environ Toxicol Chem* 15(4):460-465.
- Rule JH, Alden RW. 1996b. Interactions of Cd and Cu in anaerobic estuarine sediments. II. Bioavailability, body burdens and respiration effects as related to geochemical partitioning. *Environ Toxicol Chem* 15(4):466-471.
- *Rutzke MA, Gutenmann WH, Lisk DJ, et al. 2000. Toxic and nutrient element concentrations in soft tissues of zebra and quagga mussels from Lakes Erie and Ontario. *Chemosphere* 40:1353-1356.
- *Saenko, EL, Yaropolov AI, Harris ED. 1994. The biological functions of ceruloplasmin expressed through copper-binding sites and a cellular receptor. *J Trace Elem Exp Med* 7:69-88.

9. REFERENCES

- Saleh MA, Wilson BL. 1999. Analysis of metal pollutants in the Houston Ship Channel by inductively coupled plasma/mass spectrometry. *Ecotoxicol Environ Saf* 44:113-117.
- *Saltzer EI, Wilson JW. 1968. Allergic contact dermatitis due to copper. *Arch Dermatol* 98:375-376.
- *Saltzman BE, Gross SB, Yeager DW, et al. 1990. Total body burdens and tissue concentrations of lead, cadmium, copper, zinc, and ash in 55 human cadavers. *Environ Res* 52:126-145.
- Sanstead HH. 1982. Copper bioavailability and requirements. *Am J Clin Nutr* 35:809-814.
- Santon A, Irato P, Medici V, et al. 2003. Effect and possible role of Zn treatment in LEC rats, an animal model of Wilson's disease. *Biochim Biophys Acta* 1637:91-97.
- *Santschi PH, Nixon S, Pilson M, et al. 1984. Accumulation of sediments, trace metals (Pb, Cu) and total hydrocarbons in Narragansett Bay, Rhode Island. *Estuarine Coastal Shelf Sci* 19:427-449.
- *Sanudo-Wilhemly SA, Gill GA. 1999. Impact of the clean water act on the levels of toxic metals in urban estuaries: The Hudson River estuary revisited. *Environ Sci Technol* 33(20):1999.
- Sarkar B, Kruck TPA. 1967. Separation of Cu (II) - amino acid complexes and evidence for the existence of histidine-Cu (II)-glutamine and histidine-Cu (II)- serine complexes at physiological pH. *Can J Biochem* 45:2046-2049.
- Sarkar B, Wigfield Y. 1968. Evidence for albumin- Cu(II)- amino acid ternary complex. *Can J Biochem* 46:601-607.
- Sarkar B, Lingertat-Walsh K, Clarke JT. 1993. Copper-histidine therapy for Menkes disease. *J Pediatr* 123:828-830.
- Sato M, Hachiya N, Yamaguchi Y, et al. 1993. Deficiency of HOLO-, but not APO-, ceruloplasmins in genetically copper-intoxicated LEC mutant rat. *Life Sci* 53(18):1411-1416.
- *Sawaki M, Enomoto K, Hattori A, et al. 1994. Role of copper accumulation and metallothionein induction in spontaneous liver cancer development in LEC rats. *Carcinogenesis* 15(9):1833-1837.
- *Scharenberg W, Ebeling E. 1996. Distribution of heavy metals in a woodland food web. *Bull Environ Contam Toxicol* 56:389-396.
- *Scheinberg IH. 1979. Human health effects of copper. In: Nriagu JO, ed. *Copper in the environment. Part II. Health effects.* New York, NY: John Wiley & Sons, 82-101.
- *Scheinberg IH, Sternlieb I. 1994. Is non-Indian childhood cirrhosis caused by excess dietary copper? *Lancet* 344:1515-1516.
- Scheinberg IH, Sternlieb I. 1996. Wilson disease and idiopathic copper toxicosis. *Am J Clin Nutr* 63(5):842S-845S.
- *Schilsky ML. 1996. Wilson disease genetic basis of copper toxicity and natural history. *Sem Liver Dis* 16:83-95.

9. REFERENCES

- Schilsky ML, Blank RR, Czaja MJ, et al. 1989. Hepatocellular copper toxicity and its attenuation by zinc. *J Clin Invest* 84(5):1562-1568.
- Schmitt RC, Darwish HM, Cheney JC, et al. 1983. Copper transport kinetics by isolated rat hepatocytes. *Am J Physiol* 244:G183-G191.
- *Schnoor JL, Sato C, McKechnie, et al. 1987. Processes, coefficients, and models for simulating toxic organics and heavy metals in surface waters. EPA/600/3-87/015. Athens, GA: U.S. Environmental Protection Agency.
- *Schock MR, Neff CH. 1988. Trace metal contamination from brass fittings. *J Am Water Works Assoc* 7:47-56.
- *Schroeder HA, Nason AP, Tipton IH, et al. 1966. Essential trace metals in man: Copper. *J Chronic Dis* 19:1007-1034.
- *Schroeder WH, Dobson M, Kane DM, et al. 1987. Toxic trace elements associated with airborne particulate matter: A review. *J Air Pollut Control Assoc* 37(11):1267-1285.
- Schumann K, Classen HG, Dieter HH, et al. 2002. Hohenheim Consensus Workshop: Copper. *Eur J Clin Nutr* 56(6):469-483.
- Schwartz J, Weiss ST. 1990. Dietary factors and their relation to respiratory symptoms. *Am J Epidemiol* 132(1):67-76.
- *Scudlark JR, Conko KM, Church TM. 1994. Atmospheric wet deposition of trace elements to Chesapeake Bay: CBAD study year 1 results. *Atmos Environ* 28(8):1487-1498.
- *Sedlak DL, Phinney JT, Bdesworth WW. 1997. Strongly complexed Cu and Ni in wastewater effluents and surface runoff. *Environ Sci Technol* 31:3010-3016.
- *Semple AB, Parry WH, Phillips DE. 1960. Acute copper poisoning: An outbreak traced to contaminated water from a corroded geyser. *Lancet* 2:700-701.
- *Setchell BP, Waites GMH. 1975. The blood-testis barrier. In: Creep RO, Astwood EB, Geiger SR, eds. Philadelphia, PA: WB Saunders, 222-238.
- *Sharda B, Bhandari B. 1984. Copper concentration in plasma, cells, liver, urine, hair and nails in hepatobiliary disorders in children. *Indian Pediatr* 21:167-171.
- Sharma A, Talukder G. 1987. Effects of metals on chromosomes of higher organisms. *Environ Mutagen* 9:191-226.
- *Sharma VK, Millero FJ. 1988. Oxidation of copper (I) in seawater. *Environ Sci Technol* 22(7):768-771.
- Sharunda D, Diseker RA, Sinks T, et al. 1999. Copper in drinking water, Nebraska, 1994. *Int J Occup Environ Health* 5:256-261.
- Shibuya S, Takase Y, Sharma N. 1992. Esophageal ulcer due to ingestion of melted copper. *Dig Dis Sci* 37(11):1785-1790.

9. REFERENCES

- Shiraishi N, Taguchi T, Kinebuchi H. 1991. Copper-induced toxicity in *Macular* mutant mouse: An animal model for menkes' Kinky-Hair disease. *Toxicol Appl Pharmacol* 10(1):89-96.
- Shiraishi N, Taguchi T, Kinebuchi H. 1993. Effect of age and sex on copper-induced toxicity in the macular mutant mouse. An animal model fro Menkes' kinky-hair disease. *Biol Trace Elem Res* 39:129-137.
- *Sideris EG, Charalambous SC, Tsolomyty A, et al. 1988. Mutagenesis, carcinogenesis and the metal elements - DNA interaction. *Prog Clin Biol Res* 259:13-25.
- *Sina JF, Bean CL, Dysart GR, et al. 1983. Evaluation of the alkaline elution/rat hepatocyte assay as a predictor of carcinogenic/mutagenic potential. *Mutat Res* 113:357-391.
- *Singh I. 1983. Induction of reverse mutation and mitotic gene conversion by some metal compounds in *Saccharomyces cerevisiae*. *Mutat Res* 117:149-152.
- Singh SP, Ma LQ, Tack FMG, et al. 2000. Trace metal leachability of land-disposed dredged sediments. *J Environ Qual* 29:1124-1132.
- *Sirover MA, Loeb LA. 1976. Infidelity of DNA synthesis in vitro: Screening for potential metal mutagens of carcinogens. *Science* 94:1434-1436.
- *Smith CH, Bidlack WR. 1980. Interrelationship of dietary ascorbic acid and iron on the tissue distribution of ascorbic acid, iron and copper in female guinea pigs. *J Nutr* 110:1398-1408.
- *Sokol RJ, Devereaux M, Mierau GW, et al. 1990. Oxidant injury to hepatic mitochondrial lipids in rats with dietary copper overload. *Gastroenterology* 99(4):1061-1071.
- *Sokol RJ, Devereaux MW, O'Brien K, et al. 1993. Abnormal hepatic mitochondrial respiration and cytochrome C oxidase activity in rats with long-term copper overload. *Gastroenterology* 105(1):178-187.
- Sokol RJ, Devereaux MW, Traber MG, et al. 1989. Copper toxicity and lipid peroxidation in isolated rat hepatocytes: effect of Vitamin E. *Pediatr Res* 25(1):55-62.
- *Sora S, Carbone MLA, Pacciarini M, et al. 1986. Disomic and diploid meiotic products induced in *Saccharomyces cerevisiae* by the salts of 27 elements. *Mutagenesis* 1(1):21-28.
- Speisky H, Navarro P, Cherian MG, et al. 2003. Copper-binding proteins in human erythrocytes: Searching for potential biomarkers of copper over-exposure. *Biometals* 16:113-123.
- *Spitalny KC, Brondum J, Vogt RL, et al. 1984. Drinking-water-induced copper intoxication in a Vermont family. *Pediatrics* 74(6):1103-1106.
- *Stark P. 1981. Vineyard sprayer's lung-a rare occupational disease. *J Can Assoc Radiol* 32:183-184.
- Steinebach OM, Wolterbeek HT. 1994. Role of cytosolic copper, metallothionein and glutathione in copper toxicity in rat hepatoma tissue culture cells. *Toxicology* 9:75-90.

9. REFERENCES

- Steinkuhler C, Sapora O, Carri MT, et al. 1991. Increase of Cu, Zn-superoxide dimutase activity during differentiation of human K562 cells involves activation by copper of a constantly expressed copper-deficient protein. *J Biol Chem* 266:24580-24587.
- Steinnes E. 1990. Lead, cadmium and other metals in Scandinavian surface waters, with emphasis on acidification and atmospheric deposition. *Environ Toxicol Chem* 9:825-831.
- Steinnes E. 1995. A critical evaluation of the use of naturally growing moss to monitor the deposition of atmospheric metals. *Sci Total Environ* 160/161:243-249.
- *Stephenson T, Lester JN. 1987. Heavy metal behaviour during the activated sludge process. I. Extent of soluble and insoluble metal removal. *Sci Total Environ* 63:199-214.
- *Sternlieb I, Scheinberg IH. 1977. Human copper metabolism. In: *Medical and biologic effects of environmental pollutants-Copper*. Washington, DC: National Academy of Sciences.
- Sternlier I, Quintana N, Volenberg I, et al. 1995. An array of mitochondrial alterations in the hepatocytes of Long-Evans Cinnamon rats. *Hepatology* 22:1782-1787.
- *Stewart JH, Lassiter JV. 2001. Copper. In: Bingham E, Cohn B, Powell CH, eds. *Patty's toxicology*. New York, NY: John Wiley & Sons, Inc., 598-599.
- *Stiff MJ. 1971. The chemical states of copper in polluted fresh water and a scheme of analysis to differentiate them. *Water Res* 5:585-599.
- Stilwell DE, Gorny KD. 1997. Contamination of soil with copper, chromium, and arsenic under decks built from pressure treated wood. *Bull Environ Contam Toxicol* 58:22-29.
- Stockert RJ, Grushoff PS, Morell AG, et al. 1986. Transport and intracellular distribution of copper in a human hepatoblastoma cell line, HepG2. *Hepatology* 6:60-64.
- Stokinger HE. 1981. Copper. In: Clayton GD, Clayton FE, eds. *Patty's industrial hygiene and toxicology*. Chapter 11. New York, NY: John Wiley & Sons.
- *Strain WH, Hershey CO, McInnes S, et al. 1984. Hazards to groundwater from acid rain. *Trace Subst Environ Health* 18:178-184.
- Strand S, Hofmann WJ, Grambihler A, et al. 1998. Hepatic failure and liver cell damage in acute Wilson's disease involve CD95 (APO-1/Fas) mediated apoptosis. *Nat Med* 4(5):588-593.
- Strickland GT, Leu ML. 1975. Wilson's disease: Clinical and laboratory manifestations in 40 patients. *Am J Med* 54(2):113-137.
- *Strickland GT, Beckner WM, Leu ML. 1972. Absorption of copper in homozygotes and heterozygotes for Wilson's disease and controls: Isotope tracer studies with ⁶⁷Cu and ⁶⁴Cu. *Clin Sci* 43:617-625.
- *Suciu I, Prodan L, Lazar V, et al. 1981. Research on copper poisoning. *Med Lav* 3:190-197.
- *Sugawara N, Li D, Katakura M. 1994. Biliary excretion of copper in Fischer rats treated with copper salt and in Long-Evans Cinnamon (LEC) rats with an inherently abnormal copper metabolism. *Biol Trace Elem Res* 46:125-134.

9. REFERENCES

- *Sugawara N, Li D, Sugawara C, et al. 1995. Response of hepatic function to hepatic copper deposition in rats fed a diet containing copper. *Biol Trace Elem Res* 49:161-169.
- *Sugawara N, Sugawara C, Katakura M, et al. 1991. Harmful effect of administration of copper on LEC rats. *Res Commun Chem Pathol Pharmacol* 73(3):289-297.
- *Sugawara N, Sugawara C, Li D, et al. 1992. Copper metabolism in new mutant Long-Evans Cinnamon (LEC) rats causing hereditary hepatitis: Gastrointestinal absorption and distribution of radiostopic copper (^{64}Cu). *Res Commun Chem Pathol Pharmacol* 76(2):233-243.
- Sullivan JM, Janovitz EB, Robinson FR. 1991. Copper toxicosis in veal calves. *J Vet Diagn Invest* 3(2):161-164.
- Sunderic D, Dozic S, Hosovski E. 1990. Histologic changes in peripheral nerves in rats with chronic copper poisoning. *Srp Arh Celok Lek* 118(7-8):261-264. (Serbo-Croatian, Cyrillic)
- Sunderic D, Hosovsk E, Dasic S. 1990. Histologic changes in the rat brain during chronic copper poisoning. *Srp Arh Celok Lek* 118(5-6):171-174. (Cyrillic)
- *Suttle NF, Mills CF. 1966a. Studies of the toxicity of copper to pigs. 1. Effects of oral supplements of zinc and iron salts on the development of copper toxicosis. *Br J Nutr* 20:135-148.
- *Suttle NF, Mills CF. 1966b. Studies of the toxicity of copper to pigs. 2. Effect of protein source and other dietary components on the response to high and moderate intakes of copper. *Br J Nutr* 20:149-161.
- *Suzuki KT, Kanno S, Misawa S, et al. 1995. Copper metabolism leading to and following acute hepatitis in LEC rats. *Toxicology* 97(1-3):81-92.
- Suzuki KT, Karasawa A, Sunaga H, et al. 1989. Uptake of copper from the bloodstream and its relation to induction of metallothionein synthesis in the rat. *Comp Biochem Physiol C* 94(1):93-97.
- *Sweet CW, Vermette SJ, Landsberger S. 1993. Sources of toxic trace elements in urban air in Illinois. *Environ Sci Technol* 27(12):2502-2510.
- *Sylva RN. 1976. The environmental chemistry of copper (II) in aquatic systems. *Water Res* 10:789-792.
- *Tan WT, Tan GS, Khan ISAN. 1988. Solubilities of trace copper and lead species and the complexing capacity of river water in the Linggi River Basin. *Environ Pollut* 52:221-235.
- *Tanner MS. 1998. Role of copper in Indian childhood cirrhosis. *Am J Clin Nutr* 67:1074S-1081S.
- Tao TY, Liu F, Klomp L, et al. 2003. The copper toxicosis gene product *murr1* directly interacts with the Wilson disease protein. *J Biol Chem* 278(43):41593-41596.
- *Taylor GJ, Crowder AA. 1983. Accumulation of atmospherically deposited metals in wetland soils of Sudbury, Ontario. *Water Air Soil Pollut* 19:29-42.
- Tennant J, Stansfield M, Yamaji S, et al. 2002. Effects of copper on the expression of metal transporters in human intestinal Caco-2 cells. *FEBS Lett* 527(1-3):239-244.

9. REFERENCES

Theis TL, Young TC, Huang M, et al. 1994. Leachate characteristics and composition of cyanide-bearing wastes from manufactured gas plants. *Environ Sci Technol* 28:99-106.

*Tinwell H, Ashby J. 1990. Inactivity of copper sulphate in a mouse bone-marrow micronucleus assay. *Mutat Res* 245(3):233-236.

Tkeshelashvili LK, McBride T, Spence K, et al. 1992. Mutation spectrum of copper-induced DNA damage. Additions and corrections. *J Biol Chem* 267(19):13778.

Tollestrup K, Frost FJ, Harter L, et al. 2002. Mortality in children residing near the Asarco Copper Smelter in Ruston, Washington [Abstract]. *Am J Epidemiol* 155(11):S39.

*Town RM, Filella M. 2000. A comprehensive systematic compilation of complexation parameters reported for trace metals in natural waters. *Aquat Sci* 62:252-295.

Toyokuni S, Sagripanti JL. 1994. Increased 8-hydroxydeoxyguanosine in kidney and liver of rats continuously exposed to copper. *Toxicol Appl Pharmacol* 126(1):91-97.

*Toyokuni S, Tanaka T, Nishiyama Y, et al. 1996. Induction of renal cell carcinoma in male wistar rats treated with cupric nitrilotriacetate. *Lab Invest* 75:239-248.

*TRI01. 2003. TRI explorer: Providing access to EPA's toxics release inventory data. U.S. Environmental Protection Agency. Toxic Release Inventory. <http://www.epa.gov/triexplorer>. August 24, 2003.

Trollmann R, Neureiter D, Lang T, et al. 1999. Late manifestation of Indian childhood cirrhosis in a 3-year-old German girl. *Eur J Pediatr* 158(5):375-378.

*Truitt RE, Weber JH. 1981. Copper(II)- and cadmium(II)-binding abilities of some New Hampshire freshwaters determined by dialysis titration. *Environ Sci Technol* 15:1204-1208.

Tshiashala MD, Kabengele K, Lumu BM. 1990. Trace element determination in scalp hair of people working at a copper smelter. *Biol Trace Elem Res* 26-27:287-294.

*Tso W-W, Fung W-P. 1981. Mutagenicity of metallic cations. *Toxicol Lett* 8:195-200.

Tsuda T, Inoue T, Kojima M, et al. 1995. Market basket and duplicate portion estimation of dietary intakes of cadmium, mercury, arsenic, copper, manganese, and zinc by Japanese adults. *J AOAC Int* 78(6):1363-1367.

*Tuddenham WM, Dougall PA. 1978. Copper. In: Kirk Othmer's encyclopedia of chemical technology. Vol. 6, 3rd ed. New York, NY: John Wiley & Sons, 819-869.

*Turnlund JR. 1989. Stable isotope studies of the effect of dietary copper on copper absorption and excretion. *Adv Exp Med Biol* 258:21-28.

Turnlund JR. 1998. Human whole-body copper metabolism. *Am J Clin Nutr* 67:960S-964S.

*Turnlund JR, Keyes WR, Anderson HL, et al. 1989. Copper absorption and retention in young men at three levels of dietary copper by use of the stable isotope ⁶⁵Cu. *Am J Clin Nutr* 49:870-878.

9. REFERENCES

- *Turnlund JR, King JC, Gong B, et al. 1985. A stable isotope study of copper absorption in young men: Effect of phytate and α -cellulose. *Am J Clin Nutr* 42:18-23.
- *Turnlund JR, Michel MC, Keyes WR, et al. 1982. Copper absorption in elderly men determined by using stable ^{65}Cu . *Am J Clin Nutr* 36:587-591.
- *Turnlund JR, Reager RD, Costa F. 1988a. Iron and copper absorption in young and elderly men. *Nutr Res* 8:333-343.
- *Turnlund JR, Swanson CA, King JC. 1983. Copper absorption and retention in pregnant women fed diets based on animal and plant proteins. *J Nutr* 113:2346-2352.
- *Turnlund JR, Wada L, King JC, et al. 1988b. Copper absorption in young men fed adequate and low zinc diets. *Biol Trace Elem Res* 17:31-41.
- *Tyler LD, McBride MB. 1982. Mobility and extractability of cadmium, copper, nickel, and zinc in organic and mineral soil columns. *Soil Sci* 134(3):198-205.
- *Underwood EJ. 1977. Trace elements in human and animal nutrition. 4th ed. New York, NY: Academic Press, 43-87.
- Unlu K. 1998. Transport of metals leaching from land-disposed oil field wastes. *Waste Manage Res* 16(6):541-554.
- *USGS. 1986. Extractable cadmium, mercury, copper, lead, and zinc in the lower Columbia River Estuary, Oregon and Washington. U.S. Geological Survey Water Resources Investigations Report 86(4088):Portland, Oregon: U.S. Department of Interior.
- *USGS. 1994. Copper minerals yearbook. United States Geological Survey. <http://minerals.usgs.gov/minerals/pubs/commodity/copper>.
- *USGS. 2000. Copper minerals yearbook. United States Geological Survey. <http://www.usgs.gov/minerals/pubs/commodity/copper>.
- *USGS. 2001. Copper. U.S. Geological Survey minerals yearbook. United States Geological Survey. <http://www.usgs.gov/minerals/pubs/commodity/copper/coppmyb01>.
- *USGS. 2002. Mineral commodity summary. United States Geological Survey. <http://www.usgs.gov/minerals/pubs/commodity/copper>.
- *Van Campen DR, Mitchell EA. 1965. Absorption of Cu^{64} , Zn^{65} , Mo^{99} , and Fe^{59} from ligated segments of the rat gastrointestinal tract. *J Nutr* 86:120-124.
- Van Campen DR, Scaife PU. 1967. Zinc interference with copper absorption in rats. *Nutrition* 91:473-476.
- Van den Berg GJ, van den Hamer CJA. 1984. Trace metal uptake in liver cells. 1. Influence of albumin in the medium on the uptake of copper by hepatoma cells. *J Inorg Biochem* 22:73-84.

9. REFERENCES

- Van den Broeck K, Helsen L, Vandecasteele C, et al. 1997. Determination and characterization of copper, chromium and arsenic in chromated copper arsenate (CCA) treated wood and its pyrolysis residues by inductively coupled plasma mass spectrometry. *Analyst* 122:695-700.
- van de Sluis B, Rothuizen J, Pearson PL, et al. 2002. Identification of a new copper metabolism gene by positional cloning in a purebred dog population. *Hum Mol Gen* 11(2):165-173.
- *van Ryssen JB. 1994. The effectiveness of using supplementary zinc and molybdenum to reduce the copper content in the liver of hypercuprotic sheep. *J S Afr Vet Assoc* 65(2):59-63.
- Van Veen E, Burton N, Comber S, et al. 2002. Speciation of copper in sewage effluents and its toxicity to *Daphnia Magna*. *Environ Toxicol Chem* 21(2):275-280.
- *Varada KR, Harper RG, Wapnir RA. 1993. Development of copper intestinal absorption in the rat. *Biochem Med Metab Biol* 50(3):277-283.
- *Vaughan LA, Weber CW, Kemberling SR. 1979. Longitudinal changes in the mineral content of human milk. *Am J Clin Nutr* 32:2301-2306.
- Verloo M, Eeckhout M. 1990. Metal species transformations in soils: An analytical approach. *Int J Environ Anal Chem* 39:179-186.
- Videla LA, Fernández V, Tapia G, et al. 2003. Oxidative stress-mediated hepatotoxicity of iron and copper: Role of Kupffer cells. *Biometals* 16:103-111.
- *Vieira I, Sonnier M, Cresteil T. 1996. Developmental expression of CYP2E1 in the human liver: Hypermethylation control of gene expression during the neonatal period. *Eur J Biochem* 238:476-483.
- *Villar TG. 1974. Vineyard Sprayer's Lung: Clinical aspects. *Am Rev Respir Dis* 110:545-555.
- *Villar TG, Nogueira T. 1980. Radiology and respiratory function in Vineyard Sprayer's Lung. 30:61-70.
- Viklander M. 1999. Dissolved and particle-bound substances in urban snow. *Water Sci Technol* 39(12):37-32.
- Vinlove MP, Britt J, Cornelius J. 1992. Copper toxicity in a rabbit. *Lab Anim Sci* 42(6):614-615.
- Vodichenska T. 1988. [The biological effect of copper taken up by the body via drinking water.] *Probl Khig* 13:29-36. (Bulgarian)
- Vodichenskaia TS, Dinoeva SK. 1989. [Experimental study of the atherogenic effect of copper after its intake with drinking water.] *Gig Sanit* 2:64-66. (Russian)
- *Vong RJ, Baker BM, Brechtel FJ, et al. 1997. Ionic and trace element composition of cloud water collected on the Olympic peninsula of Washington state. *Atmos Environ* 31(13):1991-2001.
- Von Gunten HR, Sturm M, Moser RN. 1997. 200-Year record of metals in lake sediments and natural background concentrations. *Environ Sci Technol* 31:2193-2197.

9. REFERENCES

- von Muhlen Dahl KE. 1996. Copper tubings, home wells and early childhood cirrhosis. *Eur J Pediatr* 155(12):1061-1062.
- Vrzgulova M. 1993. Histological and submicroscopical findings on the seminiferous parenchyma in rams after copper oxide intoxication from industrial emissions. *Funct Dev Morphol* 3(2):115-119.
- Vulpe CD, Packman S. 1995. Cellular copper transport. *Annu Rev Nutr* 15:293-322.
- Wada H, Zhou XJ, Ishizuki T, et al. 1992. Direct determination of copper in serum by flow-injection analysis. *Anal Chim Acta* 261(1-2):87-95.
- *Wake SA, Mercer JFB. 1985. Induction of metallothionein mRNA in rat liver and kidney after copper chloride injection. *Biochem J* 228:425-432.
- *Walker WR, Reeves RR, Brosnan M, et al. 1977. Perfusion of intact skin by a saline solution of bis(glycinato) copper(II). *Bioinorg Chem* 7:271-276.
- *Walsh FM, Crosson FJ, Bayley J, et al. 1977. Acute copper intoxication. *Am J Dis Child* 131:149-151.
- *Walshe JM. 1996. Treatment of Wilson's disease: the historical background. *Q J Med* 89:252-263.
- *Walshe JM, Yealland M. 1993. Chelation treatment of neurological Wilson's disease. *Q J Med* 86:197-204.
- Wang J, Tian B. 1992. Screen-printed stripping voltammetric/potentiometric electrodes for decentralized testing of trace lead. *Anal Chem* 64(15):1706-1709.
- Wang J, Larson D, Foster N, et al. 1995. Remote electrochemical sensor for trace metal contaminants. *Anal Chem* 67(8):1481-1485.
- Wapnir RA. 1998. Copper absorption and bioavailability. *Am J Clin Nutr* 67:1054S-1060S.
- *Wapnir RA, Devas G, Solans CV. 1993. Inhibition of intestinal copper absorption by divalent cations and low-molecular weight ligands in the rat. *Biol Trace Elem Res* 36:291-305.
- Wataha JC, Lockwood PE, Schedle A, et al. 2002. Ag, Cu, Hg and Ni ions alter the metabolism of human monocytes during extended low-dose exposures. *J Oral Rehabil* 29:133-139.
- *Weant GE. 1985. Sources of copper air emissions. Research Triangle Park, NC. Air and Energy Engineering Research Laboratory, U.S. Environmental Protection Agency. EPA 600/2/-85-046.
- *Weast RC. 1980. CRC handbook of chemistry and physics. 61st edition. Boca Raton, FL: CRC Press, B13, B97-B100.
- *Weber PM, O'Reilly S, Pollycove M, et al. 1969. Gastrointestinal absorption of copper: Studies with ⁶⁴Cu, ⁹⁵Zr, a whole-body counter and the scintillation camera. *J Nucl Med* 10(9):591-596.
- Weiner AL, Cousins RJ. 1980. Copper accumulation and metabolism in primary monolayer cultures of rat liver parenchymal cells. *Biochim Biophys Acta* 629:113-125.

9. REFERENCES

- Weisberg SB, Wilson HT, Heimbuch DG, et al. 2000. Comparison of sediment metal: Aluminum relationships between the eastern and Gulf coasts of the United States. *Environ Monit Assess* 61:373-385.
- *Weiss KC, Linder MC. 1985. Copper transport in rats involving a new plasma protein. *Am J Physiol Endocrinol Metab* 249(12):E77-E88.
- *West JR, Smith HW, Chasis H. 1948. Glomerular filtration rate, effective renal blood flow, and maximal tubular excretory capacity in infancy. *J Pediatr* 32:10-18.
- *Whanger PD, Weswig PH. 1971. Effect of supplementary zinc on the intracellular distribution of hepatic copper in rats. *J Nutr* 101:1093-1098.
- White AR, Cappai R. 2003. Neurotoxicity from glutathione depletion is dependent on extracellular trace copper. *J Neurosci Res* 71(6):889-897.
- *WHO. 1996. Trace elements in human nutrition and health. Copper. Geneva, Switzerland: World Health Organization, 123-143.
- *WHO. 1998. Copper. Geneva: International Programme on Chemical Safety, World Health Organization. Environmental Health Criteria 200.
<http://www.inchem.org/documents/ehc/ehc/ehc200.html>. July 29, 2004.
- *Widdowson EM, Dickerson JWT. 1964. Chemical composition of the body. In: Comar CL, Bronner F, eds. *Mineral metabolism: An advanced treatise. Volume II: The elements Part A*. New York: Academic Press.
- *Widdowson EM, Dauncey J, Shaw JCL. 1974. Trace elements in fetal and early postnatal development. *Proc Nutr Soc* 33:275-284.
- *Wijmenga C. 2002. Non-Indian childhood cirrhosis. Using a founder population to identify the underlying genetic defect. In: Massaro EJ, ed. *Handbook of copper pharmacology and toxicology*. Totowa, NJ: Humana Press, 369-382.
- *Wilhelm M, Hafner D, Lombeck I, et al. 1991. Monitoring of cadmium, copper, lead and zinc status in young children using toenails: Comparison with scalp hair. *Sci Total Environ* 103:199-207.
- Williams DE, Vlamis J, Pukite AH, et al. 1984. Metal movement in sludge-treated soils after six years of sludge addition: 1. Cadmium, copper, lead, and zinc. *Soil Sci* 137(5):351-359.
- *Williams DM. 1982. Clinical significance of copper deficiency and toxicity in the world population. *Clinical, biochemical, and nutritional aspects of trace elements*. Chapter 15. New York, NY: Alan R. Liss, Inc.
- Wilson JC. 1989. A prospective New Zealand study of fertility after removal of copper intrauterine contraceptive devices for conception and because of complications: A four-year study. *Am J Obstet Gynecol* 160(2):391-396.
- *Windholz M. 1983. *The Merck Index*. 10th ed. Rahway, NJ: Merck & Co., 358-359; 2484-2485.

9. REFERENCES

- *Wise SA, Zeisler R. 1984. The pilot environmental specimen bank program. *Environ Sci Technol* 18(10):302A-307A.
- *Wong PK. 1988. Mutagenicity of heavy metals. *Bull Environ Contam Toxicol* 40:597-603.
- Wu J, Boyle EA. 1997. Low bank preconcentration technique for the determination of lead, copper, and cadmium in small-volume seawater samples by isotope dilution ICPMS. *Anal Chem* 69:2464-2470.
- *Wu J, Laird DA, Thompson ML. 1999. Sorption and desorption of copper on soil clay components. *J Environ Qual* 28:334-338.
- *Wyllie J. 1957. Copper poisoning at a cocktail party. *Am J Public Health* 47:617.
- Xue H, Sunda WG. 1997. Comparison of Cu^{2+} measurements in lake water determined by ligand exchange and cathodic stripping voltammetry and by ion-selective electrode. *Environ Sci Technol* 31:1902-1909.
- *Xue H, Lurdes MS, Reutlinger M, et al. 1991. Copper (I) in fogwater: Determination and interactions with sulfite. *Environ Sci Technol* 25:1716-1722.
- *Yadrick MK, Kenney MA, Winterfeldt EA. 1989. Iron, copper, and zinc status: Response to supplementation with zinc or zinc and iron in adult females. *Am J Clin Nutr* 49:145-150.
- Yamada T, Kim JK, Suzuki Y. 1993. Reduced efficiency of copper transport from cytosolic to monocytoplasmic fractions in LEC mutant rat. *Res Commun Chem Pathol Pharmacol* 81(2):243-246.
- Yamada T, Sogawa K, Kim J-K, et al. 1998. Increased polyploidy, delayed mitosis and reduced protein phosphatase -1 activity associated with excess copper in the Long Evans Cinnamon rat. *Res Commun Chem Pathol Pharmacol* 99(3):283-304.
- *Yamane Y, Sakai K, Umeda T, et al. 1984. Suppressive effect of cupric acetate on DNA alkylation, DNA synthesis and tumorigenesis in the liver of dimethylnitrosamine-treated rats. *Gann* 75(12):1062-1069.
- Yamauchi T, Yamamoto I. 1990. CuSO_4 as an inhibitor of B cell proliferation. *Jpn J Pharmacol* 54(4):455-460.
- *Yannoni CC, Piorkowski T. 1995. Profile of lead and copper levels in house plumbing and service pipe. *J New Engl Water Works Assoc* 109(3):192-210.
- *Yeats PA. 1988. The distribution of trace metals in ocean waters. *Sci Total Environ* 72:131-149.
- *Yelin G, Taff ML, Sadowski GE, et al. 1987. Copper toxicity following massive ingestion of coins. *Am J Forensic Med Pathol* 8(1):78-85.
- Yu S, Van Der Meer R, Beynen AC. 2002. Excessive hepatic copper accumulation in jaundiced rats fed a high-copper diet. *Biol Trace Elem Res* 88(3):255-269.
- *Yu S, Wests CE, Beynen AC. 1994. Increasing intakes of iron reduces status, absorption and biliary excretion of copper in rats. *Br J Nutr* 71:887-895.

9. REFERENCES

- Zabel M, Lindscheid KR, Mark H. 1990. [Copper sulfate allergy with special reference to internal exposure.] *Z Hautkr* 65(5):481-482; 485-486. (German)
- Zerounian NR, Redekosky C, Malpe R, et al. 2003. Regulation of copper absorption by copper availability in the Caco-2 cell intestinal model. *Am J Physiol Gastrointest Liver Physiol* 284(5):G739-G747.
- Zhang SZ, Noordin MM, Rahman S-O, et al. 2000. Effects of copper overload on hepatic lipid peroxidation and antioxidant defense in rats. *Vet Hum Toxicol* 42(5):261-264.
- Zhou W, Kornegay ET, Lindemann MD, et al. 1994. Stimulation of growth by intravenous injection of copper in weanling pigs. *J Anim Sci* 72:2395-2403.
- *Ziegler EE, Edwards BB, Jensen RL, et al. 1978. Absorption and retention of lead by infants. *Pediatr Res* 12:29-34.
- *Zietz BP, deVergara JD, Dunkelberg H. 2003b. Copper concentrations in tap water and possible effects on infant's health-results of a study in Lower Saxony, Germany. *Environ Res* 92(2):129-138.
- *Zietz BP, Dieter HH, Lakomek M, et al. 2003a. Epidemiological investigation on chronic copper toxicity to children exposed via the public drinking water supply. *Sci Total Environ* 301(1-3):127-144.
- Zietz BP, Kessler-Gaedtke B, Schneider H, et al. 2002. Results of an investigation on chronic copper toxicity to children exposed via the public drinking water supply [Abstract]. *Naunyn-Schmiedeberg's Arch Pharmacol* 365(Suppl 1):R135.
- Zipper J, Medel M, Prager R. 1969. Suppression of fertility by intrauterine copper and zinc in rabbits. *Am J Obstet Gynecol* 105(4):529-534.

10. GLOSSARY

Absorption—The taking up of liquids by solids, or of gases by solids or liquids.

Acute Exposure—Exposure to a chemical for a duration of 14 days or less, as specified in the Toxicological Profiles.

Adsorption—The adhesion in an extremely thin layer of molecules (as of gases, solutes, or liquids) to the surfaces of solid bodies or liquids with which they are in contact.

Adsorption Coefficient (K_{oc})—The ratio of the amount of a chemical adsorbed per unit weight of organic carbon in the soil or sediment to the concentration of the chemical in solution at equilibrium.

Adsorption Ratio (K_d)—The amount of a chemical adsorbed by sediment or soil (i.e., the solid phase) divided by the amount of chemical in the solution phase, which is in equilibrium with the solid phase, at a fixed solid/solution ratio. It is generally expressed in micrograms of chemical sorbed per gram of soil or sediment.

Benchmark Dose (BMD)—Usually defined as the lower confidence limit on the dose that produces a specified magnitude of changes in a specified adverse response. For example, a BMD10 would be the dose at the 95% lower confidence limit on a 10% response, and the benchmark response (BMR) would be 10%. The BMD is determined by modeling the dose response curve in the region of the dose response relationship where biologically observable data are feasible.

Benchmark Dose Model—A statistical dose-response model applied to either experimental toxicological or epidemiological data to calculate a BMD.

Bioconcentration Factor (BCF)—The quotient of the concentration of a chemical in aquatic organisms at a specific time or during a discrete time period of exposure divided by the concentration in the surrounding water at the same time or during the same period.

Biomarkers—Broadly defined as indicators signaling events in biologic systems or samples. They have been classified as markers of exposure, markers of effect, and markers of susceptibility.

Cancer Effect Level (CEL)—The lowest dose of chemical in a study, or group of studies, that produces significant increases in the incidence of cancer (or tumors) between the exposed population and its appropriate control.

Carcinogen—A chemical capable of inducing cancer.

Case-Control Study—A type of epidemiological study that examines the relationship between a particular outcome (disease or condition) and a variety of potential causative agents (such as toxic chemicals). In a case-controlled study, a group of people with a specified and well-defined outcome is identified and compared to a similar group of people without outcome.

Case Report—Describes a single individual with a particular disease or exposure. These may suggest some potential topics for scientific research, but are not actual research studies.

Case Series—Describes the experience of a small number of individuals with the same disease or exposure. These may suggest potential topics for scientific research, but are not actual research studies.

10. GLOSSARY

Ceiling Value—A concentration of a substance that should not be exceeded, even instantaneously.

Chronic Exposure—Exposure to a chemical for 365 days or more, as specified in the Toxicological Profiles.

Cohort Study—A type of epidemiological study of a specific group or groups of people who have had a common insult (e.g., exposure to an agent suspected of causing disease or a common disease) and are followed forward from exposure to outcome. At least one exposed group is compared to one unexposed group.

Cross-sectional Study—A type of epidemiological study of a group or groups of people that examines the relationship between exposure and outcome to a chemical or to chemicals at one point in time.

Data Needs—Substance-specific informational needs that if met would reduce the uncertainties of human health assessment.

Developmental Toxicity—The occurrence of adverse effects on the developing organism that may result from exposure to a chemical prior to conception (either parent), during prenatal development, or postnatally to the time of sexual maturation. Adverse developmental effects may be detected at any point in the life span of the organism.

Dose-Response Relationship—The quantitative relationship between the amount of exposure to a toxicant and the incidence of the adverse effects.

Embryotoxicity and Fetotoxicity—Any toxic effect on the conceptus as a result of prenatal exposure to a chemical; the distinguishing feature between the two terms is the stage of development during which the insult occurs. The terms, as used here, include malformations and variations, altered growth, and *in utero* death.

Environmental Protection Agency (EPA) Health Advisory—An estimate of acceptable drinking water levels for a chemical substance based on health effects information. A health advisory is not a legally enforceable federal standard, but serves as technical guidance to assist federal, state, and local officials.

Epidemiology—Refers to the investigation of factors that determine the frequency and distribution of disease or other health-related conditions within a defined human population during a specified period.

Genotoxicity—A specific adverse effect on the genome of living cells that, upon the duplication of affected cells, can be expressed as a mutagenic, clastogenic, or carcinogenic event because of specific alteration of the molecular structure of the genome.

Half-life—A measure of rate for the time required to eliminate one half of a quantity of a chemical from the body or environmental media.

Immediately Dangerous to Life or Health (IDLH)—The maximum environmental concentration of a contaminant from which one could escape within 30 minutes without any escape-impairing symptoms or irreversible health effects.

Incidence—The ratio of individuals in a population who develop a specified condition to the total number of individuals in that population who could have developed that condition in a specified time period.

10. GLOSSARY

Intermediate Exposure—Exposure to a chemical for a duration of 15–364 days, as specified in the Toxicological Profiles.

Immunologic Toxicity—The occurrence of adverse effects on the immune system that may result from exposure to environmental agents such as chemicals.

Immunological Effects—Functional changes in the immune response.

In Vitro—Isolated from the living organism and artificially maintained, as in a test tube.

In Vivo—Occurring within the living organism.

Lethal Concentration(Lo) (LC_{Lo})—The lowest concentration of a chemical in air that has been reported to have caused death in humans or animals.

Lethal Concentration(50) (LC₅₀)—A calculated concentration of a chemical in air to which exposure for a specific length of time is expected to cause death in 50% of a defined experimental animal population.

Lethal Dose(Lo) (LD_{Lo})—The lowest dose of a chemical introduced by a route other than inhalation that has been reported to have caused death in humans or animals.

Lethal Dose(50) (LD₅₀)—The dose of a chemical that has been calculated to cause death in 50% of a defined experimental animal population.

Lethal Time(50) (LT₅₀)—A calculated period of time within which a specific concentration of a chemical is expected to cause death in 50% of a defined experimental animal population.

Lowest-Observed-Adverse-Effect Level (LOAEL)—The lowest exposure level of chemical in a study, or group of studies, that produces statistically or biologically significant increases in frequency or severity of adverse effects between the exposed population and its appropriate control.

Lymphoreticular Effects—Represent morphological effects involving lymphatic tissues such as the lymph nodes, spleen, and thymus.

Malformations—Permanent structural changes that may adversely affect survival, development, or function.

Minimal Risk Level (MRL)—An estimate of daily human exposure to a hazardous substance that is likely to be without an appreciable risk of adverse noncancer health effects over a specified route and duration of exposure.

Modifying Factor (MF)—A value (greater than zero) that is applied to the derivation of a Minimal Risk Level (MRL) to reflect additional concerns about the database that are not covered by the uncertainty factors. The default value for a MF is 1.

Morbidity—State of being diseased; morbidity rate is the incidence or prevalence of disease in a specific population.

Mortality—Death; mortality rate is a measure of the number of deaths in a population during a specified interval of time.

10. GLOSSARY

Mutagen—A substance that causes mutations. A mutation is a change in the DNA sequence of a cell's DNA. Mutations can lead to birth defects, miscarriages, or cancer.

Necropsy—The gross examination of the organs and tissues of a dead body to determine the cause of death or pathological conditions.

Neurotoxicity—The occurrence of adverse effects on the nervous system following exposure to a chemical.

No-Observed-Adverse-Effect Level (NOAEL)—The dose of a chemical at which there were no statistically or biologically significant increases in frequency or severity of adverse effects seen between the exposed population and its appropriate control. Effects may be produced at this dose, but they are not considered to be adverse.

Octanol-Water Partition Coefficient (K_{ow})—The equilibrium ratio of the concentrations of a chemical in *n*-octanol and water, in dilute solution.

Odds Ratio (OR)—A means of measuring the association between an exposure (such as toxic substances and a disease or condition) which represents the best estimate of relative risk (risk as a ratio of the incidence among subjects exposed to a particular risk factor divided by the incidence among subjects who were not exposed to the risk factor). An odds ratio of greater than 1 is considered to indicate greater risk of disease in the exposed group compared to the unexposed group.

Organophosphate or Organophosphorus Compound—A phosphorus-containing organic compound and especially a pesticide that acts by inhibiting cholinesterase.

Permissible Exposure Limit (PEL)—An Occupational Safety and Health Administration (OSHA) allowable exposure level in workplace air averaged over an 8-hour shift of a 40-hour workweek.

Pesticide—General classification of chemicals specifically developed and produced for use in the control of agricultural and public health pests.

Pharmacokinetics—The dynamic behavior of a material in the body, used to predict the fate (disposition) of an exogenous substance in an organism. Utilizing computational techniques, it provides the means of studying the absorption, distribution, metabolism, and excretion of chemicals by the body.

Pharmacokinetic Model—A set of equations that can be used to describe the time course of a parent chemical or metabolite in an animal system. There are two types of pharmacokinetic models: data-based and physiologically-based. A data-based model divides the animal system into a series of compartments, which, in general, do not represent real, identifiable anatomic regions of the body, whereas the physiologically-based model compartments represent real anatomic regions of the body.

Physiologically Based Pharmacodynamic (PBPD) Model—A type of physiologically based dose-response model that quantitatively describes the relationship between target tissue dose and toxic end points. These models advance the importance of physiologically based models in that they clearly describe the biological effect (response) produced by the system following exposure to an exogenous substance.

Physiologically Based Pharmacokinetic (PBPK) Model—Comprised of a series of compartments representing organs or tissue groups with realistic weights and blood flows. These models require a

10. GLOSSARY

variety of physiological information: tissue volumes, blood flow rates to tissues, cardiac output, alveolar ventilation rates, and possibly membrane permeabilities. The models also utilize biochemical information such as air/blood partition coefficients, and metabolic parameters. PBPK models are also called biologically based tissue dosimetry models.

Prevalence—The number of cases of a disease or condition in a population at one point in time.

Prospective Study—A type of cohort study in which the pertinent observations are made on events occurring after the start of the study. A group is followed over time.

q1*—The upper-bound estimate of the low-dose slope of the dose-response curve as determined by the multistage procedure. The q1* can be used to calculate an estimate of carcinogenic potency, the incremental excess cancer risk per unit of exposure (usually $\mu\text{g/L}$ for water, mg/kg/day for food, and $\mu\text{g/m}^3$ for air).

Recommended Exposure Limit (REL)—A National Institute for Occupational Safety and Health (NIOSH) time-weighted average (TWA) concentrations for up to a 10-hour workday during a 40-hour workweek.

Reference Concentration (RfC)—An estimate (with uncertainty spanning perhaps an order of magnitude) of a continuous inhalation exposure to the human population (including sensitive subgroups) that is likely to be without an appreciable risk of deleterious noncancer health effects during a lifetime. The inhalation reference concentration is for continuous inhalation exposures and is appropriately expressed in units of mg/m^3 or ppm.

Reference Dose (RfD)—An estimate (with uncertainty spanning perhaps an order of magnitude) of the daily exposure of the human population to a potential hazard that is likely to be without risk of deleterious effects during a lifetime. The RfD is operationally derived from the no-observed-adverse-effect level (NOAEL—from animal and human studies) by a consistent application of uncertainty factors that reflect various types of data used to estimate RfDs and an additional modifying factor, which is based on a professional judgment of the entire database on the chemical. The RfDs are not applicable to nonthreshold effects such as cancer.

Reportable Quantity (RQ)—The quantity of a hazardous substance that is considered reportable under the Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA). Reportable quantities are (1) 1 pound or greater or (2) for selected substances, an amount established by regulation either under CERCLA or under Section 311 of the Clean Water Act. Quantities are measured over a 24-hour period.

Reproductive Toxicity—The occurrence of adverse effects on the reproductive system that may result from exposure to a chemical. The toxicity may be directed to the reproductive organs and/or the related endocrine system. The manifestation of such toxicity may be noted as alterations in sexual behavior, fertility, pregnancy outcomes, or modifications in other functions that are dependent on the integrity of this system.

Retrospective Study—A type of cohort study based on a group of persons known to have been exposed at some time in the past. Data are collected from routinely recorded events, up to the time the study is undertaken. Retrospective studies are limited to causal factors that can be ascertained from existing records and/or examining survivors of the cohort.

Risk—The possibility or chance that some adverse effect will result from a given exposure to a chemical.

10. GLOSSARY

Risk Factor—An aspect of personal behavior or lifestyle, an environmental exposure, or an inborn or inherited characteristic that is associated with an increased occurrence of disease or other health-related event or condition.

Risk Ratio—The ratio of the risk among persons with specific risk factors compared to the risk among persons without risk factors. A risk ratio greater than 1 indicates greater risk of disease in the exposed group compared to the unexposed.

Short-Term Exposure Limit (STEL)—The American Conference of Governmental Industrial Hygienists (ACGIH) maximum concentration to which workers can be exposed for up to 15 minutes continually. No more than four excursions are allowed per day, and there must be at least 60 minutes between exposure periods. The daily Threshold Limit Value - Time Weighted Average (TLV-TWA) may not be exceeded.

Standardized Mortality Ratio (SMR)—A ratio of the observed number of deaths and the expected number of deaths in a specific standard population.

Target Organ Toxicity—This term covers a broad range of adverse effects on target organs or physiological systems (e.g., renal, cardiovascular) extending from those arising through a single limited exposure to those assumed over a lifetime of exposure to a chemical.

Teratogen—A chemical that causes structural defects that affect the development of an organism.

Threshold Limit Value (TLV)—An American Conference of Governmental Industrial Hygienists (ACGIH) concentration of a substance to which most workers can be exposed without adverse effect. The TLV may be expressed as a Time Weighted Average (TWA), as a Short-Term Exposure Limit (STEL), or as a ceiling limit (CL).

Time-Weighted Average (TWA)—An allowable exposure concentration averaged over a normal 8-hour workday or 40-hour workweek.

Toxic Dose(50) (TD50)—A calculated dose of a chemical, introduced by a route other than inhalation, which is expected to cause a specific toxic effect in 50% of a defined experimental animal population.

Toxicokinetic—The absorption, distribution, and elimination of toxic compounds in the living organism.

Uncertainty Factor (UF)—A factor used in operationally deriving the Minimal Risk Level (MRL) or Reference Dose (RfD) or Reference Concentration (RfC) from experimental data. UFs are intended to account for (1) the variation in sensitivity among the members of the human population, (2) the uncertainty in extrapolating animal data to the case of human, (3) the uncertainty in extrapolating from data obtained in a study that is of less than lifetime exposure, and (4) the uncertainty in using lowest-observed-adverse-effect level (LOAEL) data rather than no-observed-adverse-effect level (NOAEL) data. A default for each individual UF is 10; if complete certainty in data exists, a value of 1 can be used; however, a reduced UF of 3 may be used on a case-by-case basis, 3 being the approximate logarithmic average of 10 and 1.

Xenobiotic—Any chemical that is foreign to the biological system.

APPENDIX A. ATSDR MINIMAL RISK LEVELS AND WORKSHEETS

The Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA) [42 U.S.C. 9601 et seq.], as amended by the Superfund Amendments and Reauthorization Act (SARA) [Pub. L. 99–499], requires that the Agency for Toxic Substances and Disease Registry (ATSDR) develop jointly with the U.S. Environmental Protection Agency (EPA), in order of priority, a list of hazardous substances most commonly found at facilities on the CERCLA National Priorities List (NPL); prepare toxicological profiles for each substance included on the priority list of hazardous substances; and assure the initiation of a research program to fill identified data needs associated with the substances.

The toxicological profiles include an examination, summary, and interpretation of available toxicological information and epidemiologic evaluations of a hazardous substance. During the development of toxicological profiles, Minimal Risk Levels (MRLs) are derived when reliable and sufficient data exist to identify the target organ(s) of effect or the most sensitive health effect(s) for a specific duration for a given route of exposure. An MRL is an estimate of the daily human exposure to a hazardous substance that is likely to be without appreciable risk of adverse noncancer health effects over a specified duration of exposure. MRLs are based on noncancer health effects only and are not based on a consideration of cancer effects. These substance-specific estimates, which are intended to serve as screening levels, are used by ATSDR health assessors to identify contaminants and potential health effects that may be of concern at hazardous waste sites. It is important to note that MRLs are not intended to define clean-up or action levels.

MRLs are derived for hazardous substances using the no-observed-adverse-effect level/uncertainty factor approach. They are below levels that might cause adverse health effects in the people most sensitive to such chemical-induced effects. MRLs are derived for acute (1–14 days), intermediate (15–364 days), and chronic (365 days and longer) durations and for the oral and inhalation routes of exposure. Currently, MRLs for the dermal route of exposure are not derived because ATSDR has not yet identified a method suitable for this route of exposure. MRLs are generally based on the most sensitive chemical-induced end point considered to be of relevance to humans. Serious health effects (such as irreparable damage to the liver or kidneys, or birth defects) are not used as a basis for establishing MRLs. Exposure to a level above the MRL does not mean that adverse health effects will occur.

MRLs are intended only to serve as a screening tool to help public health professionals decide where to look more closely. They may also be viewed as a mechanism to identify those hazardous waste sites that

APPENDIX A

are not expected to cause adverse health effects. Most MRLs contain a degree of uncertainty because of the lack of precise toxicological information on the people who might be most sensitive (e.g., infants, elderly, nutritionally or immunologically compromised) to the effects of hazardous substances. ATSDR uses a conservative (i.e., protective) approach to address this uncertainty consistent with the public health principle of prevention. Although human data are preferred, MRLs often must be based on animal studies because relevant human studies are lacking. In the absence of evidence to the contrary, ATSDR assumes that humans are more sensitive to the effects of hazardous substance than animals and that certain persons may be particularly sensitive. Thus, the resulting MRL may be as much as 100-fold below levels that have been shown to be nontoxic in laboratory animals.

Proposed MRLs undergo a rigorous review process: Health Effects/MRL Workgroup reviews within the Division of Toxicology, expert panel peer reviews, and agency-wide MRL Workgroup reviews, with participation from other federal agencies and comments from the public. They are subject to change as new information becomes available concomitant with updating the toxicological profiles. Thus, MRLs in the most recent toxicological profiles supersede previously published levels. For additional information regarding MRLs, please contact the Division of Toxicology, Agency for Toxic Substances and Disease Registry, 1600 Clifton Road NE, Mailstop F-32, Atlanta, Georgia 30333.

APPENDIX A

MINIMAL RISK LEVEL (MRL) WORKSHEET

Chemical Name: Copper and Compounds
CAS Number:
Date: September 8, 2004
Profile Status: Final Post-Public Comment
Route: Inhalation Oral
Duration: Acute Intermediate Chronic
Key to Figure: 11
Species: Humans

Minimal Risk Level: 0.01 mg copper/kg/day ppm

Reference: Pizarro F, Olivares M, Uauy R, et al. 1999. Acute gastrointestinal effects of graded levels of copper in drinking water. Environ Health Perspect 107:117-121.

Experimental design: (human study details or strain, number of animals per exposure/control groups, sex, dose administration details): A group of 60 healthy women (mean ages of 32.9–36.3 years) were divided into four groups. Each group consumed water containing 0, 1, 3, or 5 mg/L ionic copper as copper sulfate (0.006, 0.0272, 0.0731, and 0.124 mg Cu/kg/day) for a 2-week period with a 1-week rest between copper exposures. Each week, the subjects received a bottle containing copper sulfate solution and were asked to mix the contents of the bottle with 3 L water; this water was then used for drinking and cooking. The subjects recorded daily water consumption and any symptoms. Blood samples were collected 1 week before the study, at the end of the first 2-week exposure period, and at the end of the study; the blood was analyzed for serum copper, aspartate aminotransferase, alanine aminotransferase, and gamma glutamyl transferase activities, and hemoglobin levels. The average copper dietary intake, based on a 24-hour dietary recall, was 1.7 mg Cu/day (0.0266 mg Cu/kg/day using an average body weight of 64 kg).

Effects noted in study and corresponding doses: No significant alterations in serum copper, ceruloplasmin, hemoglobin, or liver enzymes were observed. Twenty-one subjects reported gastrointestinal symptoms, predominantly nausea. Nine subjects reported diarrhea with or without abdominal pain, no association between copper level and diarrhea was found. Six of these episodes of diarrhea occurred during the first week of the study independent of copper concentration. Twelve subjects reported abdominal pain, nausea, and/or vomiting; the incidences were 3/60, 1/60, 10/60, and 9/60 in the control, 0.0272, 0.0731, and 0.124 mg Cu/kg/day groups, respectively. There was a significant difference between in the incidences at concentrations of ≤ 1 mg/L (0.0272 mg/kg/day) versus ≥ 3 mg/L (0.0731 mg/kg/day). No other differences between groups were found.

Dose and end point used for MRL derivation: The MRL is based on the NOAEL of 0.0272 mg Cu/kg/day for gastrointestinal effects in women ingesting copper sulfate in drinking water for 2 weeks (Pizarro et al. 1999).

NOAEL LOAEL

Uncertainty Factors used in MRL derivation:

10 for use of a extrapolation from animals to humans
 3 for human variability; a partial uncertainty factor was used because toxicokinetic differences among individuals should not affect the sensitivity of this direct contact effect.

APPENDIX A

Was a conversion factor used from ppm in food or water to a mg/body weight dose? Yes. Daily doses of copper from drinking water were calculated using reported daily copper intakes (0.04, 1.74, 4.68, and 7.94 mg) and the average of the mean reported body weights (64 kg).

If an inhalation study in animals, list conversion factors used in determining human equivalent dose: NA

Was a conversion used from intermittent to continuous exposure? No

Other additional studies or pertinent information that lend support to this MRL: Numerous experimental studies and case reports support the identification of the gastrointestinal tract as the most sensitive end point of toxicity in humans acutely exposed to copper in drinking water or in contaminated beverages (Araya et al. 2001, 2003a, 2003b, 2003c; Chuttani et al. 1965; Gotteland et al. 2001; Knobeloch et al. 1994; Nicholas and Brist 1968; Olivares et al. 2001; Pizarro et al. 1999, 2001; Spitalny et al. 1984). In single exposure experiments in which subjects ingested copper sulfate in drinking water following an overnight fast, nausea, vomiting, and/or abdominal pain were reported at doses ranging from 0.011 to 0.03 mg Cu/kg/day as copper sulfate (Araya et al. 2001, 2003a, 2003c; Gotteland et al. 2001; Olivares et al. 2001). NOAEL values identified in these studies ranged from 0.0057 to 0.011 mg Cu/kg. Daily exposure to 0.096 mg Cu/kg/day as copper sulfate or copper oxide for 1 week also resulted in an increased occurrence of nausea, vomiting, and/or abdominal pain (Pizarro et al. 2001). Animal studies support the identification of the gastrointestinal tract as the most sensitive target of toxicity following acute-duration oral exposure. Hyperplasia of the forestomach mucosa was observed in rats exposed to 44 mg Cu/kg/day as copper sulfate in the diet (NTP 1993) and in mice exposed to 197 mg Cu/kg/day as copper sulfate in the diet (NTP 1993). At higher doses, liver and kidney damage have been observed (Haywood 1980; Haywood and Comerford 1980; Haywood et al. 1985b; NTP 1993).

Although the LOAEL values identified in the single exposure studies are lower than the NOAEL identified in the Pizarro et al. (1999) study, the Pizarro et al. (1999) study was selected as the critical study because it is a longer-duration study and it more closely mimics an exposure scenario of a population drinking copper-contaminated drinking water.

Agency Contact (Chemical Manager): Alfred Dorsey, DVM

APPENDIX A

MINIMAL RISK LEVEL (MRL) WORKSHEET

Chemical Name: Copper and Compounds
CAS Number:
Date: September 8, 2004
Profile Status: Final Post-Public Comment
Route: [] Inhalation [X] Oral
Duration: [] Acute [X] Intermediate [] Chronic
Key to Figure: 21
Species: Humans

Minimal Risk Level: 0.01 [X] mg copper/kg/day [] ppm

Reference: Araya M, Olivares M, Pizarro F, et al. 2003b. Gastrointestinal symptoms and blood indicators of copper load in apparently healthy adults undergoing controlled copper exposure. *Am J Clin Nutr* 77:646-650.

Experimental design (human study details or strain, number of animals per exposure/control groups, sex, dose administration details): Groups of 327–340 men and women (mean age 32.9 years) were exposed to 0, 2, 4, or 6 mg Cu/L in drinking water for 2 months. The subjects prepared the copper sulfate solution daily using tap water and a stock copper sulfate solution. The copper solution was used for drinking and preparation of beverages and soups. The subjects completed a daily survey on gastrointestinal and other symptoms. Blood samples were analyzed for a subset of 48–49 subjects for red blood cell copper, monocyte copper, serum copper, serum ceruloplasmin, superoxide dismutase, aspartate aminotransferase, alanine amino transferases, γ -glutamyltransferase, and hemoglobin levels. Reported copper intakes from water in the subset of subjects were 0, 42.5, 92.9, and 177.9 $\mu\text{mol/day}$ (0, 2.7, 5.9, and 11.3 mg/day).

Doses were calculated using reported copper intakes for the subset of subjects and a reference body weight of 65 kg. 2 ppm: $2.7 \text{ mg Cu/day} \times 1/65 \text{ kg} = 0.042 \text{ mg Cu/kg/day}$; 4 ppm: $5.9 \text{ mg/day} = 0.091 \text{ mg Cu/kg/day}$; 6 ppm: $11.3 \text{ mg/day} = 0.17 \text{ mg Cu/kg/day}$

Effects noted in study and corresponding doses: The incidences of gastrointestinal symptoms were 11.7, 15.3, 18.3, and 19.7% in the 0, 0.042, 0.091, and 0.17 mg Cu/kg/day groups, respectively. Using a chi-square test with Bonferroni correction, the incidence of gastrointestinal symptoms was significantly increased in the 0.17 mg Cu/kg/day group. However, if the Bonferroni correction was not used, the incidence was also significantly increased in the 0.091 mg Cu/kg/day group. Only one test is used to assess whether exposure to copper results in adverse gastrointestinal effects (reported symptoms); thus, the Bonferroni correction is not needed for this end point. No significant alterations in copper status parameters or biomarkers of liver disease were observed.

Dose and end point used for MRL derivation: The MRL is based on the NOAEL of 0.042 mg Cu/kg/day for gastrointestinal effects in men and women ingesting copper sulfate in drinking water for 2 months.

[X] NOAEL [] LOAEL

APPENDIX A

Uncertainty Factors used in MRL derivation:

[] 10 for use of a extrapolation from animals to humans

[X] 3 for human variability; a partial uncertainty factor was used because toxicokinetic differences among individuals should not affect the sensitivity of this direct contact effect

Was a conversion factor used from ppm in food or water to a mg/body weight dose? Yes. Daily doses were calculated using reported daily copper intakes (2.7, 5.9, and 11.3 mg) and a reference body weight of 65 kg.

If an inhalation study in animals, list conversion factors used in determining human equivalent dose: NA

Was a conversion used from intermittent to continuous exposure? No

Other additional studies or pertinent information that lend support to this MRL: There are limited data on the intermediate-duration toxicity of copper in humans. The Araya et al. (2003b) is the only human study located that examined the gastrointestinal tract following intermediate-duration exposure to copper. A number of acute-duration studies (Araya et al. 2001, 2003a, 2003c; Gotteland et al. 2001; Nicholas and Brist 1968; Olivares et al. 2001) support the identification of the gastrointestinal tract as the sensitive target of toxicity in humans. An intermediate-duration study by Pratt et al. (1985) did not find alterations in serum biomarkers of liver damage (cholesterol and triglyceride levels and aspartate aminotransferase, alkaline phosphatase, γ -glutamyl transferase, and lactate dehydrogenase activities) in seven adults administered 10 mg Cu/day (0.14 mg Cu/kg/day) as copper gluconate in a capsule for 12 weeks. Three intermediate-duration studies in infants also found no significant evidence of liver damage (Olivares et al. 1998; Zietz et al. 2003a, 2003b).

Numerous animal studies have examined the toxicity of copper following intermediate-duration oral exposure (Epstein et al. 1982; Fuentealba et al. 2000; Haywood 1980, 1985; Haywood and Comerford 1980; Haywood and Loughran 1985; Haywood et al. 1985a, 1985b; Kumar and Sharma 1987; NTP 1993; Rana and Kumar 1980). Most of these studies have focused on the liver and kidneys, with adverse effect levels of at least 100 times higher than the adverse effect level for gastrointestinal effects in humans. Gastrointestinal tract alterations were observed in rats and mice. Hyperplasia of the squamous mucosa on the limiting ridge separating the forestomach from the glandular stomach was observed in rats and mice exposed to 33 or 267 mg Cu/kg/day, respectively, as copper sulfate in the diet for 13 weeks (NTP 1993).

Agency Contact (Chemical Manager): Alfred Dorsey, DVM

APPENDIX B. USER'S GUIDE

Chapter 1

Public Health Statement

This chapter of the profile is a health effects summary written in non-technical language. Its intended audience is the general public, especially people living in the vicinity of a hazardous waste site or chemical release. If the Public Health Statement were removed from the rest of the document, it would still communicate to the lay public essential information about the chemical.

The major headings in the Public Health Statement are useful to find specific topics of concern. The topics are written in a question and answer format. The answer to each question includes a sentence that will direct the reader to chapters in the profile that will provide more information on the given topic.

Chapter 2

Relevance to Public Health

This chapter provides a health effects summary based on evaluations of existing toxicologic, epidemiologic, and toxicokinetic information. This summary is designed to present interpretive, weight-of-evidence discussions for human health end points by addressing the following questions:

1. What effects are known to occur in humans?
2. What effects observed in animals are likely to be of concern to humans?
3. What exposure conditions are likely to be of concern to humans, especially around hazardous waste sites?

The chapter covers end points in the same order that they appear within the Discussion of Health Effects by Route of Exposure section, by route (inhalation, oral, and dermal) and within route by effect. Human data are presented first, then animal data. Both are organized by duration (acute, intermediate, chronic). *In vitro* data and data from parenteral routes (intramuscular, intravenous, subcutaneous, etc.) are also considered in this chapter.

The carcinogenic potential of the profiled substance is qualitatively evaluated, when appropriate, using existing toxicokinetic, genotoxic, and carcinogenic data. ATSDR does not currently assess cancer potency or perform cancer risk assessments. Minimal Risk Levels (MRLs) for noncancer end points (if derived) and the end points from which they were derived are indicated and discussed.

Limitations to existing scientific literature that prevent a satisfactory evaluation of the relevance to public health are identified in the Chapter 3 Data Needs section.

Interpretation of Minimal Risk Levels

Where sufficient toxicologic information is available, ATSDR has derived MRLs for inhalation and oral routes of entry at each duration of exposure (acute, intermediate, and chronic). These MRLs are not

APPENDIX B

meant to support regulatory action, but to acquaint health professionals with exposure levels at which adverse health effects are not expected to occur in humans.

MRLs should help physicians and public health officials determine the safety of a community living near a chemical emission, given the concentration of a contaminant in air or the estimated daily dose in water. MRLs are based largely on toxicological studies in animals and on reports of human occupational exposure.

MRL users should be familiar with the toxicologic information on which the number is based. Chapter 2, "Relevance to Public Health," contains basic information known about the substance. Other sections such as Chapter 3 Section 3.9, "Interactions with Other Substances," and Section 3.10, "Populations that are Unusually Susceptible" provide important supplemental information.

MRL users should also understand the MRL derivation methodology. MRLs are derived using a modified version of the risk assessment methodology that the Environmental Protection Agency (EPA) provides (Barnes and Dourson 1988) to determine reference doses (RfDs) for lifetime exposure.

To derive an MRL, ATSDR generally selects the most sensitive end point which, in its best judgement, represents the most sensitive human health effect for a given exposure route and duration. ATSDR cannot make this judgement or derive an MRL unless information (quantitative or qualitative) is available for all potential systemic, neurological, and developmental effects. If this information and reliable quantitative data on the chosen end point are available, ATSDR derives an MRL using the most sensitive species (when information from multiple species is available) with the highest no-observed-adverse-effect level (NOAEL) that does not exceed any adverse effect levels. When a NOAEL is not available, a lowest-observed-adverse-effect level (LOAEL) can be used to derive an MRL, and an uncertainty factor (UF) of 10 must be employed. Additional uncertainty factors of 10 must be used both for human variability to protect sensitive subpopulations (people who are most susceptible to the health effects caused by the substance) and for interspecies variability (extrapolation from animals to humans). In deriving an MRL, these individual uncertainty factors are multiplied together. The product is then divided into the inhalation concentration or oral dosage selected from the study. Uncertainty factors used in developing a substance-specific MRL are provided in the footnotes of the levels of significant exposure (LSE) tables.

Chapter 3

Health Effects

Tables and Figures for Levels of Significant Exposure (LSE)

Tables and figures are used to summarize health effects and illustrate graphically levels of exposure associated with those effects. These levels cover health effects observed at increasing dose concentrations and durations, differences in response by species, MRLs to humans for noncancer end points, and EPA's estimated range associated with an upper-bound individual lifetime cancer risk of 1 in 10,000 to 1 in 10,000,000. Use the LSE tables and figures for a quick review of the health effects and to locate data for a specific exposure scenario. The LSE tables and figures should always be used in conjunction with the text. All entries in these tables and figures represent studies that provide reliable, quantitative estimates of NOAELs, LOAELs, or Cancer Effect Levels (CELs).

APPENDIX B

The legends presented below demonstrate the application of these tables and figures. Representative examples of LSE Table 3-1 and Figure 3-1 are shown. The numbers in the left column of the legends correspond to the numbers in the example table and figure.

LEGEND**See Sample LSE Table 3-1 (page B-6)**

- (1) Route of Exposure. One of the first considerations when reviewing the toxicity of a substance using these tables and figures should be the relevant and appropriate route of exposure. Typically when sufficient data exist, three LSE tables and two LSE figures are presented in the document. The three LSE tables present data on the three principal routes of exposure, i.e., inhalation, oral, and dermal (LSE Tables 3-1, 3-2, and 3-3, respectively). LSE figures are limited to the inhalation (LSE Figure 3-1) and oral (LSE Figure 3-2) routes. Not all substances will have data on each route of exposure and will not, therefore, have all five of the tables and figures.
- (2) Exposure Period. Three exposure periods—acute (less than 15 days), intermediate (15–364 days), and chronic (365 days or more)—are presented within each relevant route of exposure. In this example, an inhalation study of intermediate exposure duration is reported. For quick reference to health effects occurring from a known length of exposure, locate the applicable exposure period within the LSE table and figure.
- (3) Health Effect. The major categories of health effects included in LSE tables and figures are death, systemic, immunological, neurological, developmental, reproductive, and cancer. NOAELs and LOAELs can be reported in the tables and figures for all effects but cancer. Systemic effects are further defined in the "System" column of the LSE table (see key number 18).
- (4) Key to Figure. Each key number in the LSE table links study information to one or more data points using the same key number in the corresponding LSE figure. In this example, the study represented by key number 18 has been used to derive a NOAEL and a Less Serious LOAEL (also see the two "18r" data points in sample Figure 3-1).
- (5) Species. The test species, whether animal or human, are identified in this column. Chapter 2, "Relevance to Public Health," covers the relevance of animal data to human toxicity and Section 3.4, "Toxicokinetics," contains any available information on comparative toxicokinetics. Although NOAELs and LOAELs are species specific, the levels are extrapolated to equivalent human doses to derive an MRL.
- (6) Exposure Frequency/Duration. The duration of the study and the weekly and daily exposure regimens are provided in this column. This permits comparison of NOAELs and LOAELs from different studies. In this case (key number 18), rats were exposed to "Chemical x" via inhalation for 6 hours/day, 5 days/week, for 13 weeks. For a more complete review of the dosing regimen, refer to the appropriate sections of the text or the original reference paper (i.e., Nitschke et al. 1981).
- (7) System. This column further defines the systemic effects. These systems include respiratory, cardiovascular, gastrointestinal, hematological, musculoskeletal, hepatic, renal, and dermal/ocular. "Other" refers to any systemic effect (e.g., a decrease in body weight) not covered

APPENDIX B

in these systems. In the example of key number 18, one systemic effect (respiratory) was investigated.

- (8) NOAEL. A NOAEL is the highest exposure level at which no harmful effects were seen in the organ system studied. Key number 18 reports a NOAEL of 3 ppm for the respiratory system, which was used to derive an intermediate exposure, inhalation MRL of 0.005 ppm (see footnote "b").
- (9) LOAEL. A LOAEL is the lowest dose used in the study that caused a harmful health effect. LOAELs have been classified into "Less Serious" and "Serious" effects. These distinctions help readers identify the levels of exposure at which adverse health effects first appear and the gradation of effects with increasing dose. A brief description of the specific end point used to quantify the adverse effect accompanies the LOAEL. The respiratory effect reported in key number 18 (hyperplasia) is a Less Serious LOAEL of 10 ppm. MRLs are not derived from Serious LOAELs.
- (10) Reference. The complete reference citation is given in Chapter 9 of the profile.
- (11) CEL. A CEL is the lowest exposure level associated with the onset of carcinogenesis in experimental or epidemiologic studies. CELs are always considered serious effects. The LSE tables and figures do not contain NOAELs for cancer, but the text may report doses not causing measurable cancer increases.
- (12) Footnotes. Explanations of abbreviations or reference notes for data in the LSE tables are found in the footnotes. Footnote "b" indicates that the NOAEL of 3 ppm in key number 18 was used to derive an MRL of 0.005 ppm.

LEGEND

See Sample Figure 3-1 (page B-7)

LSE figures graphically illustrate the data presented in the corresponding LSE tables. Figures help the reader quickly compare health effects according to exposure concentrations for particular exposure periods.

- (13) Exposure Period. The same exposure periods appear as in the LSE table. In this example, health effects observed within the acute and intermediate exposure periods are illustrated.
- (14) Health Effect. These are the categories of health effects for which reliable quantitative data exists. The same health effects appear in the LSE table.
- (15) Levels of Exposure. Concentrations or doses for each health effect in the LSE tables are graphically displayed in the LSE figures. Exposure concentration or dose is measured on the log scale "y" axis. Inhalation exposure is reported in mg/m³ or ppm and oral exposure is reported in mg/kg/day.
- (16) NOAEL. In this example, the open circle designated 18r identifies a NOAEL critical end point in the rat upon which an intermediate inhalation exposure MRL is based. The key number 18 corresponds to the entry in the LSE table. The dashed descending arrow indicates the

APPENDIX B

extrapolation from the exposure level of 3 ppm (see entry 18 in the table) to the MRL of 0.005 ppm (see footnote "b" in the LSE table).

- (17) CEL. Key number 38m is one of three studies for which CELs were derived. The diamond symbol refers to a CEL for the test species-mouse. The number 38 corresponds to the entry in the LSE table.
- (18) Estimated Upper-Bound Human Cancer Risk Levels. This is the range associated with the upper-bound for lifetime cancer risk of 1 in 10,000 to 1 in 10,000,000. These risk levels are derived from the EPA's Human Health Assessment Group's upper-bound estimates of the slope of the cancer dose response curve at low dose levels (q_1^*).
- (19) Key to LSE Figure. The Key explains the abbreviations and symbols used in the figure.

SAMPLE


Table 3-1. Levels of Significant Exposure to [Chemical x] – Inhalation

Key to figure ^a	Species	Exposure frequency/ duration	System	NOAEL (ppm)	LOAEL (effect)		Reference	
					Less serious (ppm)	Serious (ppm)		
INTERMEDIATE EXPOSURE								
3	Systemic	↓	↓	↓	↓	↓	↓	
4	18	Rat	13 wk 5 d/wk 6 hr/d	Resp	3 ^b	10 (hyperplasia)	Nitschke et al. 1981	
CHRONIC EXPOSURE								
Cancer								
						11		
						↓		
	38	Rat	18 mo 5 d/wk 7 hr/d			20	(CEL, multiple organs)	Wong et al. 1982
	39	Rat	89–104 wk 5 d/wk 6 hr/d			10	(CEL, lung tumors, nasal tumors)	NTP 1982
	40	Mouse	79–103 wk 5 d/wk 6 hr/d			10	(CEL, lung tumors, hemangiosarcomas)	NTP 1982

^a The number corresponds to entries in Figure 3-1.
^b Used to derive an intermediate inhalation Minimal Risk Level (MRL) of 5×10^{-3} ppm; dose adjusted for intermittent exposure and divided by an uncertainty factor of 100 (10 for extrapolation from animal to humans, 10 for human variability).

SAMPLE

Figure 3-1. Levels of Significant Exposure to [Chemical X] - Inhalation



APPENDIX B

APPENDIX C. ACRONYMS, ABBREVIATIONS, AND SYMBOLS

ACGIH	American Conference of Governmental Industrial Hygienists
ACOEM	American College of Occupational and Environmental Medicine
ADI	acceptable daily intake
ADME	absorption, distribution, metabolism, and excretion
AED	atomic emission detection
AFID	alkali flame ionization detector
AFOSH	Air Force Office of Safety and Health
ALT	alanine aminotransferase
AML	acute myeloid leukemia
AOAC	Association of Official Analytical Chemists
AOEC	Association of Occupational and Environmental Clinics
AP	alkaline phosphatase
APHA	American Public Health Association
AST	aspartate aminotransferase
atm	atmosphere
ATSDR	Agency for Toxic Substances and Disease Registry
AWQC	Ambient Water Quality Criteria
BAT	best available technology
BCF	bioconcentration factor
BEI	Biological Exposure Index
BMD	benchmark dose
BMR	benchmark response
BSC	Board of Scientific Counselors
C	centigrade
CAA	Clean Air Act
CAG	Cancer Assessment Group of the U.S. Environmental Protection Agency
CAS	Chemical Abstract Services
CDC	Centers for Disease Control and Prevention
CEL	cancer effect level
CELDS	Computer-Environmental Legislative Data System
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act
CFR	Code of Federal Regulations
Ci	curie
CI	confidence interval
CL	ceiling limit value
CLP	Contract Laboratory Program
cm	centimeter
CML	chronic myeloid leukemia
CPSC	Consumer Products Safety Commission
CWA	Clean Water Act
DHEW	Department of Health, Education, and Welfare
DHHS	Department of Health and Human Services
DNA	deoxyribonucleic acid
DOD	Department of Defense
DOE	Department of Energy
DOL	Department of Labor
DOT	Department of Transportation

APPENDIX C

DOT/UN/ NA/IMCO	Department of Transportation/United Nations/ North America/International Maritime Dangerous Goods Code
DWEL	drinking water exposure level
ECD	electron capture detection
ECG/EKG	electrocardiogram
EEG	electroencephalogram
EEGL	Emergency Exposure Guidance Level
EPA	Environmental Protection Agency
F	Fahrenheit
F ₁	first-filial generation
FAO	Food and Agricultural Organization of the United Nations
FDA	Food and Drug Administration
FEMA	Federal Emergency Management Agency
FIFRA	Federal Insecticide, Fungicide, and Rodenticide Act
FPD	flame photometric detection
fpm	feet per minute
FR	Federal Register
FSH	follicle stimulating hormone
g	gram
GC	gas chromatography
gd	gestational day
GLC	gas liquid chromatography
GPC	gel permeation chromatography
HPLC	high-performance liquid chromatography
HRGC	high resolution gas chromatography
HSDB	Hazardous Substance Data Bank
IARC	International Agency for Research on Cancer
IDLH	immediately dangerous to life and health
ILO	International Labor Organization
IRIS	Integrated Risk Information System
K _d	adsorption ratio
kg	kilogram
kkg	metric ton
K _{oc}	organic carbon partition coefficient
K _{ow}	octanol-water partition coefficient
L	liter
LC	liquid chromatography
LC ₅₀	lethal concentration, 50% kill
LC _{Lo}	lethal concentration, low
LD ₅₀	lethal dose, 50% kill
LD _{Lo}	lethal dose, low
LDH	lactic dehydrogenase
LH	lutinizing hormone
LOAEL	lowest-observed-adverse-effect level
LSE	Levels of Significant Exposure
LT ₅₀	lethal time, 50% kill
m	meter
MA	<i>trans,trans</i> -muconic acid
MAL	maximum allowable level
mCi	millicurie
MCL	maximum contaminant level

APPENDIX C

MCLG	maximum contaminant level goal
MF	modifying factor
MFO	mixed function oxidase
mg	milligram
mL	milliliter
mm	millimeter
mmHg	millimeters of mercury
mmol	millimole
mppcf	millions of particles per cubic foot
MRL	Minimal Risk Level
MS	mass spectrometry
NAAQS	National Ambient Air Quality Standard
NAS	National Academy of Science
NATICH	National Air Toxics Information Clearinghouse
NATO	North Atlantic Treaty Organization
NCE	normochromatic erythrocytes
NCEH	National Center for Environmental Health
NCI	National Cancer Institute
ND	not detected
NFPA	National Fire Protection Association
ng	nanogram
NHANES	National Health and Nutrition Examination Survey
NIEHS	National Institute of Environmental Health Sciences
NIOSH	National Institute for Occupational Safety and Health
NIOSHTIC	NIOSH's Computerized Information Retrieval System
NLM	National Library of Medicine
nm	nanometer
nmol	nanomole
NOAEL	no-observed-adverse-effect level
NOES	National Occupational Exposure Survey
NOHS	National Occupational Hazard Survey
NPD	nitrogen phosphorus detection
NPDES	National Pollutant Discharge Elimination System
NPL	National Priorities List
NR	not reported
NRC	National Research Council
NS	not specified
NSPS	New Source Performance Standards
NTIS	National Technical Information Service
NTP	National Toxicology Program
ODW	Office of Drinking Water, EPA
OERR	Office of Emergency and Remedial Response, EPA
OHM/TADS	Oil and Hazardous Materials/Technical Assistance Data System
OPP	Office of Pesticide Programs, EPA
OPPT	Office of Pollution Prevention and Toxics, EPA
OPPTS	Office of Prevention, Pesticides and Toxic Substances, EPA
OR	odds ratio
OSHA	Occupational Safety and Health Administration
OSW	Office of Solid Waste, EPA
OTS	Office of Toxic Substances
OW	Office of Water

APPENDIX C

OWRS	Office of Water Regulations and Standards, EPA
PAH	polycyclic aromatic hydrocarbon
PBPD	physiologically based pharmacodynamic
PBPK	physiologically based pharmacokinetic
PCE	polychromatic erythrocytes
PEL	permissible exposure limit
pg	picogram
PHS	Public Health Service
PID	photo ionization detector
pmol	picomole
PMR	proportionate mortality ratio
ppb	parts per billion
ppm	parts per million
ppt	parts per trillion
PSNS	pretreatment standards for new sources
RBC	red blood cell
REL	recommended exposure level/limit
RfC	reference concentration
RfD	reference dose
RNA	ribonucleic acid
RQ	reportable quantity
RTECS	Registry of Toxic Effects of Chemical Substances
SARA	Superfund Amendments and Reauthorization Act
SCE	sister chromatid exchange
SGOT	serum glutamic oxaloacetic transaminase
SGPT	serum glutamic pyruvic transaminase
SIC	standard industrial classification
SIM	selected ion monitoring
SMCL	secondary maximum contaminant level
SMR	standardized mortality ratio
SNARL	suggested no adverse response level
SPEGL	Short-Term Public Emergency Guidance Level
STEL	short term exposure limit
STORET	Storage and Retrieval
TD ₅₀	toxic dose, 50% specific toxic effect
TLV	threshold limit value
TOC	total organic carbon
TPQ	threshold planning quantity
TRI	Toxics Release Inventory
TSCA	Toxic Substances Control Act
TWA	time-weighted average
UF	uncertainty factor
U.S.	United States
USDA	United States Department of Agriculture
USGS	United States Geological Survey
VOC	volatile organic compound
WBC	white blood cell
WHO	World Health Organization

APPENDIX C

>	greater than
≥	greater than or equal to
=	equal to
<	less than
≤	less than or equal to
%	percent
α	alpha
β	beta
γ	gamma
δ	delta
μm	micrometer
μg	microgram
q ₁ *	cancer slope factor
-	negative
+	positive
(+)	weakly positive result
(-)	weakly negative result

APPENDIX D. INDEX

absorbed dose.....	85
adsorbed.....	135, 143, 145, 146, 152, 156, 184
adsorption.....	121, 142, 146, 147, 151, 152, 155, 193, 197
aerobic.....	121, 145
alanine aminotransferase.....	14, 17, 18, 57, 59, 60, 61, 79, 80, 83, 88, 96
ambient air.....	11, 182, 187, 202, 205
anaerobic.....	121, 152, 157
anemia.....	12, 56, 65, 82, 87, 89
aspartate aminotransferase.....	14, 17, 18, 57, 61, 62, 83, 88
bioaccumulation.....	149, 150, 176, 187
bioavailability.....	101, 154, 158, 187, 190
bioconcentration factor.....	149, 150
biomarker.....	85, 100
biomarkers.....	13, 14, 17, 57, 83, 85, 87, 100, 193, 198
body weight effects.....	22, 62, 65
breast milk.....	175, 183
cancer.....	6, 7, 13, 31, 64, 82, 98
carcinogen.....	7, 205
carcinogenic.....	15, 21, 64, 98, 99
carcinogenicity.....	13, 31, 64, 66, 98, 205
carcinoma.....	66
carcinomas.....	64
cardiovascular.....	22, 54, 65, 87
cardiovascular effects.....	54, 66
ceruloplasmin.....	58, 71, 72, 73, 78, 80, 89, 91, 100
chromosomal aberrations.....	66
clearance.....	23, 60
death.....	7, 21, 22, 59, 64, 87
deoxyribonucleic acid.....	69
DNA.....	68, 69, 85, 98
dopamine.....	12, 63, 99
endocrine.....	22, 32, 65, 80
endocrine effects.....	24
fetus.....	1, 81, 84
gastrointestinal distress.....	12, 87
gastrointestinal effects.....	12, 13, 14, 16, 17, 18, 19, 23, 24, 54, 55, 56, 83, 86, 95, 97
general population.....	3, 11, 12, 85, 123, 158, 171, 183, 184, 188
genotoxic.....	21, 98
genotoxicity.....	66, 68, 98
groundwater.....	3, 4, 139, 141, 147, 148, 155, 163, 164, 165, 187, 190, 197
half-life.....	73, 85, 155
hematological effects.....	24, 56, 65
hematopoietic.....	56
hepatic effects.....	14, 16, 24, 57, 61, 89, 96
hydroxyl radical.....	76
immune system.....	13
immunological.....	21, 25
immunological effects.....	25

APPENDIX D

K _{ow}	109
leukopenia	12
metabolic effects	32
metallothionen	12, 71, 84
micronuclei	66, 68
milk	14, 58, 82, 90, 91, 95, 97, 102, 174, 175, 188, 203
musculoskeletal effects	56
neonatal	84
neurobehavioral	81, 99
neutropenia	82
norepinephrine	63
nuclear	78, 138, 154
ocular effects	22, 65
partition coefficients	145, 156
pharmacodynamic	75
pharmacokinetic	74, 75, 76, 77
pharmacokinetics	81
pulmonary fibrosis	12, 22
renal effects	61, 95, 96, 100
retention	71, 166
sequestered	78, 88
solubility	147, 153, 185
spermatozoa	66
toxicokinetic	17, 21, 60, 84
toxicokinetics	101
tremors	89
tumors	64
weanling	71

