



Uploaded to the VFC Website

▶▶▶ 2016 ◀◀◀

This Document has been provided to you courtesy of Veterans-For-Change!

Feel free to pass to any veteran who might be able to use this information!

For thousands more files like this and hundreds of links to useful information, and hundreds of "Frequently Asked Questions, please go to:

[Veterans-For-Change](#)

If Veterans don't help Veterans, who will?

Note:

VFC is not liable for source information in this document, it is merely provided as a courtesy to our members & subscribers.



Tailored new compound can dramatically reduce brain damage resulting from stroke

Published on March 10, 2016 at 5:59 AM

After suffering a stroke, about three-fourths of patients exhibit some disability. The extent of a patient's symptoms depends on the degree and location of brain tissue damage following the stroke event. This week in *ACS Central Science*, researchers show that by using a tailored small molecule to turn off the production of a key neuromodulator in the brain, they can dramatically reduce brain damage in stroke models in rats.

The neuromodulator is the gas hydrogen sulfide (H₂S). Its production is carefully controlled in the brain. After a stroke, levels of H₂S appear to be elevated, leading to brain tissue damage, but the details of how that happens are still a bit of a mystery. So, David B. Berkowitz and coworkers designed a quick way to synthesize molecules they deduced would inhibit the production of H₂S. They showed in vitro that these compounds block an enzyme called CBS from making H₂S by mimicking one of its other products. Peter T. H. Wong and colleagues then tested the compounds in rats. When the new compound was injected an hour after the simulation of a stroke, the authors observed about a 70 percent reduction in the severity of the observed stroke damage. The results were even more striking with pretreatment. The authors conclude that using molecules like the ones they made will help researchers dissect the mechanism underlying H₂S-mediated neuronal damage and will serve as an important starting point for the development of even more drug-like compounds that act in a similar manner.

Source:

American Chemical Society
