



---

## Uploaded to the VFC Website

▶▶▶▶ 2022 ◀◀◀◀

---

This Document has been provided to you, courtesy of Veterans-For-Change!

Feel free to pass to any Veteran who might be able to use this information!

For thousands more files like this and hundreds of links to useful information, and hundreds of "Frequently Asked Questions, please go to:

[Veterans-For-Change](#)

---

***If Veterans don't help Veterans, who will?***

---

**Note:**

VFC is not liable for source information in this document, it is merely provided as a courtesy to our members & subscribers.



# Bipolar disorder may involve accelerated epigenetic aging, study finds

Dec 12 2017

Bipolar disorder may involve accelerated epigenetic aging, which could explain why persons with the disorder are more likely to have - and die from - age-related diseases, according to researchers from The University of Texas Health Science Center at Houston (UTHealth).

The findings were published in yesterday's issue of *Translational Psychiatry*, a Nature Publishing Group journal.

While chronological age is measured in the amount of time that a person has been alive, epigenetic age measures molecular markers of chemical modifications to DNA.

"Bipolar disorder has been previously associated with accelerated aging but the mechanisms are largely unknown," said Gabriel R. Fries, Ph.D., first author and post-doctoral research fellow in the Department of Psychiatry and Behavioral Sciences at McGovern Medical School at UTHealth. "We aimed to understand from our study the biology of what's driving the accelerated aging. What we found is that patients with bipolar disorder showed an accelerated epigenetic aging compared to healthy controls."

The chemical modifications could be precipitated by the disorder itself or by poor lifestyle habits in diet, exercise, tobacco use and illegal substance use.

"Controlling these factors is just as important as taking medications," Fries said.

Senior author of the study was Joao L. de Quevedo, M.D., Ph.D., professor and director of the *Translational Psychiatry* Program in the Department of Psychiatry and Behavioral Sciences at McGovern Medical School.

Using blood samples, the researchers compared 22 patients with bipolar disorder, 16 siblings of bipolar patients and 20 healthy controls. They also found that while older bipolar disorder patients had significantly accelerated epigenetic aging compared to controls, no difference was found in younger

patients.

"We believe a difference wasn't detected in younger patients because they haven't had as much exposure to stressful events," Fries said. "This gave us a hint that cumulative chronic exposure to stress would relate to accelerated aging. We would see it more in older people who have experienced a lifetime of stress in dealing with the disease."

Along with the epigenetic clock, the study included two other biologic clocks: telomere length and mitochondrial DNA copy numbers.

"The epigenetic acceleration correlated with the number of copies of mitochondrial DNA, suggesting that the cross-talk between the nucleus and the mitochondria might be underlying the premature aging in bipolar disorder," Fries said.

---

**Source:**

<https://www.uth.edu/media/story.htm?id=805e6f7f-3a33-4806-ba2f-505d1ba8e949>

---